

Research Article

Spatial Variation in Bird Community Composition in Relation to Topographic Gradient and Forest Heterogeneity in a Central Amazonian Rainforest

Renato Cintra¹ and Luciano N. Naka^{1,2}

¹ Coordenação de Biodiversidade, Instituto Nacional de Pesquisas da Amazônia (INPA), 69067-375 Manaus, AM, Brazil

² Centro de Estudos da Biodiversidade, Universidade Federal de Roraima (UFRR), Boa Vista, RR, Brazil

Correspondence should be addressed to Renato Cintra, rcintrasoares@gmail.com

Received 11 March 2011; Revised 16 May 2011; Accepted 8 July 2011

Academic Editor: Bradford Hawkins

Copyright © 2012 R. Cintra and L. N. Naka. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigated the effects of landscape features and forest structure on the avian community at the Reserva Florestal Adolpho Ducke near Manaus, in the Brazilian Amazon. We sampled the landscape and forest in 72 50 × 50 m plots systematically distributed in the reserve, covering an area of 6,400 ha. The avifauna was sampled using mist nets and acoustic surveys near the plots. We found no significant relationships between landscape features and forest components in the plots and the number of bird species and individuals sampled. Results of Principal Coordinate Analyses, however, showed that bird species composition changes along a topographic gradient (plateau-slope-valley), and also in relation to leaf litter depth and distance to forest streams. We also found compositional differences in the avian community on the eastern and western water basins that compose the reserve. Our results suggest that although most bird species occur throughout the reserve, many species track differences in the landscape and the forest structure.

1. Introduction

Understanding how the structure of the habitat influences avian populations and communities is a key factor to link habitat and niche selection with species diversity [1, 2]. Although the mechanisms that determine habitat use in tropical forests remain poorly understood [3, 4], it is clear that different species use their surrounding environment differently. This use can change over space and time due to individual movements, but also as a result of differences in habitat structure [5–9]. Several studies have focused on the factors affecting avian species richness and abundance in tropical forests, investigating the relationship between vegetation structure and bird communities [2, 8–10], but few have investigated how natural variations in the structure of the forest may affect entire species assemblages (but see [4, 7, 11]). When available, such studies are often done at small spatial scales [9, 12, 13], and the effects of habitat structure are rarely considered.

Most bird species in Amazonia are resident and sedentary [12, 14, 15], and are relatively long-lived compared to those in temperate regions [16]. Longevity can result in a more specialized use of the environment, and birds may adjust their microhabitat use according to local variation in the forest structure [7, 11, 17]. Avian community composition, microhabitat selection, and guild assemblages can be affected by landscape features such as elevation, topography (valleys, slopes, and plateaus), and proximity to water. Furthermore, the natural heterogeneity found in many components of the forest structure, such as the number of trees, the amount of light that reaches the forest understory (canopy openness), and even the amount of litter on the ground [4, 7, 11] can also affect avian communities. For example, it has been documented that the number of tree-fall gaps in a forest increases with selective logging, affecting the abundance of insectivorous and nectarivorous birds [13, 18]. Similarly, the spatial variation in the depth of forest leaf litter can affect the use of the forest floor by insectivorous birds [10]. Although

in recent years a great body of knowledge has been gathered on general aspects of the biology and behaviour of many tropical species [19], little is known about the function of species in the ecosystem, and how a significant part of the bird community responds to the local variation observed in the structure of the forest.

In this study, we investigate how the natural forest heterogeneity influences avian communities. We selected seven variables, which include local topography (elevation), number of trees, fallen logs and standing dead trees (snags), depth of leaf litter, percentage of canopy openness, and the distance to the nearest stream. These variables may be important because (i) many bird species rely on a shady understory for foraging, and areas with more trees can offer a more complex structure, sustaining more species [1]; (ii) the abundance of logs and snags may provide more area for foraging and nesting, while providing a particular substrate for specialist species; (iii) the abundance of small vertebrates and large invertebrates (prime avian food resources) are directly related to the abundance of leaf litter [20]; (iv) the amount of light reaching the understory may affect fruit production, attracting arthropods and increasing food availability [17]; (v) streams are an important component of the forest, serving as a microhabitat for several avian specialists; (vi) elevation in central Amazonia is known to be directly related to soil texture (one of the best predictors of forest structure and tree composition) [21].

We tested the effect of these variables on bird species richness, abundance, and species' composition, using data based on mist-net captures and acoustic transect surveys. Specifically, we tested the hypothesis that the spatial variation in the forest structure and some landscape features should influence the occurrence, abundance, and distribution of bird species in a tropical forest. Because within and among-year seasonality can also determine the spatial distribution of bird species richness in a tropical forest [8, 12], we also tested the effect of seasonality on the avifauna.

In particular, we ask the following questions: (1) does the bird community composition change seasonally (dry versus wet season) and among years in our study area?; (2) what are the effects of some selected forest structural components on the richness, abundance, and species' composition of birds?; (3) does bird species composition change along a topographic gradient that includes plateaus, slopes, and valleys?; (4) what, if any, are the differences in bird species composition between the two main drainage basins in the study area?

2. Material and Methods

2.1. Study Area and Sampling Design. The Reserva Florestal Adolpho Ducke (RFAD), a 10,000 ha forest reserve, is located 30 km north of the city of Manaus (02°55'–03°01' S, 59°53'–59°59' W), in the Brazilian state of Amazonas. Although the spread of Manaus, the largest city in the Amazon basin, has reached the southern and western boundaries of the reserve in recent years, RFAD is still connected to continuous forest on its eastern side. The reserve can be divided into an eastern and a western water basin (with six microbasins: Acará, Barro

Branco, Bolívia, Ipiranga, Tinga, and Ubere) divided by a central plateau. The dominant vegetation type in the reserve is primary Terra Firme forest, never inundated by seasonal river fluctuations. Oxisols are predominant in the reserve, and small streams are abundant in the area, resulting in an undulated terrain of plateaus of up to 140 m above sea level, crossed by many valleys [22]. The mean precipitation in the area is ~2,300 mm/yr, with most of the rainfall falling between November and May, and a short dry season between June and October [22].

We sampled the avifauna of the RFAD with mist-nets from January 2002 to July 2004, monthly from January to December 2002 (except June 2002, November 2003, and January to April 2004), whereas acoustic survey data were collected during March and May 2002. The RFAD is particularly suited for studies investigating the general heterogeneity of a tropical forest because it has a 64 km² grid, criss-crossed by trails that allow access to the entire reserve (Figure 1). Lines of mist-nets were systematically placed across the entire grid, along the nine parallel 8 km-long trails that cross the RFAD from east to west, covering 6,400 ha (Figure 1). In total, we sampled 72 individual areas, whereas acoustic transect surveys were randomly assigned and performed in 21 of them. Components of the forest structure were sampled in each of the 72 sample areas in 2,500 m² (50 × 50 m) plots. The 72 sample areas were located at least 1 km from one another, and at least 1 km from the borders of the RFAD, reducing possible edge effects.

2.2. Bird Surveys. Bird data was obtained using two independent methods: mist-nets and acoustic surveys along predefined transects. The use of mist-nets has several advantages over other types of survey methods [23], particularly being independent of identification skills (birds are easier to identify in the hand), easier to standardize (number of nets and/or time), and are good in capturing understory species that are moving through the forest. As its downside, it neglects an entire part of the community (i.e., all birds that forage and occur above 10 m, including most canopy birds). Acoustic surveys, on the other hand, are much more dependent on the skills and experience of the observer, and therefore its results are difficult to compare to other studies, but have the advantage of potentially recording many more species of birds with a much smaller sampling effort. Additionally, species recorded are not restricted to a particular stratum of the forest, although they rely on vocalizing individuals.

We placed lines of 20 mist-nets (12 m long, 2.5 m high, and 2.5 cm of mesh size) along existing trails, covering ~240 m. Each line started at mid-points (500 m, 1500 m, 2500 m, etc.), from the beginning of each trail that crosses the RFAD from east to west (Figure 1). Mist-nets were opened between 6 AM and 1 PM, for two consecutive days in the same place. Captured individuals were identified and marked with numbered aluminum bands (CEMAVE—Bird Migration Center, ICMBio, Brazil). Data from both days were pooled to obtain a single number of species and individuals (birds captured and marked on the first day were counted only once).

Acoustic surveys were performed by LNN along 21 1 km transects (see Figure 1). Transects started at 5:30 AM from the mid-point of the trail and included half an hour of dawn chorus census (which are mostly conducted in the dark and are important to record some species that are vocally active only during a few minutes before dawn). From 6 AM to 10 AM the entire 1 km was surveyed twice by walking slowly through the trail. Along the trails, bird vocalizations were tape-recorded, and the approximate location of each individual bird was noted. No playback was used during the surveys. The general location of each individual was noted to avoid double counting individual birds.

2.3. Forest Structure Components and Landscape Features Sampling. We established 72 50 × 50 m plots, located in the middle of each line of mist-nets (120 m from the beginning). Within plots, we quantified the following forest structure components and features of the landscape: (i) number of trees (diameter at breast height—DBH > 10 cm); (ii) number of fallen logs (>20 cm in diameter); (iii) number of snags (standing dead trees); (iv) leaf litter depth; (v) percentage of canopy openness; (vi) distance to the nearest stream; (vii) elevation.

We used the number of trees, fallen logs, and dead snags, as a measurement of tree abundance, and these were directly counted in the plots. We measured the depth of the litter from 20 subplots of 1 m² located along the sides of each 50 × 50 m plot (systematically distributed every 10 m) and from an additional subplot placed in the center of the plot. Litter depth was recorded by inserting a Swiss knife blade (7 cm length, 1 cm width, and 0.1 cm thick) into the forest ground until its tip touched the forest bare soil. The number of impaled dead leaves was used as the measurement of litter depth, using the mean value of the 21 subplots in the analyses. We recorded forest canopy openness using a Spherical Crown Densiometer (Concave—Mode C—Robert E. Lemon, Forest Densiometer—Bartlesville, OK, USA). Measurements were obtained from four readings (north, south, east, and west) at the corners and one at the center of the 50 × 50 m plots. Following factory recommendations, we multiplied each reading by 1.04. We used the mean value of five readings within a subplot as a subplot measurement and the mean value of 5 subplots in the analyses. Distance of the plots to the nearest stream and elevation (meters above sea level) of the plots were collected from a detailed topographic map.

2.4. Statistical Analyses. Although complementary, mist-net and acoustic survey data are not comparable and were analysed separately. To compare species composition among the 72 mist-net lines and for the 21 acoustic survey transects, we analysed both quantitative and qualitative data matrices using Principal Coordinates Analysis (PCoA), implemented in PC-ORD [24]. This analysis summarizes more information on one to three axes than other indirect ordination techniques, and is more robust to nonlinear effects [25]. We used the Bray-Curtis distance measure to obtain values of dissimilarity between sites. When used on presence/absence data, the Bray-Curtis index is known as the Sorensen distance

measure [26, 27], which has been used in ecological gradient studies [25, 28], including studies with plants [29], insects [30], and birds [6, 11, 13, 31, 32]. The resulting PCoA-scores were used as dependent variables in models of multivariate analysis of variance (MANOVA) and multiple regressions. We used two or three PCoA axes in the analyses because these generally explain most of the variance in the original variables for quantitative and qualitative data. We used *a posteriori* Pillai-Trace tests to verify whether MANOVA reveal significant differences among sites in relation to the topographic gradient (plateau-slope-valley), water basins (western and eastern sides of the RFAD, see Figure 1), and seasonality (dry and wet season and among years). For the water basin analysis we excluded mist-net lines and censuses located on the central plateau that divides the two basins (see Figure 1).

The Pillai-Trace statistics have been shown to be less sensitive to deviations from assumptions than other multivariate statistics [33, 34]. We also used multiple regressions followed by Pillai-Trace to evaluate the effects of the forest structure components on qualitative and quantitative bird community composition across the RFAD. These analyses were performed using GLM in Systat [35]. To verify potential problems of residual analysis in multiple regressions, we used a graphic method called partial residual plot, available in R (Core Development Team 2008). We also used R to verify possible linear relationships among predicting variables, estimating the variance inflation factor, which calculates the level of multicollinearity [36]. We used Mantel tests, implemented in PC-ORD [24], to verify spatial autocorrelation among variables, or the significance of relationships between assemblage matrices of similarity and distance between transects. We built Pearson correlation matrices to verify the correlation among forest structure components (independent variables). When these variables were significantly correlated, we included them in different analyses.

We built qualitative (presence/absence data) and quantitative (abundance data) matrices of species composition for the community analyses. We used regression models for qualitative and quantitative data, using PCoA axes as response variables in the regression models. In two regression models, the PCoA axes were regressed against five forest structure components as independent variables together (abundance of forest trees, abundance of fallen logs, abundance of snags, mean leaf litter depth, and proximity to the nearest stream), and in another two models, the PCoA-axes were regressed with just two independent variables together (elevation and canopy openness), because these two components were significantly correlated with the other five.

Throughout the study, we used three PCoA axes for quantitative data and two axes for qualitative data as dependent variables. For mist-net data, we found that three PCoA axes captured most of the variance in the original variables for quantitative data in the bird species matrix (Cumulative proportion of total variance, $C_{PV} = 0.63$), whereas two axes were enough for the presence/absence data matrix ($C_{PV} = 0.68$). Similarly, for the acoustic survey data three axes captured most of the variance for quantitative data ($C_{PV} = 0.60$), and two axes for presence/absence data ($C_{PV} = 0.56$).




FIGURE 2: Results from Principal Coordinate Analysis (PCoA) used to generate a simple ordination of 72 areas at the Reserva Florestal Adolpho Ducke (RFAD), Manaus, Brazil, based on the similarity matrix of distances generated by the Sorensen index for the qualitative matrix (presence and absence data) for the entire bird community. Letters represent the 72 areas in which the bird species were sampled using 240 m mist-net lines placed throughout the RFAD (p = plateau; s = slopes; v = valleys). The closer the points are to one another in the graph, the more similar are their bird species community composition. Note that the species composition, with a group of p in the upper part, a group of s in the middle, and a group of v in the lower part, changes significantly along the topographical gradient from plateau to valleys.

3.5. Effects of Forest Structure Components on the Avifauna.

We found no significant relation between the forest structure component parameters and the number of species (richness) and individuals (abundance) recorded in both mist-nets and acoustic surveys. Bird community composition (mist-net data) was significantly influenced by elevation, amount of leaf litter and distance to the nearest stream (Table 3). However, in all cases few species were strongly associated with the limits of the gradients, and most species occurred across the gradients of elevation, leaf litter and distance to stream (Figures 3(a), 3(b), and 3(c)). Changes in community composition (qualitative data) were significantly correlated with the amount of leaf litter, the distance to the nearest stream, and with terrain elevation (Table 3, Figures 4(a), 4(b), and 4(c)). When abundance of individuals was added to the community data (quantitative data), however, all relations lost significance (Table 3). Bird community composition (acoustic survey data) was also not significantly influenced by any of the seven forest structure components measured (data not shown).

3.6. Bird Species Composition between the Two Waterbasins (Eastern and Western Side). We found significant differences in the composition of bird species based on qualitative mist-net data between the two basins (eastern and western sides of the RFAD) (Manova, Pillai-Trace = 0.243; $F_{4,138} = 4.781$;

TABLE 1: Mantel randomization tests on forest structure components in relation to position of the sampling plots (geographical coordinates) in the 72 plots at the Reserva Florestal Adolpho Ducke, Manaus, central Amazonia.

Variable	Correlation value (r)	(Significance) P value
Tree abundance	0.048	0.040
Abundance of snags	-0.032	0.959
Abundance of fallen logs	-0.005	0.600
Mean leaf litter depth	0.065	0.020
Distance to nearest stream	0.032	0.550
Mean canopy openness	0.092	0.0001
Elevation	0.031	0.050

$P = 0.001$), but not for quantitative data (Manova, Pillai-Trace = 0.117; $F_{6,136} = 1.403$; $P = 0.218$). We also found significant differences in the composition of bird species between the two water basins using qualitative data from the acoustic survey (Manova, Pillai-Trace test = 0.600; $F_{3,17} = 8.506$; $P < 0.001$).

4. Discussion

This study adds to the growing body of evidence suggesting that the natural heterogeneity found in a tropical forest can affect the local composition of animal and plant communities [17, 21, 38–48]. Although we found that none of the variables analyzed were significantly associated with the number of bird species (richness) and individuals (abundance) recorded either on mist-nets or acoustic surveys, we found significant relationships between at least three parameters (elevation, distance to the nearest stream, and depth of the leaf litter) and bird species composition. Furthermore, we also found that changes in bird community composition can, in part, be attributed to a topographic gradient (plateau-slope-valley), and the water basins within the RFAD.

Finding composition changes without variation in species richness and bird abundance can be attributed to species turnover within the study area. In this case we are not referring to biogeographical or regional variation in the avifauna, but to local changes in the avian communities. Our surveys indicate that most bird species occur throughout the RFAD (data not shown) and the central plateau, which we found to be relevant to explain changes in bird species' composition, by no means represents a biogeographical or geographic barrier. Therefore, we are confident that our results indicate that birds are tracking differences in the landscape and the structure of the forest.

Our results show a significant response of the avian community to a topographic gradient (plateau-slope-valley), even though the range in elevation within our plots was only 75 m (39–114 m above sea level). Although this variation seems negligible, it is enough to create a topographic gradient that includes “high” flat areas (plateaus), “low” areas (valleys, which often flood on rainy days), and either gentle or steep slopes connecting the two. This topographic gradient has






FIGURE 3: Continued.



FIGURE 3: Bird species captured in 72 lines of mist-nets at Reserva Florestal Adolpho Ducke, near Manaus, Brazil, in relation to gradients in (a) elevation, (b) distance to the nearest stream, and (c) leaf litter depth. Species bars represent the presence of each species in a given line (presence/absence qualitative data), whereas height of bar at the bottom represents the quantitative value of those variables.




FIGURE 4: (a) The influence of terrain inclination (elevation), (b) stream proximity, and (c) depth of forest floor leaf litter on bird community composition in the Reserva Florestal Adolpho Ducke, Manaus, Brazil. Qualitative data (species presence/absence for the 72 transects with mist-nets), which were subjected to ordination analyses (PCoA). Only the second PCoA-axes were used to construct the graphs, the ones showing statistical significance.

been related to several soil parameters, particularly the amount of clay or sand in the soil [39], which in turn has been shown to be the best predictor of above-ground tree biomass at the RFAD [21] and other sites in central Amazonia [49]. This topographic gradient has also been shown to affect the distribution of palms [42], ants [43], and frogs [44], and is now shown for the first time in birds at the RFAD. We did not find a significant effect of the topography on the bird community sampled using acoustic surveys, and we be-

lieve that a main reason for such a difference is that, whereas lines of mist-nets covered ~ 240 m, acoustic surveys covered 1 km. It is more likely that several topographic classes are present in 1 km than in 0.25 km, and this may obscure possible relationships.

The depth of leaf litter on the forest ground can have several indirect effects on the avifauna, offering more substrate for small and large invertebrates for insectivorous birds, and even small vertebrates on which raptors (e.g., forest-falcons and owls) can prey [38]. Many bird species are known to actively search under dead leaves on the ground, such as leaf-tossers (*Sclerurus*) and foliage-gleaners (*Automolus*), and using leaf litter is a common practice in North America to attract birds to people's backyards. On the other hand, leaf litter is very dynamic, and can change dramatically in space (flat versus steep areas) and time (dry versus wet season). Hence, we are not sure which mechanisms can maintain its effect on an entire avifauna in the long term.

Forest streams are a vital component of the forest landscape and correlated to valleys. Therefore, the relationship between species composition and proximity to the microhabitats formed by the streams was expected. In fact, several bird species are restricted to valleys and streams, as shown by our ordination analysis. Within our sample, we identified almost 30 species that occur more often near streams (Figure 3(b)), including some species that are rarely recorded away from them (e.g., *Topaza pella*, *Philydor pyrrhodes*, and *Schistocichla leucostigma*). Proximity to streams is also known to influence the distribution and abundance of terrestrial herbs [41, 50] and palms [42] at the RFAD.

The number of trees, fallen logs, snags, and canopy openness are variables known to affect individual birds species [7] or groups [38], yet they had little effect on the number of species or individuals (richness and abundance), and the species composition associated with our plots. One potential problem in assessing the effect of local environmental variables on birds is that, differently from herbs, palms, or frogs, most bird species hold relatively large territories for their size. Data from throughout the Neotropics suggest that even small passerines occupy (and actively defend) between 10 and 20 ha of primary forest [2, 46]. How then can we relate landscape features and small-scale forest parameters in plots of 0.25 ha (50×50 m) with birds that may occupy an area 50 to 75 times larger? Our understanding on the use of space by tropical birds remains limited, and it is not well known how much of their territories birds actually use, or how much of their territory is actually needed for activities such as feeding or nesting. Although weak, the spatial autocorrelation that we found in variables such as tree abundance, leaf litter depth, canopy openness, and elevation may indicate that our plots may represent the forest well beyond the plots' 50×50 m boundaries and may actually describe large enough areas to hold entire territories of individual birds. Similarly, plateaus, slopes, and valleys are continuous habitats, and one can walk (or a bird fly) for several kilometers on a plateau avoiding the valleys. Whether birds do this and select their use of territory in relation to topographic gradients or simply use their entire territory is not clear yet and deserve further investigation.

TABLE 2: Pearson correlation matrix among the forest structure components recorded in 72 plots at the Reserva Florestal Adolpho Ducke, Manaus, central Amazonia, used in the avian community analyses; ** $P < 0.001$; * $P < 0.05$ resulting from the Bonferroni probability matrix.

	Abundance of forest trees	Abundance of fallen logs	Abundance of snags	Leaf litter depth	Forest canopy openness (%)	Elevation (m)
Abundance of fallen logs	0.208					
Abundance of snags	0.083	0.315				
Leaf litter depth	0.047	0.318	0.255			
Forest canopy openness ⁺	0.134	0.560**	0.474**	0.528**		
Elevation ⁺	0.118	0.093	0.035	0.290	0.242	
Distance to nearest stream	0.053	0.129	0.019	0.020	-0.058	0.370*

⁺ These two were significantly correlated to other variables; therefore, canopy openness and elevation were analyzed in a separate regression model.

TABLE 3: Results of the multiple regression analyses performed to test the effect of the seven forest structure components on qualitative and quantitative bird composition. Analyses were performed on scores from the Principal Coordinate Analyses (PCoA).

Data (matrix)	Forest components	Qualitative (presence/absence)				Quantitative (abundance)			
		Pillai-Trace	<i>F</i>	DF	<i>P</i>	Pillai-Trace	<i>F</i>	DF	<i>P</i>
Bird community	Model 1								
	Tree abundance	0.084	1.958	3; 64	0.129	0.085	1.986	3; 64	0.125
	Log abundance	0.049	1.109	3; 64	0.352	0.069	1.580	3; 64	0.203
	Snag abundance	0.023	0.500	3; 64	0.683	0.043	0.961	3; 64	0.417
	Leaf litter depth	0.111 (-)	2.667	3; 64	0.055	0.036	0.800	3; 64	0.498
	Distance to stream	0.118 (-)	2.861	3; 64	0.044	0.088	2.062	3; 64	0.114
	Model 2								
	Elevation	0.209 (+)	5.919	3; 67	0.001	0.079	1.928	3; 67	0.133
	Canopy openness	0.033	0.761	3; 67	0.520	0.015	0.351	3; 67	0.789

Because elevation and canopy openness were significantly correlated to other variables (see Table 1), they were analyzed in separate multiple linear regression models (see Section 3 and Figure 3). Three or two PCoA-axes were used in the models for quantitative and qualitative analysis, respectively (see Section 2). The negative and positive signals within parenthesis are just to indicate the directions of the relationships and are not related to Pillai-Trace values (see also Figure 4).

The central plateau that divides the RFAD in two halves, results in two relatively similar-sized basins with significantly different community compositions. Similar results were found for the community composition of herbs [42] and fish [45]. Therefore, it is possible that the differences in bird communities between basins are due to differences in forest biomass and soil, which is higher in the eastern part of the RFAD [21]. Higher biomass may result in more food resources and enhanced opportunities for foraging, but these indirect relationships remain speculative.

Despite the fact that the RFAD is rapidly becoming a large forest fragment, its avifauna seems to remain untouched, if compared to neighboring areas with continuous forests [14]. Not only did we record most species expected to be found in a healthy forest, but independently of being found in our standardized surveys, important species of conservation concern are known to occur at the RFAD (see Table 4). Among these are several large frugivores, such as guans, curassows, trumpeters, toucans, and macaws, which are the first species to be hunted to local extinction by surrounding human populations. Similarly important is the presence of several large-bodied top predators such as Harpy Eagle (*Harpia harpyja*), Crested Eagle (*Morphnus guianensis*), Black-and-white Hawk-Eagle (*Spizaetus melanoleucus*), Black Hawk-Eagle (*Spizaetus tyrannus*), and Ornate Hawk-Eagle (*Spiza-*

etus ornatus), many of which are known to nest in the RFAD (T.M. Sanaiotti, pers. comm.). Other species seem to be truly rare at the RFAD and were recorded only once in our surveys, for example, Crimson Topaz (*Topaza pella*), Royal Flycatcher (*Onychorhynchus coronatus*), Guianan Red-Cotinga (*Phoenicircus carnifex*), and Slaty-backed Forest-Falcon (*Micrastur mirandolleii*). These species seem to be generally rare or locally uncommon in central Amazonia [14], independently of the conservation condition of the forest. In fact, virtually all species considered vulnerable to fragmentation or sensitive to fragment area [47] were found at the RFAD [48].

Despite the good conditions of the RFAD's avifauna, which was last surveyed by us more than five years ago [7, 32, 38, 48], careful monitoring is important, particularly given the continuous pressure created by the growth of the city of Manaus. Therefore, we recommend that long-term monitoring of bird communities using standardized methods be done at the RFAD [54] in order to ensure that processes of species loss (if unavoidable) are well documented. The only active measure that can potentially delay or protect the RFAD from species loss is maintaining the connectivity of the RFAD on its eastern edge and creating a protected forest corridor from the RFAD to other large tracks of forest north of Manaus.

TABLE 4: Bird species recorded at the Reserva Florestal Adolpho Ducke, central Amazonia, Brazil (see codes at end). Taxonomy and systematic order follow the Comitê Brasileiro de Registros Ornitológicos (CBRO) as of August 2009 [51].

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
TINAMIDAE								
<i>Tinamus major</i> (Gmelin, 1789)	1100c	s	c	a, i, s, f	x	g	a	inhambu-de-cabeça-vermelha
<i>Crypturellus soui</i> (Hermann, 1783)	220c	s	u	a, i, s, f	x	g	o	tururim
<i>Crypturellus variegatus</i> (Gmelin, 1789)	384b	s	c	a, i, s, f	x	g	a	inhambu-anhangá
<i>Crypturellus brevirostris</i> (Pelzeln, 1863)	—	s	u	a, i, s, f	x	g	a	inhambu-carijó
CRACIDAE								
<i>Ortalis motmot</i> (Linnaeus, 1766)	445c	g	c	a, i, fr, s		u		aracua-pequeno
<i>Penelope marail</i> (Statius Muller, 1776)	840c	s, g	c	a, i, fr, s	x	c, u	a	jacumirim
<i>Crax allector</i> (Linnaeus, 1766)	3100c	s, g	u	a, i, fr, s	x	g, u		mutum-poranga
ODONTOPHORIDAE								
<i>Odontophorus guianensis</i> (Gmelin, 1789)	255b	g	u	a, i, fr, s	x	g	a	uru-corcovado
ARDEIDAE								
<i>Ardea alba</i> (Linnaeus, 1758)	885c	s	u	fo, fi		g		garça-branca-grande
CICONIIDAE								
<i>Mycteria americana</i> (Linnaeus, 1758)	2500c	s	r	fi		u		cabeça-seca
CATHARTIDAE								
<i>Coragyps atratus</i> (Bechstein, 1793)	1800c	s, g	c	d		c		urubu-de-cabeça-preta
<i>Cathartes aura</i> (Linnaeus, 1758)	2000c	s	c	d	x	c		urubu-de-cabeça-vermelha
<i>Cathartes melambrotus</i> (Wetmore, 1964)	1400c	s	u	d	x	c		urubu-da-mata
<i>Sarcoramphus papa</i> (Linnaeus, 1758)	3300c	s	r	d		c		urubu-rei
PANDIONIDAE								
<i>Pandion haliaetus</i> (Linnaeus, 1758)	1700c	s	b	fi		c		águia-pescadora
ACCIPITRIDAE								
<i>Leptodon cayanensis</i> (Latham, 1790)	605c	s	r	bi		c		gavião-de-cabeça-cinza
<i>Chondrohierax uncinatus</i> (Temminck, 1822)	300c	s	r	b, r		c		caracoleiro
<i>Elanoides forficatus</i> (Linnaeus, 1758)	445c	s	?	a, i		c		gavião-tesoura
<i>Gampsonyx swainsonii</i> (Vigors, 1825)	95c	s	r	a, i		c, u		gaviãozinho
<i>Harpagus bidens</i> (Latham, 1790)	170c	s	u	a, i, b	x	u		gavião-ripina
<i>Ictinia plumbea</i> (Gmelin, 1788)	280c	s	?	i		c		sovi
<i>Accipiter superciliosus</i> (Linnaeus, 1766)	80b	s	r	bi, ro		c, u		gavião-miudinho
<i>Accipiter bicolor</i> (Vieillot, 1817)	230c	s	r	bi, ro		c, u		gavião-bombachinha-grande
<i>Leucopternis melanops</i> (Latham, 1790)	330c	s	r	bi, sn	x	c, u	a	gavião-de-cara-preta
<i>Leucopternis albigollis</i> (Latham, 1790)	750c	s	c	bi, sn	x	c, u	o	gavião-branco
<i>Buteogallus urubitinga</i> (Gmelin, 1788)	1100c	s	u	bi, fi, sn	x	c, u		gavião-preto

TABLE 4: Continued.

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
<i>Rupornis magnirostris</i> (Gmelin, 1788)	265c	s	c	bi, li, sn	x	c, u		gavião-carijó
<i>Buteo nitidus</i> (Latham, 1790)	475c	s	c	bi, li, sn	x	c, u		gavião-pedrés
<i>Buteo platypterus</i> (Vieillot, 1823)	450c	s	b	bi, li, sn		c, u		gavião-de-asa-larga
<i>Morphnus guianensis</i> (Daudin 1800)	1800c	s	r	mo, sl, ro	x	c, u		uiiraçu-falso
<i>Harpia harpyja</i> (Linnaeus, 1758)	6500c	s	r	mo, sl, ro	x	c, u		gavião-real
<i>Spizaetus melanoleucus</i> (Vieillot, 1816)	815c	s	r	bi, mo, ro		c		gavião-pato
<i>Spizaetus tyrannus</i> (Wied, 1820)	1000c	s	r	bi, mo, ro	x	c		gavião-pegá-macaco
<i>Spizaetus ornatus</i> (Daudin, 1800)	1200c	s	u	bi, mo, ro	x	c	a	gavião-de-penacho
FALCONIDAE								
<i>Ibycter americanus</i> (Boddaert, 1783)	570c	g	c	a, i, li, bi	x	c	a	gralhão
<i>Milvago chimachima</i> (Vieillot, 1816)	325c	s	c	a, i, li, bi		u, g		carrapateiro
<i>Micrastur ruficollis</i> (Vieillot, 1817)	176	s*	c	a, i, li, bi	x	u	o	falcão-caburé
<i>Micrastur gilvicolis</i> (Vieillot, 1817)	226	s	c	a, i, li, bi	x	u	o, a	Falcão-mateiro
<i>Micrastur mirandollei</i> (Schlegel, 1862)	420c	s	u	a, i, li, bi		u	a	tanatau
<i>Micrastur semitorquatus</i> (Vieillot, 1817)	535c	s	u	a, i, li, bi		u		falcão-relógio
<i>Falco ruficularis</i> (Daudin, 1800)	140c	s	c	a, i, ba, bi	x	c		cauré
<i>Falco peregrinus</i> (Tunstall, 1771)	600c	s	b	a, i, ba, bi		c		falcão-peregrino
PSOPHIIDAE								
<i>Psophia crepitans</i> (Linnaeus, 1758)	1300c	g	u	a, i, fr, s	x	g, u	o	jacamim-de-costas-cinzentas
RALIDAE								
<i>Aramides cajanea</i> (Statius Muller, 1776)	410c	s	c	a, i	x	g		saracura-três-potes
<i>Laterallus viridis</i> (Statius Muller, 1776)	64c		u	a, i		g		sanã-castanha
EURYPIDIDAE								
<i>Eurypyga helias</i> (Pallas, 1781)	220c	s	r	a, i	x	g		pavãozinho-do-Pará
SCOLOPACIDAE								
<i>Tringa solitaria</i> (Wilson, 1813)	47c	s	b	a, i		g		maçarico-solitário
COLUMBIDAE								
<i>Columbina passerina</i> (Linnaeus, 1758)	32c	s	u	s	x	c, u		rolinha-cinza
<i>Columbina talpacoti</i> (Temminck, 1811)	47c	s	c	s	x	g, u		rolinha-roxa
<i>Patagioenas cayennensis</i> (Bonnaterre, 1792)	230c	s	u	s	x	c, u		pomba-galega
<i>Patagioenas plumbea</i> (Vieillot, 1818)	180c	s	c	s	x	c, u	a	pomba-amargosa

TABLE 4: Continued.

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
<i>Patagioenas subvinacea</i> (Lawrence, 1868)	128b	s	c	s	x	c, u	a	pomba-botafofo
<i>Leptotila verreauxi</i> (Bonaparte, 1855)	155c	s	c	s	x	g, u		juriti-pupu
<i>Geotrygon montana</i> (Linnaeus, 1758)	122	s	c	S	x	g, u	o, a	pariri
PSITTACIDAE								
<i>Ara ararauna</i> (Linnaeus, 1758)	1100c	c, g	u	fr, s		c		arara-canindé
<i>Ara macao</i> (Linnaeus, 1758)	1000c	c, g	c	fr, s		c	a	araracanga
<i>Ara chloropterus</i> (Gray, 1859)	1250c	c, g	c	fr, s	x	c	a	arara-vermelha-grande
<i>Orthopsittacus manilata</i> (Boddaert, 1783)	370c	c, g	u	fr, s	x	c		maracanã-do-buriti
<i>Aratinga leucophthalma</i> (Statius Muller, 1776)	160c	g	u	fr, s	x	c		periquitão-maracanã
<i>Brotogeris chrysoptera</i> (Linnaeus, 1766)	65c	g	c	fr, s	x	c	a	periquito-de-asa-dourada
<i>Touit purpuratus</i> (Gmelin, 1788)	60c	g	u	fr, s		c	a	apuim-de-costas-azuis
<i>Pyrilia caica</i> (Latham, 1790)	130c	g	u	fr, s	x	c	a	curica-caica
<i>Pionus menstruus</i> (Linnaeus, 1766)	245c	g	c	fr, s	x	c	a	maitaca-de-cabeça-azul
<i>Pionus fuscus</i> (Statius Muller, 1776)	205c	s	u	fr, s		c	a	maitaca-roxa
<i>Amazona autumnalis</i> (Linnaeus, 1758)	415c	g	c	fr, s	x	c	a	papagaio-diadema
<i>Amazona farinosa</i> (Boddaert, 1783)	620c	g	u	fr, s		c		papagaio-moleiro
<i>Derophtus accipitrinus</i> (Linnaeus, 1758)	240c	s	u	fr, s		c	a	anacã
CUCULIDAE								
<i>Piaya cayana</i> (Linnaeus, 1766)	95c	s	u	a, i, fo, li	x	u		alma-de-gato
<i>Piaya melanogaster</i> (Vieillot, 1817)	100c	s	c	a, i, fo, li	x	u	a	chincão-de-bico-vermelho
<i>Crotophaga ani</i> (Linnaeus, 1758)	95c	g	c	a, i, fo, li	x	u		anu-preto
STRIGIDAE								
<i>Megascops watsonii</i> (Cassin, 1849)	147b	s	c	a, i, ba, ro	x	u	a	corujinha-orelhuda
<i>Lophotrix cristata</i> (Daudin, 1800)	545c	s	c	a, i, ba, ro	x	u		coruja-de-crista
<i>Pulsatrix perspicillata</i> (Latham, 1790)	850c	s	u	a, i, ba, ro	x	c		murucututu
<i>Strix huhula</i> (Daudin, 1800)	385c	s	u	a, i, ba, ro	x	u		coruja-preta
<i>Strix virgata</i> (Cassin, 1849)	260c	s	r	a, i, ba, ro	x	u		coruja-do-mato
<i>Glauclidium hardyi</i> (Viellard, 1990)	55b	s	c	a, i, ba, ro	x	u	a	caburé-da-amazônia
NYCTIBIDAE								
<i>Nyctibius grandis</i> (Gmelin, 1789)	550c	s	r	i	x	c, u		mãe-da-lua-gigante
<i>Nyctibius aethereus</i> (Wied, 1820)	320	s	r	i		c, u		mãe-da-lua-parda
<i>Nyctibius griseus</i> (Gmelin, 1789)	35c	s	u	i	x	c, u	a	mãe-da-lua
<i>Nyctibius leucopterus</i> (Wied, 1821)	—	s	u	i	x	c, u		urutau-de-asa-branca
CAPRIMULGIDAE								
<i>Lurocalis semitorquatus</i> (Gmelin, 1789)	75c	s	u	i		g		tuju
<i>Nyctidromus albigollis</i> (Gmelin, 1789)	145	s	c	i	x	g		bacurau

TABLE 4: Continued.

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
<i>Caprimulgus nigrescens</i> (Cabanis, 1848)	136	s	u	i	x	g		bacurau-de-lajeado
APOIDIDAE								
<i>Sireptoprocne zonaris</i> (Shaw, 1796)	90c	g	r	i		a		taperuçu-de-coleira-branca
<i>Chaetura spinicaudus</i> (Temminck, 1839)	15c	g	c	i		a		andorinhão-de-sobre-branco
<i>Chaetura chapmani</i> (Hellmayr, 1907)	22c	g	c	i		a		andorinhão-de-chapman
<i>Chaetura brachyura</i> (Jardine, 1846)	20c	g	u	i		a		andorinhão-de-rabo-curto
<i>Panyptila cayennensis</i> (Gmelin, 1789)	18c	s	r	i		a		andorinhão-estofador
<i>Tachornis squamata</i> (Cassin, 1853)	11c	s	r	i	x	a		tesourinha
TROCHILIDAE								
<i>Glauis hirsutus</i> (Gmelin, 1788)	4	s	r	a, i, n	x	c, u	o	balança-rabo-de-bico-torto
<i>Phaethornis superciliosus</i> (Linnaeus, 1766)	6	s	c	a, i, n	x	c, u	a, o	rabo-branco-de-bigodes
<i>Phaethornis bourcieri</i> (Lesson, 1832)	5	s	c	a, i, n	x	c, u	a, o	rabo-branco-de-bico-reto
<i>Phaethornis ruber</i> (Linnaeus, 1758)	2c	s	r	a, i, n	x	c, u		rabo-branco-rubro
<i>Camptylopterus largipennis</i> (Boddaert, 1783)	6	s	c	a, i, n	x	c, u	a, o	asa-de-sabre-cinza
<i>Florisuga mellivora</i> (Linnaeus, 1758)	6	s	u	a, i, n	x	c, u		beija-flor-azul-de-rabo-branco
<i>Anthracothorax nigricollis</i> (Vieillot, 1817)	7c	s	r	a, i, n	x	c, u		beija-flor-de-veste-preta
<i>Topaza pella</i> (Linnaeus, 1758)	12	s	r	a, i, n	x	c, u	a, o	beija-flor-brilho-de-fogo
<i>Discosura longicaudus</i> (Gmelin, 1788)	3c	s	r	a, i, n		c, u		Bandeirinha
<i>Thalurania furcata</i> (Gmelin, 1788)	5	s	c	a, i, n	x	c, u	a, o	beija-flor-tesoura-verde
<i>Hylocharis sapphirina</i> (Gmelin, 1788)	3	s	u	a, i, n	x	c, u	a	beija-flor-safira
<i>Polytmus theresiae</i> (Da Silva Maia, 1843)	4c	s	u	a, i, n	x	c, u		beija-flor-verde
<i>Amazilia versicolor</i> (Vieillot, 1818)	—	s	r	a, i, n	x	c, u	o	beija-flor-de-banda-branca
<i>Amazilia fimbriata</i> (Gmelin, 1788)	8c	s	c	a, i, n	x	c, u	o	beija-flor-de-garganta-verde
<i>Heliothryx auritus</i> (Gmelin, 1788)	4	s	c	a, i, n	x	c, u	a, o	beija-flor-de-bochecha-azul
TROGONIDAE								
<i>Trogon melanurus</i> (Swainson, 1838)	92b	s	c	a, i, fr, s	x	c, u	a	surucua-de-cauda-preta
<i>Trogon viridis</i> (Linnaeus, 1766)	49b	s	c	a, i, fr, s	x	c, u	a	s.-grande-de-barriga-amarela
<i>Trogon violaceus</i> (Gmelin, 1788)	54b	s	c	a, i, fr, s	x	c, u	a	surucua-pequeno
<i>Trogon rufus</i> (Gmelin, 1788)	52	s	u	a, i, fr, s	x	c, u	a, o	surucua-de-barriga-amarela

TABLE 4: Continued.

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
<i>Pharomachrus pavoninus</i> (Spix, 1824)	—	s	r	a, i, fr, s	x	c, u	a	surucua-pavão
ALCENIDIDAE								
<i>Megasceryle torquata</i> (Linnaeus, 1766)	300c	s	r	fi	x	u		martim-pescador-grande
<i>Chloroceryle amazona</i> (Latham, 1790)	110c	s	r	fi	x	u		martim-pescador-verde
<i>Chloroceryle aenea</i> (Pallas, 1764)	16	s	r	fi	x	u	o	martinho
<i>Chloroceryle americana</i> (Gmelin, 1788)	27	s	r	fi	x	u		martim-pescador-pequeno
<i>Chloroceryle inda</i> (Linnaeus, 1766)	50	s	c	fi	x	u	o	martim-pescador-da-mata
MOMOTIDAE								
<i>Momotus momota</i> (Linnaeus, 1766)	152	s	c	a, i	x	g, u	a, o	udu-de-coroa-azul
GALBULIDAE								
<i>Galbula albirostris</i> (Latham, 1790)	27	s	c	a, i	x	u	a, o	ariramba-de-bico-amarelo
<i>Galbula leucogastra</i> (Vieillot, 1817)	16c	s	r	a, i	x	u	a, o	ariramba-bronzeada
<i>Galbula dea</i> (Linnaeus, 1758)	30c	s	c	a, i	x	c, u	a, o	ariramba-do-paraiso
<i>Jacamerops aureus</i> (Statius Muller, 1776)	61b	s	u	a, i	x	c, u	a, o	jacamaracu
BUCCONIDAE								
<i>Notharchus macrorhynchos</i> (Gmelin, 1788)	53b	s	c	a, i, fo, li	x	g, u	a	macuru-de-testa-branca
<i>Notharchus tectus</i> (Boddaert, 1783)	27c	s	u	a, i, fo, li		g, u		macuru-pintado
<i>Bucco tamatia</i> (Gmelin, 1788)	36	s	u	a, i, fo, li	x	g, u	a, o	rapazinho-carijó
<i>Bucco capensis</i> (Linnaeus, 1766)	52	s	r	a, i, fo, li	x	g, u	a, o	rapazinho-de-colar
<i>Malacoptila fusca</i> (Gmelin, 1788)	42	s	u	a, i, fo, li	x	g, u	o	barbudo-pardo
<i>Nonnula rubecula</i> (Spix, 1824)	21b	s	r	a, i, fo, li		g, u		macuru
<i>Monasa atra</i> (Boddaert, 1783)	85	s	c	a, i, fo, li	x	g, u	a, o	chora-chuva-de-asa-branca
<i>Chelidoptera tenebrosa</i> (Pallas, 1782)	35c	s	r	a, i, fo, li	x	g, u		urubuzinho
CAPITONIDAE								
<i>Capito niger</i> (Statius Muller, 1776)	55c	s	c	a, i, fr, s		c, u	a	capitão-de-bigode-carijó
RAMPHASTIDAE								
<i>Ramphastos tucanus</i> (Linnaeus, 1758)	600c	c, g	c	a, bi, i, fr, s	x	c, u	a	tucano-grande-de-papo-branco
<i>Ramphastos vitellinus</i> (Lichtenstein, 1823)	372	c, g	c	a, bi, i, fr, s	x	c, u	a, o	tucano-de-bico-preto
<i>Selenidera piperivora</i> (Linnaeus, 1766)	149b	c	c	a, bi, i, fr, s		c, u	a	araçari-negro
<i>Pteroglossus viridis</i> (Linnaeus, 1766)	140b	g	c	a, bi, i, fr, s	x	c, u	a, o	araçari-miudinho
PICIDAE								
<i>Picumnus exilis</i> (Lichtenstein, 1823)	9c	s, c	r	a, i, fr, s		c, u		pica-pau-anão-de-pinta-amarela
<i>Melanerpes cruentatus</i> (Boddaert, 1783)	58c	s, c, g	c	a, i, fr, s	x	c, u	a	benedito-de-testa-vermelha

TABLE 4: Continued.

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
<i>Veniliornis cassini</i> (Malherbe, 1862)	33b	s, c	c	a, i, fr, s	x	c, u	a, o	pica-pau-de-colar-dourado
<i>Piculus flavigula</i> (Boddaert, 1783)	33b	s, c	c	a, i, fr, s	x	c, u	a, o	pica-pau-bufador
<i>Piculus chrysochloros</i> (Vieillot, 1818)	83c	s, c	r	a, i, fr, s	x	c, u		pica-pau-dourado-escuro
<i>Celeus undatus</i> (Linnaeus, 1766)	64c	s, c	r	a, i, fr, s		c, u	a	pica-pau-barrado
<i>Celeus elegans</i> (Statius Muller, 1776)	128	s, c	u	a, i, fr, s	x	c, u	a, o	pica-pau-chocolate
<i>Celeus flavus</i> (Statius Muller, 1776)	105c	s, c	u	a, i, fr, s		c, u		pica-pau-amarelo
<i>Celeus torquatus</i> (Boddaert, 1783)	120c	s, c	u	a, i, fr, s		c, u	a	pica-pau-de-coleira
<i>Dryocopus lineatus</i> (Linnaeus, 1766)	200c	s, c	c	a, i, fr, s	x	c, u		pica-pau-de-banda-branca
<i>Campephilus rubricollis</i> (Boddaert, 1783)	216b	s, c	c	a, i, fr, s	x	c, u	a	pica-pau-de-barriga-vermelha
THAMNOPHILIDAE								
<i>Gymbilaimus lineatus</i> (Leach, 1814)	35b	s	c	a, i	x	u	a	papa-formiga-barrado
<i>Frederickena viridis</i> (Vieillot, 1816)	66	s	r	a, i	x	u	a, o	borralhara-do-norte
<i>Thamnophilus murinus</i> (Slater & Salvin, 1868)	16	s	c	a, i	x	u	a, o	choca-murina
<i>Thamnophilus punctatus</i> (Shaw, 1809)	20c	s	u	a, i	x	u	o	choca-bate-cabo
<i>Thamnomanes ardesiacus</i> (Slater & Salvin, 1867)	20	s	c	a, i	x	c, u	a, o	uirapuru-de-garganta-preta
<i>Thamnomanes caesi</i> (Temminck, 1820)	18	s	c	a, i	x	c, u	a, o	ipecuá
<i>Epinecrophylia gutturalis</i> (Slater & Salvin, 1881)	10	s	c	a, i	x	u	a, o	choquinha-de-barriga-parda
<i>Myrmotherula brachyura</i> (Hermann, 1783)	7c	s	c	a, i		u	a	choquinha-miúda
<i>Myrmotherula guttata</i> (Vieillot, 1825)	10b	s	u	a, i	x	u		choquinha-de-barriga-ruiva
<i>Myrmotherula axillaris</i> (Vieillot, 1817)	8	s	u	a, i	x	u	a	choquinha-de-flanco-branco
<i>Myrmotherula longipennis</i> (Pelzelin, 1868)	11	s	c	a, i	x	u	a, o	choquinha-de-asa-comprida
<i>Myrmotherula menetriesii</i> (d'Orbigny, 1837)	11	s	c	a, i	x	u	a, o	choquinha-de-garganta-cinza
<i>Herpisilochmus dorsimaculatus</i> (Pelzelin, 1868)	9c	s	c	a, i		c, u	a	chorozinho-de-costa-manchada
<i>Terenura spodiopila</i> (Slater & Salvin, 1881)	6c	s	c	a, i	x	u	a	zidedê-de-asa-cinza
<i>Cercomacra cinerascens</i> (Slater, 1857)	21	s	c	a, i	x	u	a	chororó-pocuá
<i>Cercomacra tyrannina</i> (Slater, 1855)	17	s	r	a, i	x	u	o	chororó-escuro

TABLE 4: Continued.

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
<i>Hypocnemis cantator</i> (Boddaert, 1783)	14	s	c	a, i	x	u	a	papa-formiga-cantador
<i>Pernostola rufifrons</i> (Gmelin, 1789)	30	s*	c	a, i	x	g, u	a, o	formigueiro-de-cabeça-preta
<i>Schistocichla leucostigma</i> (Pelzelin, 1868)	24	s	u	a, i	x	g, u	a, o	formigueiro-de-asa-pintada
<i>Myrmeciza ferruginea</i> (Statius Muller, 1776)	18	s	c	a, i	x	u	a, o	formigueiro-ferrugem
<i>Myrmeciza atrothorax</i> (Boddaert, 1783)	18b	s	u	a, i	x	u	o	formigueiro-de-peito-preto
<i>Myrmormis torquata</i> (Boddaert, 1783)	45b	s	u	a, i	x	g		pinto-do-mato-carijó
<i>Pithys albifrons</i> (Linnaeus, 1766)	22	s*	c	a, i	x	u	o	papa-formiga-de-topete
<i>Gymnophithys rufigula</i> (Boddaert, 1783)	34	s*	c	a, i	x	u	a, o	mãe-de-taoca-garganta-vermelha
<i>Willisornis poecilinotus</i> (Cabanis, 1847)	16c	s*	c	a, i	x	g, u	a, o	Rendadinho
CONOPOPHAGIDAE								
<i>Conopophaga aurita</i> (Hermann, 1783)	24b	s	u	a, i	x	u		chupa-dente-de-cinta
GRALLARIIDAE								
<i>Grallaria varia</i> (Boddaert, 1783)	128	s	c	a, i	x	g	a, o	tovaçu
<i>Hylopezus macularius</i> (Temminck, 1823)	42b	s	u	a, i		g	a	torom-carijó
<i>Myrmothera campanisoma</i> (Hermann, 1783)	48b	s	c	a, i	x	g	a	tovaca-patinho
FORMICARIIDAE								
<i>Formicarius colma</i> (Boddaert, 1783)	54	s	c	a, i	x	g	a, o	galinha-do-mato
<i>Formicarius analis</i> (d'Orbigny & Lafresnaye, 1837)	62b	s	c	a, i		g		pinto-do-mato-de-cara-preta
SCLERURIDAE								
<i>Sclerurus mexicanus</i> (Sclater, 1857)	28	s	u	a, i	x	u	a, o	vira-folha-de-peito-vermelho
<i>Sclerurus rufigularis</i> (Pelzelin, 1868)	21b	s	c	a, i	x	u	a, o	vira-folha-de-bico-curto
<i>Sclerurus caudacutus</i> (Vieillot, 1816)	20	s	r	a, i	x	u	o	vira-folha-pardo
DENDROCOLAPTIDAE								
<i>Dendrocincla fuliginosa</i> (Vieillot, 1818)	46	s*	c	a, i	x	c, u	a, o	arapaçu-pardo
<i>Dendrocincla merula</i> (Lichtenstein, 1829)	63	s*	c	a, i	x	c, u	o	arapaçu-da-taoca
<i>Deconychura longicauda</i> (Pelzelin, 1868)	31	s	c	a, i	x	c, u	o	arapaçu-rabudo
<i>Deconychura stictolaema</i> (Pelzelin, 1868)	18	s	c	a, i	x	c, u	a, o	arapaçu-de-garganta-pintada
<i>Sittasomus griseicapillus</i> (Vieillot, 1818)	14b	s	c	a, i	x	c, u	a	arapaçu-verde
<i>Glyphorhynchus spirurus</i> (Vieillot, 1819)	12	s	c	a, i	x	c, u	a, o	arapaçu-de-bico-de-cunha
<i>Dendrexetastes rufigula</i> (Lesson, 1844)	70b	s	u	a, i	x	c, u	a	arapaçu-galinha
<i>Hylexetastes perrotii</i> (Lafresnaye, 1844)	123	s*	u	a, i	x	c, u	a, o	arapaçu-de-bico-vermelho
<i>Dendrocolaptes certhia</i> (Boddaert, 1783)	66	s*	c	a, i	x	c, u	a, o	arapaçu-barrado

TABLE 4: Continued.

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
<i>Dendrocolaptes picumnus</i> (Lichtenstein, 1820)	35	s*	u	a, i	x	c, u	a, o	arapaçu-meio-barrado
<i>Xiphorhynchus pardalotus</i> (Viellot, 1818)	53	s	c	a, i	x	c, u	a, o	arapaçu-assobiador
<i>Lepidocolaptes albolineatus</i> (Lafresnaye, 1845)	19b	s	c	a, i		c, u	a	arapaçu-de-listras-brancas
<i>Campylorhamphus procurviroides</i> (Lafresnaye, 1850)	42	s	u	a, i	x	c, u	a, o	arapaçu-de-bico-curvo
FURNARIIDAE								
<i>Synallaxis rutilans</i> (Temminck, 1823)	17b	c	u	a, i	x	u	o	joão-teneném-castanho
<i>Philydor erythrocercus</i> (Pelzel, 1859)	23b	s	c	a, i		u	a	limpa-folha-de-sobre-ruivo
<i>Philydor pyrrhodes</i> (Cabanis, 1848)	30	s	u	a, i	x	u	a, o	limpa-folha-vermelho
<i>Automolus ochrolaemus</i> (Tschudi, 1844)	35b	s	c	a, i	x	u		barranqueiro-camurça
<i>Automolus infuscatus</i> (Sclater, 1856)	30	s	c	a, i	x	u	a, o	barranqueiro-pardo
<i>Xenops milleri</i> (Chapman, 1914)	11b	s	c	a, i	x	u	a, o	bico-virado-da-copa
<i>Xenops minutus</i> (Sparman, 1788)	14	s	c	a, i	x	u	a, o	bico-virado-miúdo
TYRANNIDAE								
<i>Mionectes oleagineus</i> (Lichtenstein, 1823)	10	s	u	a, i, fr, s	x	u	o	abre-asa
<i>Mionectes macconnelli</i> (Chubb, 1919)	15	s	c	a, i, fr, s	x	u	a, o	abre-asa-da-mata
<i>Corythopsis torquatus</i> (Tschudi, 1844)	22	s	u	a, i, fr, s	x	g, u	o	estalador-do-norte
<i>Lophotriccus villosus</i> (Bangs & Penard, 1921)	—	s	c	a, i, fr, s		c, u	a	maria-fiteira
<i>Lophotriccus galeatus</i> (Boddaert, 1783)	7c	s	r	a, i, fr, s		c, u		caga-sebino-de-penacho
<i>Hemitriccus zosterops</i> (Pelzel, 1868)	8c	s	c	a, i, fr, s		c, u	a	maria-de-olho-branco
<i>Myiornis ecaudatus</i> (d'Orbigny & Lafresnaye, 1837)	4c	s	u	a, i, fr, s		u	a	caçula
<i>Todirostrum pictum</i> (Salvin, 1897)	7c	s	c	a, i, fr, s		c, u	a	ferreirinho-de-sobrancelha
<i>Tyrannulus elatus</i> (Latham, 1790)	7c	s	c	a, i, fr, s		c	a	maria-te-viú
<i>Myiopagis gaimardii</i> (d'Orbigny, 1839)	12c	s	c	a, i, fr, s	x	c	a	maria-pechin
<i>Myiopagis caniceps</i> (Swainson, 1835)	10c	s	c	a, i, fr, s		c	a	guaracava-cinzenta
<i>Elaenia parvirostris</i> (Pelzel, 1868)	15c	s	u	a, i, fr, s		c	a	guaracava-de-bico-curto
<i>Orniithion inerme</i> (Hartlaub, 1853)	6c	s	u	a, i, fr, s		c	a	poiaieiro-de-sobrancelha
<i>Camptostoma obsoletum</i> (Temminck, 1824)	7c	s	u	a, i, fr, s	x	c	o	risadinha
<i>Phaeomyias murina</i> (Spix, 1825)	10c	s	r	a, i, fr, s	x	c	o	bagageiro
<i>Zimmerius gracilipes</i> (Sclater & Salvin, 1868)	7c	s	c	a, i, fr, s		c	a	poiaieiro-de-pata-fina
<i>Phylloscartes virescens</i> (Todd, 1925)	—	s	c	a, i, fr, s		c	a	borboletinha-guianense
<i>Rhynchocylus olivaceus</i> (Temminck, 1820)	20	s	u	a, i, fr, s	x	c, u	o	bico-chato-grande

TABLE 4: Continued.

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
<i>Tolmomyias assimilis</i> (Pelzeln, 1868)	13b	s	c	a, i, fr, s	x	u	a, o	bico-chato-da-copa
<i>Tolmomyias poliocephalus</i> (Taczanowski, 1884)	11c	s	c	a, i, fr, s		u	a	bico-chato-de-cabeça-cinza
<i>Playrinchus saturatus</i> (Salvin & Godman, 1882)	12	s	u	a, i, fr, s	x	u	o	patinho-escuro
<i>Playrinchus coronatus</i> (Sclater, 1858)	10	s	c	a, i, fr, s	x	u	o	patinho-de-coroa-dourada
<i>Playrinchus platyrhynchos</i> (Gmelin, 1788)	11c	s	u	a, i, fr, s		u		patinho-de-coroa-branca
<i>Onychorhynchus coronatus</i> (Statius Muller, 1776)	15	s	u	a, i, fr, s	x	u	o	maria-leque
<i>Myiobius barbatus</i> (Gmelin, 1789)	10b	s	c	a, i, fr, s	x	u	a, o	assanhadinho
<i>Terenotriccus erythrurus</i> (Cabanis, 1847)	12	s	c	a, i, fr, s	x	u	a, o	papa-moscas-uirapuru
<i>Neopipo cinnamomea</i> (Lawrence, 1869)	7b	s	u	a, i, fr, s		u		enferrujadinho
<i>Contopus cooperi</i> (Nuttall, 1831)	32c	s	b	a, i, fr, s		c		piui-boreal
<i>Contopus virens</i> (Linnaeus, 1766)	14c	s	b	a, i, fr, s		c		piui-verdadeiro
<i>Pyrocephalus rubinus</i> (Boddaert, 1783)	14c	s	a	a, i, fr, s		c		príncipe
<i>Legatus leucophaeus</i> (Vieillot, 1818)	23c	s	u	a, i, fr, s	x	c		bem-te-vi-pirata
<i>Myiozetetes cayanensis</i> (Linnaeus, 1766)	26c	s	c	a, i, fr, s	x	c, u	a	bentevizinho-de-asa-ferrugínea
<i>Pitangus sulphuratus</i> (Linnaeus, 1766)	63c	s	c	a, i, fr, fo, fr, s	x	c, u		bem-te-vi
<i>Conopias parvus</i> (Pelzeln, 1868)	21c	s	c	a, i, fr, s	x	c	a	bem-te-vi-da-copa
<i>Myiodynastes maculatus</i> (Statius Muller, 1776)	43c	s	a	a, i, fr, s		c		bem-te-vi-rajado
<i>Megarynchus pitangua</i> (Linnaeus, 1766)	62c	s	u	a, i, fr, s	x	c	o	neinei
<i>Tyrannopsis sulphurea</i> (Spix, 1825)	54c	s	u	a, i, fr, s	x	c	o	suiriri-de-garganta-rajada
<i>Empidonomus varius</i> (Vieillot, 1818)	25c	s	u	a, i, fr, s	x	c		petica
<i>Griseotyrannus aurantioatrocristatus</i> (d'Orbigny & Lafresnaye, 1837)	27c	s	a	a, i, fr, s		c		petica-de-chapéu-preto
<i>Tyrannus melancholicus</i> (Vieillot, 1819)	39c	s	c	a, i, fr, s	x	c		suiriri
<i>Tyrannus savana</i> (Vieillot, 1808)	42c	s	a	a, i, fr, s		c		tesourinha
<i>Rhytipterna simplex</i> (Lichtenstein, 1823)	32	s	c	a, i, fr, s		c	a, o	vissíá
<i>Syrstes sibilator</i> (Vieillot, 1818)	32c	s	c	a, i, fr, s	x	c	a, o	gritador
<i>Myiarchus tuberculifer</i> (d'Orbigny & Lafresnaye, 1837)	19c	s	u	a, i, fr, s	x	u		maria-cavaleira-pequena
<i>Myiarchus ferox</i> (Gmelin, 1789)	24c	s	u	a, i, fr, s	x	u	o	maria-cavaleira
<i>Ramphotrigon ruficauda</i> (Spix, 1825)	19b	s	r	a, i, fr, s	x	c, u	a, o	bico-chato-de-rabo-vermelho

TABLE 4: Continued.

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
<i>Attila spadiceus</i> (Gmelin, 1789)	38	s	c	a, i, fr, s	x	u	a, o	capitão-de-saíra-amarelo
COTINGIDAE								
<i>Phoenicircus carnifex</i> (Linnaeus, 1758)	51	s	u	a, i, fr, s	x	c, u		saurá
<i>Cotinga cotinga</i> (Linnaeus, 1766)	54c	s	r	a, i, fr, s		c		anambé-de-peito-roxo
<i>Cotinga cayana</i> (Linnaeus, 1766)	65c	s	r	a, i, fr, s		c		anambé-azul
<i>Lipaugus vociferans</i> (Wied, 1820)	70	s	c	a, i, fr, s	x	u	a, o	cricrió
<i>Xipholena punicea</i> (Pallas, 1764)	65c	s	u	a, i, fr, s	x	c	a	anambé-pompadora
<i>Haematoderus militaris</i> (Shaw, 1792)	—	s	r	a, i, fr, s		c		anambé-militar
<i>Perissocephalus tricolor</i> (Statius Muller, 1776)	340c	s	u	a, i, fr, s	x	c	a	maú
PIPRIDAE								
<i>Neopelma chrysocephalum</i> (Pelzelin, 1868)	10	s	u	a, i, fr, s	x	u	o	fruchu-do-carrasco
<i>Tyrannetes virescens</i> (Pelzelin, 1868)	7b	s	c	a, i, fr, s	x	c	a, o	uirapuruzinho-do-norte
<i>Piprites chloris</i> (Temminck, 1822)	17b	s	c	a, i, fr, s	x	u	a, o	papinho-amarelo
<i>Corapipo gutturalis</i> (Linnaeus, 1766)	12	s	c	a, i, fr, s	x	u	o	dancarino-de-garganta-branca
<i>Lepidothrix serena</i> (Linnaeus, 1766)	12	s	c	a, i, fr, s	x	u	a, o	uirapuru-estrela
<i>Manacus manacus</i> (Linnaeus, 1766)	16c	s	u	a, i, fr, s	x	u		rendeira
<i>Dixiphia pipra</i> (Linnaeus, 1758)	11	s	c	a, i, fr, s	x	u	a, o	cabeça-branca
<i>Pipra erythrocephala</i> (Linnaeus, 1758)	12	s	c	a, i, fr, s	x	u	a, o	cabeça-de-ouro
TITYRIDAE								
<i>Schiffornis turdina</i> (Wied, 1831)	36	s	c	a, i, fr, s	x	u	a, o	flautim-marrom
<i>Laniocera hypopyrra</i> (Vieillot, 1817)	49b	s	c	a, i, fr, s		c		chorona-cinza
<i>Tityra cayana</i> (Linnaeus, 1766)	69c	s	c	a, i, fr, s	x	c	a	anambé branco-de-rabo-preto

TABLE 4: Continued.

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
<i>Pachyrhamphus rufus</i> (Boddaert, 1783)	18c	s	u	a, i, fr, s	x	c	o	caneleiro-cinzentio
<i>Pachyrhamphus marginatus</i> (Lichtenstein, 1823)	18b	s	c	a, i, fr, s		c	a	caneleiro-bordado
<i>Pachyrhamphus surinamus</i> (Linnaeus, 1766)	20c	s	c	a, i, fr, s		c		caneleiro-da-guiana
<i>Pachyrhamphus minor</i> (Lesson, 1830)	—	s	u	a, i, fr, s	x	c	a	caneleiro-pequeno
VIREONIDAE								
<i>Cyclarhis guianensis</i> (Gmelin, 1789)	28c	s	c	a, fr, s	x	c	a	pitiguari
<i>Vireolanius leucotis</i> (Swainson, 1838)	25c	s	c	a, fr, s	x	c	a	assobiador-do-castanhal
<i>Vireo olivaceus</i> (Linnaeus, 1766)	15c	s	b	a, fr, s		c	a	juruviara
<i>Vireo altiloquus</i> (Vieillot, 1808)	19c	s	b	a, fr, s		c		juruviara-barbuda
<i>Hylophilus muscicapinus</i> (Sclater & Salvin, 1873)	10b	s	c	a, fr, s	x	c	a	vite-vite-camurça
<i>Hylophilus ochraceiceps</i> (Sclater, 1860)	10	s	c	a, fr, s	x	c	a, o	vite-vite-uirapuru
HIRUNINIDAE								
<i>Atticora tibialis</i> (Cassin, 1853)	—	g	u	i		a		calcinha-branca
<i>Stelgidopteryx ruficollis</i> (Vieillot, 1817)	15c	s	u	i		a		andorinha-serradora
<i>Progne subis</i> (Linnaeus, 1758)	47c	g	b	i		a		andorinha-azul
<i>Progne chalybea</i> (Gmelin, 1789)	39c	g	u	i		a		andorinha-doméstica-grande
<i>Hirundo rustica</i> (Linnaeus, 1758)	17c	g	b	i		a		andorinha-de-bando
TROGLODYTIDAE								
<i>Microcerculus bambla</i> (Boddaert, 1783)	14	s	c	a, i	x	g, u	a, o	uirapuru-de-asa-branca
<i>Troglodytes musculus</i> (Naumann, 1823)	11b	s	c	a, i	x	g, u		corruíra
<i>Pheugopedius coraya</i> (Gmelin, 1789)	15	c	c	a, i	x	g, u	a, o	garrinchão-coraia
<i>Cyphorhinus arada</i> (Hermann, 1783)	18	c	u	a, i	x	g, u	o	uirapuru-verdadeiro
POLIOPTILIDAE								
<i>Microbates collaris</i> (Pelzeln, 1868)	11b	s	c	a, i	x	g, u	a, o	bico-assovelado-de-coleira
<i>Ramphocaenus melanurus</i> (Vieillot, 1819)	9b	s	c	a, i		g, u		bico-assovelado

TABLE 4: Continued.

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
<i>Poliophtila guianensis</i> (Todd, 1920)	5c	c	u	a, i		g, u		balança-rabo-guianense
TURDIDAE								
<i>Catharus fuscus</i> (Stephens, 1817)	28b	s*	b	a, i, fr, s		u		sabiá-norte-americano
<i>Catharus minimus</i> (Lafresnaye, 1848)	30b	s*	b	a, i, fr, s		u		sabiá-de-cara-cinza
<i>Turdus leucomelas</i> (Vieillot, 1818)	62c	s	u	a, i, fr, s	x	u		sabiá-barranco
<i>Turdus albicollis</i> (Vieillot, 1818)	50	s*	c	a, i, fr, s	x	u	a, o	sabiá-coleira
COEREBIDAE								
<i>Coereba flaveola</i> (Linnaeus, 1758)	9b	s	c	n, fr, s	x	c	a	cambacica
THRAUPIDAE								
<i>Salpator grossus</i> (Linnaeus, 1766)	44	s	u	s, fr	x	c	a	bico-encarnado
<i>Salpator maximus</i> (Statius Muller, 1776)	42	s	u	s, fr	x	c		tempera-viola
<i>Lamprospiza melanoleuca</i> (Vieillot, 1817)	—	s	c	a, i, n, fr, s		c	a	pipira-de-bico-vermelho
<i>Eucometis penicillata</i> (Spix, 1825)	29c	s	u	a, i, n, fr, s		c, u		pipira-da-taoca
<i>Tachyphonus cristatus</i> (Linnaeus, 1766)	18b	s	c	a, i, n, fr, s		c	a	tiê-galo
<i>Tachyphonus surinamus</i> (Linnaeus, 1766)	19	s	c	a, i, n, fr, s	x	c, u	a, o	tem-tem-de-topete-ferrugíneo
<i>Lanio fulvus</i> (Boddaert, 1783)	22	s	u	a, i, n, fr, s		c	a, o	pipira-parda
<i>Ramphocelus carbo</i> (Pallas, 1764)	24b	g	c	a, i, n, fr, s	x	u		pipira-vermelha
<i>Thraupis episcopus</i> (Linnaeus, 1766)	35c	c	c	a, i, n, fr, s	x	c, u		sanhaçu-da-amazônia
<i>Thraupis palmarum</i> (Wied, 1823)	35b	c	c	a, i, n, fr, s	x	c, u	a	sanhaçu-do-coqueiro
<i>Gyanicterus cyanicterus</i> (Vieillot, 1819)	34c	s	r	a, i, n, fr, s		c	a	pipira-azul
<i>Tangara mexicana</i> (Linnaeus, 1766)	20c	c	r	a, i, n, fr, s	x	c		saiira-de-bando
<i>Tangara chilensis</i> (Vigors, 1832)	15b	c	c	a, i, n, fr, s		c		sete-cores-da-amazônia
<i>Tangara punctata</i> (Linnaeus, 1766)	14b	c	c	a, i, n, fr, s		c	a	saiira-negaça
<i>Tangara varia</i> (Statius Muller, 1776)	10c	c	r	a, i, n, fr, s		c, u	a	saiira-carijó
<i>Tangara gyrola</i> (Linnaeus, 1758)	19c	c	r	a, i, n, fr, s		c		saiira-de-cabeça-castanha
<i>Tangara velia</i> (Linnaeus, 1758)	20c	c	c	a, i, n, fr, s		c	a	saiira-diamante
<i>Dacnis lineata</i> (Gmelin, 1789)	—	c	c	a, i, n, fr, s	x	c, u	a, o	sai-de-máscara-preta
<i>Dacnis cayana</i> (Linnaeus, 1766)	12	c	c	a, i, n, fr, s	x	c, u	a, o	sai-azul
<i>Chlorophanes spiza</i> (Linnaeus, 1758)	16b	c	c	a, i, n, fr, s		c, u	a	sai-verde
<i>Gyanerpes nitiidus</i> (Hartlaub, 1847)	9c	c	u	a, i, n, fr, s		c		sai-de-bico-curto
<i>Gyanerpes caeruleus</i> (Linnaeus, 1758)	12c	c	c	a, i, n, fr, s		c		sai-de-perna-amarela
<i>Gyanerpes cyaneus</i> (Linnaeus, 1766)	14c	c	c	a, i, n, fr, s		c		sai-beija-flor
<i>Hemithraupis flavicollis</i> (Vieillot, 1818)	12c	s	c	a, i, n, fr, s		c	a	saiira-galega
EMBERIZIDAE								
<i>Ammodramus aurifrons</i> (Spix, 1825)	14b	s	c	s, fr		g, u		cigarrinha-do-campo
<i>Volatinia jacarina</i> (Linnaeus, 1766)	9c	s	c	s, fr		u		tiziu

TABLE 4: Continued.

Bird species	Wt. (g)	Soc.	Abund.	Diet	Br.	Str.	Rec.	Portuguese name
<i>Sporophila castaneiventris</i> (Cabanis, 1849)	8c	s	c	s, fr	x	u		caboclinho-de-peito-castanho
<i>Sporophila angolensis</i> (Linnaeus, 1766)	13b	s	u	s, fr	x	u		curio
<i>Arremon taciturnus</i> (Hermann, 1783)	22b	s	r	s, fr	x	u	a, o	tico-tico-de-bico-preto
CARDINALIIDAE								
<i>Pranga rubra</i> (Linnaeus, 1758)	28c	c	b	a, i, n, fr, s		c		sanhaçu-vermelho
<i>Caryothraustes canadensis</i> (Linnaeus, 1766)	33b	s	c	s, fr		c	a	furriel
<i>Cyanoxia cyanooides</i> (Lafresnaye, 1847)	28	s	c	s, fr	x	u	a, o	azulão-da-amazônia
PARULIDAE								
<i>Dendroica striata</i> (Forster, 1772)	10c	s	b	a, i		c		mariquita-de-perna-clara
<i>Phaeothlypis rivularis</i> (Wied, 1821)	12c	s	u	a, i		c		pula-pula-ribeirinho
ICTERIDAE								
<i>Psarocolius viridis</i> (Statius Muller, 1776)	375c	g	u	fr, i, s	x	c		japu-verde
<i>Cacicus haemorrhous</i> (Linnaeus, 1766)	100c	g	c	fr, i, s	x	c	a	guaxe
<i>Cacicus cela</i> (Linnaeus, 1776)	104c	g	u	fr, i, s	x	c	a	xexéu, japiim
<i>Icterus chryscephalus</i> (Linnaeus, 1766)	42c	s	u	fr, i, s	x	c		rouxinal-do-río-negro
<i>Sturnella militaris</i> (Linnaeus, 1758)	44c	s	u	fr, i, s		c		polícia-inglesa-do-norte
<i>Molothrus oryzivorus</i> (Gmelin, 1788)	180c	g	u	fr, i, s		c		irauna-grande
<i>Molothrus bonariensis</i> (Gmelin, 1789)	45c	g	u	fr, i, s		c		vira-bosta
FRINGILLIDAE								
<i>Euphonia chlorotica</i> (Linnaeus, 1766)	11c	s	u	a, i, n, fr, s	x	c		fim-fim
<i>Euphonia chrysopasta</i> (Sclater & Salvin, 1869)	14c	s	u	a, i, n, fr, s	x	c		gaturamo-verde
<i>Euphonia minuta</i> (Cabanis, 1849)	10c	s	u	a, i, n, fr, s	x	c		gaturamo-de-barriga-branca
<i>Euphonia cayennensis</i> (Gmelin, 1789)	13c	s	c	a, i, n, fr, s		c	a	gaturamo-preto

Codes: Wt.: body weight; b: mean value rounded from [52]; c: mean value rounded from [53]; Diet: a: arthropods (spiders, centipedes, millipedes, etc); b: birds; ba = bats; f: flowers; fi: fish; fr: fruits; fo: frogs; i: insects; l: leaf; li: lizards; mo: monkeys; ne: nectar; r: rodents; s: seeds; sl: sloths. Str.: Forest stratum: c: canopy; g: ground; u: understory; Soc.: Sociability: c: pairs; g: groups; s: solitary (*: army-ant follower); Abund.: Abundance: c: common, u: uncommon, r: rare, a: austral migrant, b: boreal migrant; Br.: Breeding: x: nests, eggs, nestling or juveniles recorded in the Reserve; Rec.: Record (this study); a: acoustic surveys along 1 km transects; o: mist-nets (see Section 2). Species not recorded during this study but included in the table (see Rec.) were recorded by the authors at the RFAD in the past. These species were not included in the analyses. Data on weight, sociability, abundance, diet, breeding, and strata were collected by RC during the last 20 years.

Acknowledgments

We thank Obed Barros, Sidnei Dantas, Adrianny Maruoka, Karina Amaral, Marlison Ferreira, Lucas Mergulhão and Francisco Marques Bezerra for their competent field assistance with the mist-nets and measurements of forest structure components. Pedro Santos and S. V. Wilson helped with comments in earlier versions. Victor Landeiro, Cintia Gomes, and Cristian Dambros helped with construction of figures on gradients using the program "R". William Magnusson and Charles R. Clement helped editing the final English version. The paper benefited greatly from comments by two anonymous reviewers and by B. A. Hawkins (Editor) on the manuscript. This study was supported by PNOPg-CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) and INPA (Instituto Nacional de Pesquisas da Amazônia). We thank Tito Fernandes for his help with CorelDRAW program.

References

- [1] R. MacArthur, H. Recher, and M. Cody, "On the relation between habitat selection and species diversity," *American Naturalist*, vol. 100, pp. 319–332, 1966.
- [2] J. Terborgh, S. K. Robinson, T. A. Parker III, C. A. Munn, and N. Pierpont, "Structure and organization of an Amazonian forest bird community," *Ecological Monographs*, vol. 60, no. 2, pp. 213–238, 1990.
- [3] J. R. Karr and K. E. Freemark, "Habitat selection and environmental gradients: dynamics in the 'stable' tropics," *Ecology*, vol. 64, no. 6, pp. 1481–1494, 1983.
- [4] J. Terborgh, "Habitat selection in Amazonian birds," in *Habitat Selection in Birds*, M. L. Cody, Ed., pp. 331–340, Academic Press, New York, NY, USA, 1985.
- [5] J. M. Thiollay, "Influence of selective logging on bird species diversity in a Guianan rain forest," *Conservation Biology*, vol. 6, no. 1, pp. 47–63, 1992.
- [6] R. Cintra and T. M. Sanaiotti, "Fire effects on the composition of a bird community in an Amazonian savanna (Brazil)," *Brazilian Journal of Biology*, vol. 65, no. 4, pp. 683–695, 2005.
- [7] R. Cintra, A. E. Maruoka, and L. N. Naka, "Abundance of two *Dendrocicla* woodcreepers (Aves: Dendrocolaptidae) in relation to forest structure in Central Amazonia," *Acta Amazonica*, vol. 36, no. 2, pp. 209–219, 2006.
- [8] B. A. Loiselle and J. G. Blake, "Temporal variation in birds and fruits along an elevational gradient in Costa Rica," *Ecology*, vol. 72, no. 1, pp. 180–193, 1991.
- [9] P. B. Pearman, "The scale of community structure: habitat variation and avian guilds in tropical forest understory," *Ecological Monographs*, vol. 72, no. 1, pp. 19–39, 2002.
- [10] D. L. Pearson, "The relationship of foliage complexity to ecological diversity of three Amazonian bird communities," *Condor*, vol. 77, pp. 453–466, 1975.
- [11] R. Cintra, "Spatial distribution and foraging tactics of tyrant flycatchers in two habitats in the Brazilian Amazon," *Studies on Neotropical Fauna and Environment*, vol. 32, no. 1, pp. 17–27, 1997.
- [12] R. O. Bierregaard, "Avian communities in the understory of Amazonian forest fragments," in *Biogeography and Ecology of Forest Bird Communities*, A. Keast, Ed., pp. 333–343, SPB Academic Publishing, The Hague, The Netherlands, 1990.
- [13] E. Guilherme and R. Cintra, "Effects of intensity and age of selective logging and tree girdling on an understory bird community composition in central Amazonia, Brazil," *Ecotropica*, vol. 7, pp. 77–92, 2001.
- [14] M. Cohn-Haft, A. Whittaker, and P. C. Stouffer, "A new look at the 'species-poor' Central Amazon: the avifauna north of Manaus, Brazil," *Ornithological Monographs*, vol. 48, pp. 205–235, 1997.
- [15] P. C. Stouffer and R. O. Bierregaard, "Use of Amazonian forest fragments by understory insectivorous birds," *Ecology*, vol. 76, no. 8, pp. 2429–2445, 1995.
- [16] D. W. Snow and A. Lill, "Longevity records for some Neotropical land birds," *Condor*, vol. 76, pp. 262–267, 1974.
- [17] C. Banks and R. Cintra, "The heterogeneity of Amazonian treefall gaps and bird community composition," *Ecotropica*, vol. 14, pp. 1–13, 2008.
- [18] A. Aleixo, "Effects of selective logging on a bird community in the Brazilian Atlantic forest," *Condor*, vol. 101, no. 3, pp. 537–548, 1999.
- [19] J. del Hoyo, A. Elliott, and J. Sargatal, *Handbook of the Birds of the World*, vol. 1–16, Lynx Edicions, Barcelona, Spain, 1992–2011.
- [20] J. Adis, "On the abundance and density of terrestrial arthropods in Central Amazonian dryland forests," *Journal of Tropical Ecology*, vol. 4, no. 1, pp. 19–24, 1988.
- [21] C. V. de Castilho, W. E. Magnusson, R. N. O. de Araújo et al., "Variation in aboveground tree live biomass in a central Amazonian Forest: effects of soil and topography," *Forest Ecology and Management*, vol. 234, no. 1–3, pp. 85–96, 2006.
- [22] J. E. L. S. Ribeiro, M. J. G. Hopkins, A. Vincentini et al., *Flora da Reserva Ducke—Guia de Identificação das Plantas Vasculares de uma Floresta de Terra-Firme na Amazônia Central*, Instituto Nacional de Pesquisas da Amazônia, Manaus, Amazonas, Brazil, 1999.
- [23] J. R. Karr, "Surveying birds with mist nets," *Studies in Avian Biology*, vol. 6, pp. 62–67, 1981.
- [24] B. McCune and M. J. Mefford, *Multivariate Analysis of Ecological Data, Version 4.25*, MjM Software, Gleneden Beach, Okla, USA, 1999.
- [25] P. R. Minchin, "An evaluation of the relative robustness of techniques for ecological ordination," *Vegetatio*, vol. 69, no. 1–3, pp. 89–107, 1987.
- [26] M. J. Anderson and T. J. Willis, "Canonical analysis of principal coordinates: a useful method of constrained ordination for ecology," *Ecology*, vol. 84, no. 2, pp. 511–525, 2003.
- [27] P. Legendre and L. Legendre, *Numerical Ecology*, Elsevier Press, Amsterdam, Netherlands, 1998.
- [28] R. C. Mac Nally, "On characterizing foraging versatility, illustrated by using birds," *Oikos*, vol. 69, no. 1, pp. 95–106, 1994.
- [29] F. R. C. Costa and W. E. Magnusson, "Selective logging effects on abundance, diversity, and composition of tropical understory herbs," *Ecological Applications*, vol. 12, no. 3, pp. 807–819, 2002.
- [30] A. P. Lima, A. C. Cordeiro-Duarte, F. J. Luizão, and N. Higuchi, "Effect of selective logging intensity on two termite species of the genus *Syntermes* in Central Amazonia," *Forest Ecology and Management*, vol. 137, no. 1–3, pp. 151–154, 2000.
- [31] R. Cintra, T. M. Sanaiotti, and M. Cohn-Haft, "Spatial distribution and habitat of the Anavilhanas Archipelago bird community in the Brazilian Amazon," *Biodiversity and Conservation*, vol. 16, no. 2, pp. 313–336, 2007.

- [32] R. Cintra, P. M. R. S. Dos Santos, and C. B. Leite, "Composition and structure of the lacustrine bird communities of seasonally flooded wetlands of Western Brazilian Amazonia at high water," *Waterbirds*, vol. 30, no. 4, pp. 521–540, 2007.
- [33] C. R. Johnson and C. A. Field, "Using fixed-effects model multivariate analysis of variance in marine biology and ecology," *Oceanography and Marine Biology*, vol. 31, pp. 177–221, 1993.
- [34] C. L. Olson, "On choosing a test statistic in multivariate analysis of variance," *Psychological Bulletin*, vol. 83, no. 4, pp. 579–586, 1976.
- [35] L. Wilkinson, *Systat: The System for Statistics, Version 12*, SYSTAT Software, Evanston, Ill, USA, 2007.
- [36] J. Fox, *An R and S-Plus Companion to Applied Regression*, SAGE Publications, London, UK, 2002.
- [37] E. O. Willis, "Preliminar list of the birds from western part and neighboring areas of Ducke Reserve, Amazonas, Brazil," *Revista Brasileira de Biologia*, vol. 37, pp. 585–601, 1977.
- [38] O. G. Barros and R. Cintra, "The effects of forest structure on occurrence and abundance of three owl species (Aves: Strigidae) in the Central Amazon forest," *Zoologia*, vol. 26, no. 1, pp. 85–96, 2009.
- [39] A. Chauvel, Y. Lucas, and R. Boulet, "On the genesis of the soil mantle of the region of Manaus, Central Amazonia, Brazil," *Experientia*, vol. 43, no. 3, pp. 234–241, 1987.
- [40] V. F. Kinupp and W. E. Magnusson, "Spatial patterns in the understory shrub genus *Psychotria* in central Amazonia: effects of distance and topography," *Journal of Tropical Ecology*, vol. 21, no. 4, pp. 363–374, 2005.
- [41] F. R. C. Costa, W. E. Magnusson, and R. C. Luizão, "Mesoscale distribution patterns of Amazonian understory herbs in relation to topography, soil and watersheds," *Journal of Ecology*, vol. 93, no. 5, pp. 863–878, 2005.
- [42] F. R. C. Costa, J. L. Guillaumet, A. P. Lima, and O. S. Pereira, "Gradients within gradients: the mesoscale distribution patterns of palms in a central amazonian forest," *Journal of Vegetation Science*, vol. 20, no. 1, pp. 69–78, 2009.
- [43] P. Y. de Oliveira, J. L. P. de Souza, F. B. Baccaro, and E. Franklin, "Ant species distribution along a topographic gradient in a "terra-firme" forest reserve in Central Amazonia," *Pesquisa Agropecuaria Brasileira*, vol. 44, no. 8, pp. 852–860, 2009.
- [44] M. Menin, A. P. Lima, W. E. Magnusson, and F. Waldez, "Topographic and edaphic effects on the distribution of terrestrially reproducing anurans in Central Amazonia: mesoscale spatial patterns," *Journal of Tropical Ecology*, vol. 23, no. 5, pp. 539–547, 2007.
- [45] F. P. Mendonça, W. E. Magnusson, and J. Zuanon, "Relationships between habitat characteristics and fish assemblages in small streams of Central Amazonia," *Copeia*, vol. 2005, no. 4, pp. 751–764, 2005.
- [46] J. M. Thiollay, "Structure, density and rarity in an Amazonian rainforest bird community," *Journal of Tropical Ecology*, vol. 10, no. 4, pp. 449–481, 1994.
- [47] G. Ferraz, J. D. Nichols, J. E. Hines, P. C. Stouffer, R. O. Bierregaard, and T. E. Lovejoy, "A large-scale deforestation experiment: effects of patch area and isolation on Amazon birds," *Science*, vol. 315, no. 5809, pp. 238–241, 2007.
- [48] L. N. Naka, "Structure and organization of canopy bird assemblages in Central Amazonia," *Auk*, vol. 121, no. 1, pp. 88–102, 2004.
- [49] W. F. Laurance, P. M. Fearnside, S. G. Laurance et al., "Relationship between soils and Amazon forest biomass: a landscape-scale study," *Forest Ecology and Management*, vol. 118, no. 1–3, pp. 127–138, 1999.
- [50] D. P. Drucker, F. R. C. Costa, and W. E. Magnusson, "How wide is the riparian zone of small streams in tropical forests? A test with terrestrial herbs," *Journal of Tropical Ecology*, vol. 24, no. 1, pp. 65–74, 2008.
- [51] *Lista das aves do Brasil. Comitê Brasileiro de Registros Ornitológicos (CBRO)*, <http://www.cbro.org.br/CBRO/listabr.htm>.
- [52] R. O. Bierregaard, "Morphological data from understory birds in terra firme forest in the Central Amazon basin," *Revista Brasileira de Biologia*, vol. 48, pp. 169–178, 1988.
- [53] S. L. Hilty, *Birds of Venezuela*, Princeton University Press, Princeton, NJ, USA, 2003.
- [54] A. S. Bueno, R. S. Bruno, T. P. Pimentel, T. M. Sanaiotti, and W. E. Magnusson, "The width of riparian habitats for understory birds in an Amazonian forest," *Ecological Applications*. In press.