



UNIVERSIDADE FEDERAL DE MATO GROSSO
INSTITUTO DE BIOCÊNCIAS
PROGRAMA DE PÓS-GRADUAÇÃO EM ECOLOGIA E CONSERVAÇÃO DA
BIODIVERSIDADE

**STRUCTURE OF COMPLEX NETWORKS AND THE ROLE OF SPATIAL
VARIATIONS AND ABUNDANCE IN PLANT-ANT INTERACTIONS**

WESLEY FRANCISCO DÁTILLO DA CRUZ

CUIABÁ – MATO GROSSO

March 2012



UNIVERSIDADE FEDERAL DE MATO GROSSO
INSTITUTO DE BIOCÊNCIAS
PROGRAMA DE PÓS-GRADUAÇÃO EM ECOLOGIA E CONSERVAÇÃO DA
BIODIVERSIDADE

**STRUCTURE OF COMPLEX NETWORKS AND THE ROLE OF SPATIAL
VARIATIONS AND ABUNDANCE IN PLANT-ANT INTERACTIONS**

WESLEY FRANCISCO DÁTILLO DA CRUZ

Dissertação apresentada ao Programa de
Pós- Graduação, do Instituto de Biociências,
para obtenção do título de Mestre em
Ecologia e Conservação da Biodiversidade.

CUIABÁ – MATO GROSSO

March 2012

Cruz, Wesley Francisco Dáttilo da.

S586s

Structure of complex networks and the role of spatial variations and abundance in plant-ant interactions. / Wesley Francisco Dáttilo da Cruz. Cuiabá, Mato Grosso, Brazil, 2012.

99f.: Il.

Dissertação (Mestrado em Ecologia e Conservação da Biodiversidade) – Programa de Pós-Graduação em Ciências Biológicas, Instituto de Biociências, Universidade Federal de Mato Grosso, Cuiabá, Mato Grosso, 2011.

Orientador: Prof. Dr. Thiago Junqueira Izzo.

Co-orientador: Prof. Dr. Paulo Roberto Guimarães Júnior.

1. Aninhamento. 2. Neutralidade. 3. Coevolução. 4. Interações planta-animal. 5. Substituição espacial. I. Título. 574

CDU: 574 (817.2)

ADVISOR: Dr. Thiago Junqueira Izzo

Departamento de Ecologia e Botânica

Instituto de Biociências - Universidade Federal de Mato Grosso.

CO-ADVISOR: Dr. Paulo Roberto Guimarães Júnior


Departamento de Ecologia Geral

Instituto de Biociências - Universidade de São Paulo.

DISSERTATION COMMITTEE

Dr. Fernando Zagury Vaz de Mello

Universidade Federal de Mato Grosso – Departamento de Biologia e Zoologia,
Instituto de Biociências
(Internal Examiner)



Dr. Mário Almeida Neto

Universidade Federal de Goiás – Departamento de Ecologia
Instituto de Ciências Biológicas
(External Examiner)



Dr. Jerry Magno Ferreira Penha

Universidade Federal de Mato Grosso – Departamento de Ecologia e Botânica
Instituto de Biociências
(Internal Examiner)

Dr. Roberto de Moraes Lima Silveira

Universidade Federal de Mato Grosso – Departamento de Ecologia e Botânica
Instituto de Biociências
(Alternate Examiner)

I dedicate this dissertation to all those who supported me throughout this journey. In particular my parents and my wife, who were often deprived of attention, but always stayed by my side.

“Interactions between species are as evolutionarily malleable
as the species themselves and have played a central role in the
diversification and organization of life.”
(John N. Thompson 1999)

ACKNOWLEDGEMENTS

The work presented in this dissertation benefited from many people's comments, discussion, collaboration and encouragement.

First and foremost, I thank my advisor, Dr. Thiago J. Izzo, for their guidance, support and inspiration throughout. Besides an excellent researcher is a great friend I made in 2007 at the Instituto Nacional de Pesquisas da Amazônia (INPA) where I was his intern working with ant-plant symbiosis. Despite his busy schedule, Thiago Izzo always finds the time to discuss anything from intriguing experimental results to soccer. His dedication and commitment to science is truly inspiring and remarkable. I would also like to thank my co-advisor, Dr. Paulo R. Guimarães-Jr, I am very appreciative of his generosity with his time, advice, data analysis and all help with this dissertation. Thanks for also welcoming me into his lab at the Universidade de São Paulo (USP). Without his support, this project would not have been possible. Thiago Izzo and Paulo Guimarães both brought unique perspectives to my research, enriching it greatly. Their encouragement and enthusiasm were important for the completion of this project.

To Dra. Soraia Diniz and Dr. Roberto Silveira, I express my gratefulness for the discussion and interpretation of some results presented in this thesis. Many additional colleagues provided valuable information that helped my dissertation, and I am grateful to all of them for their assistance.

I cannot forget to thank my dissertation committee members for their valuable comments and constructive criticism; thank you Dr. Mário Almeida-Neto (Universidade Federal de Goiás), Dr. Jerry Penha (UFMT) and Dr. Fernando Vaz-de-Mello (UFMT).

I am grateful also to my true friends at the Universidade Federal de Mato Grosso who over the years have patiently endured my highs and lows and have kept me sane and entertained. In particular, I thank Fernando Tortato (Galdério), Mateus Souza (Ventania), Marcela Gigliotti (Marcelita), Miquéias Ferrão (Kiki), Rafael Nunes (Bino), Vitor Azarias (Zé Colméia). Sorry for something. I extend my heartfelt salute to all my colleagues at the Community Ecology Lab. of UFMT (Ricardo Vicente, Jéssica Falcão and Daniela Monteiro). It was an honour and a pleasure to work among folks of such commitment and fun.

My studies were funded by a scholarship from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), with additional funding from the

Brazilian Program for Biodiversity Research (PPBio) and the logistical support from the Office National des Forêts Brazil. I gratefully acknowledge this support.

Finally, I would like to express my deepest gratitude to Jéssica Falcão (my wife) for everything from technical and scientific support to emotional support. I could not have done it without you. Thank you for being there for me from the very beginning.

Thank you for believing in me.

SUMMARY

| | |
|------------------------|-------------|
| FIGURES..... | XI |
| APPENCIDES..... | XIII |
| ABSTRACT..... | XIV |
| RESUMO..... | XV |

| | |
|----------------------------------------------|-----------|
| CHAPTER 1 - General introduction..... | 17 |
| References..... | 21 |

| | |
|-----------------------------------------------------------------------------------------------|-----------|
| CHAPTER 2 - The structure of ant-plant mutualistic networks: is abundance enough?..... | 24 |
| Abstract..... | 26 |
| 1. Introduction..... | 27 |
| 2. Material and Methods..... | 30 |
| 2.1) <i>Study area</i> | 30 |
| 2.2) <i>Data Collection</i> | 30 |
| 2.3) <i>Network topology</i> | 32 |
| 2.4) <i>Overrepresentation of species interactions</i> | 33 |
| 2.5) <i>Additive partitioning of diversity</i> | 35 |
| 3. Results..... | 36 |
| 4. Discussion..... | 37 |
| Acknowledgements..... | 40 |
| Literature cited..... | 40 |

| | |
|------------------------------------------------------------------------------|-----------|
| CHAPTER 3 - Spatial structure of ant-plant mutualistic networks | 53 |
| Abstract..... | 55 |
| 1. Introduction..... | 56 |
| 2. Material and Methods..... | 58 |
| 2.1) <i>Study area</i> | 58 |
| 2.2) <i>Data Collection</i> | 58 |
| 2.3) <i>Network topology</i> | 59 |
| 2.4) <i>Spatial turnover of species composition</i> | 60 |
| 2.5) <i>Statistical analysis</i> | 61 |

| | |
|------------------------------|-----------|
| 3. Results..... | 61 |
| 4. Discussion..... | 62 |
| Acknowledgements..... | 65 |
| References..... | 66 |
| FINAL CONCLUSION..... | 74 |

FIGURES

Chapter 1

Figure 1. Main ways to view the structure of an interaction networks nested and asymmetric: A) bipartite graphs and B) ordered matrices. Both forms of representation are ordered according to interactions number of the species, where species that have fewer interactions are subsets of species with more interactions. Cells painted of gray in Figure 1B represent the core of generalists species (Figures modified from Lewinsohn et al. 2006). 19

Chapter 2

Figure 1. Relationship among the different metrics used and the type of interaction involved (mutualistic or neutral): A) mean number of links for plant specie; B) mean number of links for ant specie; C) connectance; D) modularity*; E) nestedness (NODF metric); F) nestedness (Z-Score metric). Each line represents one of the 12 plots paired with the type of interaction. It is also shown the boxplots with the distribution of the data set based on their descriptive parameters. Only the mean number of links for ant specie (B) was not significant (paired t-test: $t = 0.258$, $df = 11$, $P = 0.801$). * In any of the plots studied were observed modularity significantly higher than expected by the heterogeneity of interactions. 49

Figure 2. Average percentage of additive partitioning of total diversity of ants collected on plants with (mutualistic interactions) and without (neutral interactions) extra-floral nectaries in 12 plots of an *terra-firme* dense rainforest in the Brazilian Meridional Amazon. α -diversity represents the mean of ants richness collected on plants with and without extra-floral nectaries. β -diversity represent the turnover of ant species composition collected on plants with and without extra-floral nectaries in the plots. The total ants richness found in mutualistic and neutral networks in the plots was determined by γ -diversity. The values of the additive partitioning of diversity expected were calculated excluding the effect of the presence of EFNs through 1000 randomizations of plants classification (presence or absence of EFN). 51

Figure 3. Graphical representation of one of the 12 interaction matrices among ants with extra-floral nectaries (EFNs) plants studied. This matrix shows the probability of an interaction between ants and plants occur mediated by the relative species abundance of both plants and ants. The red cells represents that the interaction occurs less than expected by the abundance, blue cells represents the interaction occurs more than expected by the abundance, gray cells represents ant species collected in plants without EFNs was not collected in plants with EFNs. 52

Figure 4. Non-metric multidimensional scaling of ants collected in plants with (triangles) and without extrafloral nectaries (squares) in 12 plots of an *terra-firme* dense rainforest in the Brazilian Meridional Amazon. This ordination analysis was calculated from the (A) Sorensen's dissimilarity index (Stress= 0.351 ; Axis 1 + Axis 2= 34.9% of explanation) and (B) Bray-Curtis's dissimilarity index (Stress= 0.318 ; Axis 1 + Axis 2= 43.7% of explanation). 54

Chapter 3

Figure 1. Relationship among the dissimilarity of: a) connectance, b) network specialization, c) nestedness by NODF metric and d) nestedness by Z-Score metric, with dissimilarity of geographic distance of 12 plots collected in the São Nicolau Farm, Mato Grosso State, Brazilian Meridional Amazon. Correlation coefficient (r) and significance (P computed using Mantel tests) also are shown. (n= 66 points in each of the metrics calculated). 72

Figure 2. Relationship among the dissimilarity of: a) plants species composition, b) ants species composition, c) plants core composition and b) ants core composition, with dissimilarity of geographic distance of 12 plots collected in the São Nicolau Farm, Mato Grosso State, Brazilian Meridional Amazon. Correlation coefficient (r) and significance (P computed using Mantel tests) also are shown. (n= 66 points in each of the metrics calculated). 73

APPENDICES

APPENDIX 1. List of 238 plant species with and without extrafloral nectaries (EFNs) collected in 12 plots (250 x 25 m) in an terra-firme forest in the Brazilian Meridional Amazon located in the municipality of Cotriguaçu, north of Mato Grosso State, Brazil. The plants were collected between between December 2010 and January 2011. 75

APPENDIX 2. List of 149 ant species collected on plants with and without extrafloral nectaries (EFNs) collected in 12 plots (250 x 25 m) in an terra-firme forest in the Brazilian Meridional Amazon located in the municipality of Cotriguaçu, north of Mato Grosso State, Brazil. The plants were collected between between December 2010 and January 2011. ..84

APPENDIX 3. Instructions for Authors – *Ecology*..... 90

APPENDIX 4. Instructions for Authors – *Oikos*..... 93

ABSTRACT: The structural organization of mutualism networks provides insights into processes shaping biodiversity. Understanding the mechanisms that shape this organization is essential for us to understand the ecological and evolutionary dynamics of the interacting species. Recent studies have suggested that species abundance is one of the most fundamental criteria shaping mutualistic networks. However, the role of species abundance on mutualistic networks is still unclear. Moreover, only recently has the spatial effect been incorporated as a mechanism structuring mutualistic networks. In this work I studied plants with extrafloral nectaries and associated ants to show that the natural abundance of ants on vegetation explained just a part of the frequency of mutualistic interactions and that it is independent of ant species compositions. In addition, the generalist core of these networks interacts more among themselves than expected by their abundances. This generalist core was formed by competitively superior ants that have behavioral and ecophysiological traits to use liquid food and are that exclude other species from the same resource. I also showed that even the fact that ant and plant composition of networks changes over space, the generalist core species and the topological structure of networks remain unaltered. This finding indicates that independently of local and landscape environmental factors the nonrandom pattern of community organization is not changed. Such generalist core conformation being stable over space and time could have serious implications on coevolutionary process of the system. In short, contributing thus to our understanding of the maintenance of biodiversity.

Keywords: nestedness, neutrality, coevolution, plant-animal interactions, spatial turnover.

RESUMO: A organização estrutural das redes mutualísticas provê idéias sobre processos que modulam a biodiversidade. Conhecer os mecanismos que modulam essa organização é essencial para entendermos a dinâmica ecológica e evolutiva das espécies que interagem. Recentes estudos têm sugerido que a abundância das espécies é um dos principais critérios que modulam essas redes. Entretanto, o papel da abundância relativa das espécies nas redes mutualísticas ainda é incerto. Além disso, apenas recentemente o efeito espacial tem sido incorporado como mecanismo que estrutura redes mutualísticas. Neste trabalho, eu estudei plantas com nectários extra-florais para mostrar que a abundância natural das formigas sobre a vegetação explica apenas uma parte da frequência das interações mutualísticas independente da composição de espécies de formigas. Adicionalmente, o núcleo generalista dessas redes interage mais entre si do que esperado pelas suas abundâncias. Este núcleo generalista foi formado por formigas competitivamente superiores que tem características comportamentais e ecofisiológicas para utilizar alimentos líquidos e que excluem as demais espécies do mesmo recurso. Eu também mostrei que mesmo que a composição de formigas e plantas das redes muda ao longo do espaço, o núcleo de espécies generalistas e a estrutura topológica das redes permanecem inalterados. Esse resultado indica que independente de fatores locais ou ambientais o padrão não-aleatório da organização da comunidade não é mudado. Tal conformação do núcleo generalista sendo estável ao longo do espaço e do tempo poderia ter sérias implicações sobre os processos coevolutivos do sistema. Por fim, contribuindo dessa forma para nosso entendimento da manutenção da biodiversidade.

Palavras-chave: aninhamento, neutralidade, coevolução, interações planta-animal, substituição espacial.

CHAPTER 1

GENERAL INTRODUCTION

CHAPTER 1 - GENERAL INTRODUCTION

Most interspecific interactions involving multiple species, resulting in the interactions networks that may be beneficial, neutral or harmful among the individuals involved (Lewinsohn et al. 2006). Among the various forms of interactions, the study of mutualism in community level was overlooked for a long time (Stanton 2003). However, recently the study of cooperative interactions among species has become one of the central issues in the community ecology (Bronstein 2001; Thompson 2005; Bronstein 2006, Ferrière et al. 2007). Nevertheless, it is likely that all organisms on Earth are involved in at least one event of mutualistic interactions throughout their life history (Bronstein 2001; Toby-Kiers et al. 2010). In this type of interaction, a species provides a service or product that the partner can not achieve alone, and in exchange receive some type of reward (Janzen 1985; Hoeksema & Bruna 2000). This makes mutualistic interactions influences directly the ecological and evolutionary dynamics of interacting species in different ecosystems around the world (Bronstein 2001; Thompson 2005; Bascompte et al. 2006; Montoya et al. 2006; Guimarães et al. 2007). Thus, understand how these factors act on the interaction patterns is helpfull on the management and conservation of the interacting species (Janzen 1974; Burslem et al. 2005; Del-Claro & Torezan-Silingardi 2009; Dyer et al. 2010)

Recent studies have focused on the structure of mutualistic networks among free-living species. This studies found non-random patterns of interaction on different systems and habitats throughout the Earth (Bascompte et al. 2003, Thompson 2005; Lewinsohn et al. 2006; Guimarães et al. 2007; Vázquez 2009). The metrics used in the study of interaction networks are derived from graph theory, first proposed in 1735 by Swiss mathematician Leonhard Euler (Mello 2010). Among the main features found in such networks is that they are highly nested and exhibit asymmetrical pattern of interactions. This means that species associated with few other species (specialists) preferentially interact with species that interact with many others (generalists), causing asymmetric specialization between interacting partners (Bascompte et al. 2003; Thompson 2005; Burgos et al. 2009). However, generalist species tend to interact with one another, forming a dense core of symmetric interactions (Bascompte et al. 2006; Vázquez et al. 2007; Guimarães et al. 2011). The nested pattern has already been observed in different mutualistic networks, such as: plant-pollinator, plant-ant, plant-disperser, clownfish-anemone and marine fish cleaning symbioses (Bascompte et al. 2003; Guimarães et al.

2006; Lewinsohn et al. 2006; Guimarães et al. 2007; Ollerton et al. 2007). Several factors have been proposed to explain the origin of these patterns in mutualistic networks (for more information, please see: Medan et al. 2007; Morales & Vázquez 2008; Nielsen & Bascompte 2007; Rezende et al. 2007; Stang et al. 2009; Vázquez et al. 2009). However, symbiotic networks are not nested, because they present great compartmentalization of interactions among the species involved (Guimarães et al. 2007).

Mutualistic interactions networks can be viewed in different ways, however, there are two main ways to visualize these interactions through: bipartite graphs (Figure 1A) or/and ordered matrices (Figure 1B). Using as example a network of interactions between plant-animals, in bipartite graphs, the left nodes represent different animal species, and the right nodes correspond plant species that interact positively with the animals (Figure 1A). Lines, also called "links", connect positively interacting species; On the other hand, the visualization of interactions in ordered matrices indicates that each column represents one plant species and each row represents a animal species, and filled cells represent positive interactions among species. (Figure 1B).




Figure 1. Main ways to view the structure of an interaction networks nested and asymmetric: A) bipartite graphs and B) ordered matrices. Both forms of representation are ordered according to interactions number of the species, where species that have fewer interactions are subsets of species with more interactions. Cells painted of gray in Figure

1B represent the core of generalists species (Figures modified from Lewinsohn et al. 2006).

According to Thompson (2005) and Stanton (2003), studies of interactions only among pairs of species are insufficient to understand the evolutionary and coevolutionary processes in mutualistic interactions. Several authors have studied the implications of coevolution in mutualistic networks of interacting species (Jordano et al. 2003; Thompson 2005, Bascompte et al. 2006; Guimarães et al. 2007; Jordano et al. 2010; Guimarães et al. 2011). The focus of these studies is mainly in the core of generalist species, due the symmetric strength of the interacting species (Jordano et al. 2003; Bascompte et al. 2006; Guimarães et al. 2011). The generalist core can drive the evolution of the whole community because the species of the core interact with virtually all species of the matrix (Jordano et al. 2003; Bascompte et al. 2006; Guimarães et al. 2007; Jordano et al. 2010; Guimarães et al. 2011). Moreover, the generalist core can act as a coevolutionary vortex of more specialized interactions, where the convergence of traits on both sides (e.g. animals and plants) of the interaction increases over evolutionary time (Thompson 2005; Guimarães et al. 2007). Finally, understand the feature and factors that influence the interactions of the species generalist core is essential to understand the current view of coevolutionary process in mutualistic interactions (Bascompte et al. 2003; Thompson, 2005; Bascompte et al. 2006; Jordano et al. 2010).

My aim of this dissertation was, to 1) evaluate the topology of mutualistic (ants and plants with extrafloral nectaries) and neutral (ants and plants without extrafloral nectaries) networks in ant-plant interactions; 2) evaluate the turnover of species composition among these networks; 3) determine through a simples mathematical model the role of relative abundance of species in the probability of pairwise interactions in ant-plant mutualistic networks; 4) evaluate how the topological properties and the spatial turnover of species composition of ant-plant mutualistic vary along an spatial scale. My dissertation was divided into two interrelated chapters. In Chapter 1, I show that the topology of mutualistic and neutral networks are different, and which the core of generalist species interact more among themselves than expected by their abundance in mutualistic networks. Moreover, is little the turnover of ants composition among the mutualistic and neutral networks. In Chapter 2, I show that in mutualisms between ants and plants with EFNs the networks topology remains unaltered at the spatial scale studied (5.099 m). In addition, even that ants and plants composition of networks changes over space, the species of core generalist

remains the same along a regional scale on a *terra-firme* forest in Brazilian Meridional Amazon. Finally, the list of ants and plants species collected in this study addition of the instructions for authors of the journals that I will submit the manuscripts are included in four appendices at the end of this dissertation.

References

- Bascompte, J.; Jordano, P.; Melián, C.J.; Olesen, J.M. (2003). The nested assembly of plant-animal mutualistic networks. *Proceedings of the National Academy of Sciences USA* 100: 9383-87.
- Bascompte, J.; Jordano, P.; Olesen, J.M. (2006). Asymmetric coevolutionary networks facilitate biodiversity maintenance. *Science* 312: 431-433.
- Bronstein, J.L. (2001) Mutualisms. In: Fox, C.; Fairbairn, D.; Roff, D. (Eds.). *Evolutionary ecology: perspectives and synthesis*. Oxford University Press: New York. pp. 315-330.
- Bronstein, J.L.; Alarcon, R.; Geber, M. (2006). Tansley Review: Evolution of insect/plant mutualisms. *New Phytologist* 172(3): 412-428.
- Burgos, E.; Ceva, H.; Hernández, L.; Perazzo, R.P.J. (2009). Understanding and characterizing nestedness in mutualistic bipartite networks. *Computer Physics Communications* 180: 532-535.
- Burslem, D.F.R.P.; Pinard, M.A.; Hartley, S.E. (2005). *Biotic interactions in the tropics. Their role in the maintenance of species diversity*. Cambridge University Press: Cambridge. 580p.
- Del-Claro, K. & Torezan-Silingardi, H.M. (2009). Insect-plant interactions: new pathways to a better comprehension of ecological communities in Neotropical savannas. *Neotropical Entomology* 38(2): 159-164.
- Dyer, L.A.; Walla, T.R.; Greeney, H.F.; Stireman, J.O.; Hazen, R.F. (2010). Diversity of Interactions: A Metric for Studies of Biodiversity. *Biotropica* 42(3): 281-289.
- Ferrière, R.; Gauduchon, M.; Bronstein, J.L. (2007). Evolution and persistence of obligate mutualists and exploiters: competition for partners and evolutionary immunization. *Ecology Letters* 10: 115-126.
- Guimarães, P.R.; Rico-Gray, V.; Reis, S.F.; Thompson, J.N. (2006). Asymmetries in specialization in ant-plant mutualistic networks. *Proceedings of the Royal Society, B, Biological Sciences* 273: 2041-2047

- Guimarães, P.R.; Sazima, C.; Reis, S.F.; Sazima, I. (2007). The nested structure of marine cleaning symbiosis: is it like flowers and bees? *Biology Letters* 3:51-54.
- Guimarães, P.R.; Jordano, P.; Thompson, J.N. (2011). Evolution and coevolution in mutualistic networks. *Ecology Letters* 14: 877-88.
- Hoeksema, J.D. & Bruna, E.M. (2000). Pursuing the big questions about interspecific mutualism: a review of theoretical approaches. *Oecologia* 125:321-330.
- Janzen, D.H. (1974). The deflowering of Central America. *Natural History* 83(4): 49-53.
- Janzen, D.H. (1985). The natural history of mutualisms. In: Boucher, D.H. (Ed.). *The biology of mutualism*. Oxford University Press: New York. pp. 40-99.
- Jordano, P.; Bascompte, J.; Olesen, J.M. (2003). Invariant properties in coevolutionary networks of plant-animal interactions. *Ecology Letters* 6: 69-81.
- Jordano, P. (2010). Coevolution in multispecific interactions among free-living species. *Evolution: Education and Outreach* 3(1): 40-46.
- Lewinsohn, T.M.; Prado, P.I.; Jordano, P.; Bascompte, J.; Olesen, J.M. (2006). Structure in plant-animal interaction assemblages. *Oikos* 113:174-84.
- Medan, D.; Perazzo, R.P.J.; Devoto, M.; Burgos, E.; Zimmermann, M.G.; Ceva, H.; Delbue, A.M. (2007). Analysis and assembling of network structure in mutualistic systems. *Journal of Theoretical Biology* 246: 510-521.
- Montoya, J.M.; Pimm, S.L.; Sole, R.V. (2006). Ecological networks and their fragility. *Nature* 442: 259-264.
- Morales, J.M. & Vázquez, D.P. (2008). The effect of space in plant-animal mutualistic networks: insights from a simulation study. *Oikos* 117: 1362-1370.
- Nielsen, A. & Bascompte, J. (2007). Ecological networks, nestedness and sampling effort. *Journal of Ecology* 95: 1134-1141.
- Ollerton, J.; McCollin, D.; Fautin, D.G.; Allen, G.R. (2007) Finding NEMO: nestedness engendered by mutualistic organization in anemone-fish and their hosts. *Proceedings of the Royal Society, B, Biological Sciences* 274: 591-598.
- Rezende, E.L.; Jordano, P.; Bascompte, J. (2007). Effects of phenotypic complementarity and phylogeny on the nested structure of mutualistic networks. *Oikos* 116: 1919-1929.
- Stang, M.; Klinkhamer, P.G.L.; Waser, N.M.; Stang, I.; Van-der-Meijden, E. (2009). Size-specific interaction patterns and size matching in a plant-pollinator interaction web. *Annals of Botany* 103: 1459-1469.

- Stanton, M.L. (2003). Interacting guilds: moving beyond the pairwise perspective on mutualisms. *American Naturalist* 162: S10-S23.
- Thompson, J.N. (2005). *The geographic mosaic of coevolution*. University of Chicago Press: Chicago. 443p.
- Toby-Kiers, E.; Palmer, T.M.; Ives, A.R.; Bruno, J.F.; Bronstein, J.L. (2010). Mutualisms in a changing world: an evolutionary perspective. *Ecology Letters* 13(12): 1459-1474.
- Vázquez, D.P.; Melián, C.J.; Williams, N.M.; Blüthgen, N.; Krasnov, B.R.; Poulin, R. (2007). Species abundance and asymmetric interaction strength in ecological networks. *Oikos* 116: 1120-1127.
- Vázquez, D.P.; Blüthgen, N.; Cagnolo, L.; Chacoff, N.P. (2009). Uniting pattern and process in plant-animal mutualistic networks: a review. *Annals of Botany* 103: 1445-1457.

Chapter 2

The structure of ant-plant mutualistic networks: is abundance enough?*

** This manuscript will be submitted to Ecology*

Submitted to Ecology

Running head: *The effect of abundance in ant-plant networks*

The structure of ant-plant ecological networks: is abundance enough?

Wesley Dáttilo ^{1*}; Thiago J. Izzo ¹; Flávia M.D. Marquitti ² and Paulo R.

Guimarães Jr ²

¹ Departament of Ecology and Botany, Community Ecology Lab.,
Universidade Federal de Mato Grosso, Cuiabá, MT, CEP 78060-900, Brazil.

² Departament of Ecology, Instituto de Biociências, Universidade de São
Paulo, São Paulo, SP, CEP 05508-900, Brazil.

** Corresponding author:*

Wesley Dáttilo

UFMT-IB-PPGECB

Avenida Fernando Correa/ SN

Cuiabá, Mato Grosso State, 78068-900

Brazil

E-mail wdattilo@hotmail.com

ABSTRACT.

The structural organization of mutualism networks provides insights into processes shaping biodiversity. To know the mechanisms that shape this organization is essential for us to understand the ecological and evolutionary dynamics of the interacting species. Recent studies have suggested that species abundance is one of the most fundamental criteria shaping mutualistic networks. However, the role of species abundance on mutualistic networks is still unclear. In this work we studied plants with extrafloral nectaries and associated ants to show that the natural abundance of ants on vegetation explained just a part of the frequency of mutualistic interactions and that it is independent of ant species compositions. In ant-plant mutualistic networks, the nestedness was higher than predicted by the abundance. In addition, the generalist core of these networks interacts more among themselves than expected by their abundances. This generalist core was formed by competitively superior ants that have behavioral and ecophysiological traits to use liquid food and are that exclude other species from the same resource. Such generalist core conformation being stable over space and time could have serious implications on coevolutionary patterns of the system.

KEYWORDS: coevolution, nestedness, neutrality, plant-animal interactions.

1. INTRODUCTION

A central goal of ecology is to understand the mechanisms that determine the structure of ecological communities at different spatial and temporal scales (Turner 1990, Levin 1992, Williams and Martinez 2008, Zhou and Zhang 2008). Recent studies have focused on the network structure of mutualisms and found non-random patterns of interaction on a wide range of ecosystems (Bascompte et al. 2003, Lewinsohn et al. 2006, Guimarães et al. 2007, Vázquez 2009a). These non-random patterns influence the whole ecological and evolutionary dynamics of the interacting species (Jordano et al. 2003, Thompson 2005, Bascompte et al. 2006, Montoya et al. 2006, Guimarães et al. 2011).

Relative species abundance is one of the most fundamental criteria shaping the ecological networks, and it seems to be an important factor in the probability of interactions (Vázquez et al. 2007, 2009b, Verdú and Valiente-Banuet 2011). In this case, abundant species should interact most frequently with each other and with other less abundant species, but less abundant species will rarely interact with them (Krishna et al. 2008, Vázquez et al. 2007, 2009b). However, the role of species abundance on mutualistic networks is still unclear.

A strictly system to study questions on abundance and dominance on interaction networks is the ant-plant system. Ant-plant interactions are commonly found in tropical rainforests, in which more than 94% of arthropods and 86% of the biomass collected in canopies are ants (Majer 1990, Tobin 1995). The observed high frequency of ant foraging on the surface of plants is due to the high availability of different food and nesting sites within their structures (Andersen 1990, Blüthgen et al. 2000, Davidson et al. 2003). The interactions between ants and plants with extrafloral nectaries (EFNs) are key-ecological interactions in tropical rainforests and well documented in literature. In these associations, plants produce nutritious liquid to attract ants (Baker et al. 1978, Koptur et al. 1998, Rico-Gray and Oliveira 2007). In exchange for food, the ants defend plants against potential

herbivores (Del-Claro et al. 1996, Oliveira et al. 1999, Rico-Gray and Oliveira 2007). However, despite nectar being a key resource for ants foraging on plants, there is an adaptive filter in which only a few ant species have adaptations for the acquisition, storage and digestion of liquid resources (Fowler et al. 1991, Oliveira and Brandão 1991, Davidson et al. 2003, 2004).

With the use of the network theory in ecological interaction studies in recent years, some authors have found the existence of a dense core of symmetric interactions in ant-plant mutualistic networks, where generalist species tend to interact with one another (Guimarães et al. 2006, Chamberlain et al. 2010, Sugiura 2010). Moreover, these studies show that species associated with few other species (specialists) preferentially interact with species that interact with many others (generalists), forming an asymmetrical and nested pattern of interaction (Guimarães et al. 2006, Chamberlain et al. 2010, Sugiura 2010). Several factors have been proposed to explain the origin of these non-random patterns in mutualistic networks, such as spatial distribution of individuals and species (Morales and Vázquez 2008, Burkle and Alarcón 2011), species richness (Medan et al. 2007), foraging behavior (Vázquez et al. 2009a), sampling effects (Nielsen and Bascompte 2007), phylogeny (Rezende et al. 2007), and phenotypic traits of interacting individuals (Stang et al. 2007). However, species abundance seems to be the best predictor in ant-plant mutualistic networks (Vázquez et al. 2007, Chamberlain et al. 2010). Thus, abundant species tend to find individuals of other abundant species more often than individuals of rare species (Vázquez et al. 2009b). As EFNs are a food resource, their discovery and dominance by ants being dependent only of abundance, imply no actual competition. However, this is not supported by literature once the nectar secreted is a predictable resource and rich in energy, and different ant species compete for the same resource (Hölldobler and Wilson 1990, Dreisig 2000, Blüthgen and Fiedler 2004ab).

On the other hand, among the arboreal ant communities, some species do not have adaptations for feeding liquid resources, but forage on vegetation for prey, for example (Andersen 1990). In plants without the presence of EFNs and honeydew secreted by homopterous insects, resources cannot be predicted, and ants randomly forage using the plant only as substrate (Blüthgen et al. 2000). The presence of such ants in a plant may be driven by neutral, not-deterministic factors, such as the available foraging area. Thus, we expect the existence of two subsets with different composition of ant species foraging on plants with and without EFNs. These subsets should be maintained possibly by the adaptive physiological characteristics of ants, as well as competition where some ants monopolize EFNs plants.

In this study, we hypothesized that there is a particular ant species assembly foraging on plants with EFNs determined by behavior and ecophysiological factors (adaptations for liquid diets and monopolizing the resource) that differ from those ants randomly foraging in the foliage, generating two different subsets. This assembly of nectar-feeding ants monopolizes the resource and interacts with plants more among themselves than expected by their abundance. Moreover, we expected that, due to the liquid-energy food source of EFNs, ants would not interact randomly with plants. Therefore, the topological structure of networks generated through deterministic (plants with EFNs) and non-deterministic (plants without EFNs) processes in ant-plant interactions could be different. In order to test our hypothesis, we evaluated the topology of the ecological networks between ants and plants with and without EFNs, and the turnover of ant species composition among these networks through the additive partitioning of diversity. We also determined through a simple mathematical model the role of the relative species abundance in the pairwise probability interactions in networks of ants and plants with EFNs in a tropical rainforest inserted in the Brazilian Meridional Amazon.

2. METHODS

2.1) Study area

We conducted this study at the São Nicolau Farm (9°48'S and 58°15'W, elev. 254m), located in the municipality of Cotriguaçu, north of Mato Grosso, Brazil. According to the Köppen classification, the climate is tropical wet (Am) with annual average of: 24 °C temperature, 85% humidity, and 2.300 mm precipitation (Camargo et al. 2010, Dáttilo et al. 2012). The study region has two distinct seasons, a rainy season between November and April and a dry season between May and October. The area is characterized as a *terra-firme* dense rainforest inserted in the Brazilian Meridional Amazon (Veloso et al. 1991, Camargo et al. 2010). The reserve area covers 7.000 ha of continuous forest, surrounded by a much larger area of continuous forest. The terrain is undulating with altitudinal variation of 50 m between the plateaus and the river side terrains. Canopy trees range from 30-40 m high with some emergent trees reaching 50 m. The understory is relatively open, with high frequency of *Orbignya phalerata* Mart. (Arecaceae).

2.2) Data Collection

We collected data in a module managed by the Brazilian Research Program in Biodiversity (PPBio) inserted at São Nicolau Farm. The module consists of two 5 km parallel trails East-West one kilometer apart. In both trails, one sampling point with 250 m x 25 m (6.250 m²) was made every km., totaling 12 sampling points in the entire module. The central trail of each plot was established minimizing variations of soil and altitude, and increasing the precision of estimates for predictor variables (Magnusson et al. 2005) (additional details about module and trails please see <http://ppbio.inpa.gov.br>). Here, we considered each of the 12 sampling points as independent samples of ants and plants, generating 12 different ant-plant interaction networks. We assume that ants and plants are

sessile organisms (Fourcassié et al. 2003) and the distance between sampling points is enough so as to guarantee that organisms of a given plot can never interact with the organisms of another plot.

We collected ants and plants in December 2010 and January 2011 always between 9:00 h and 15:00 h. In each of the 12 sampling points, we collected ants foraging in all plants with EFNs that were accessible to the collector (from 0.5 m to 3 m). For ant collection, we used a method similar to the entomological umbrella in which the branches were shaken and all the ants that fell were collected in a white squared cloth board of 1.2 m² previously placed under the branch (Bestelmeyer et al. 2000). This method is very effective as some species, particularly of the genera *Camponotus* and *Ectatomma* drop from the plant at the slightest sign of disturbance made by the collector (W. Dáttilo, pers. obs.). We observed and collected additional ants, especially more secretive species before and after performing this procedure in order to register ants feeding on EFNs. For each plant with EFNs where ants were collected, we selected a plant without EFNs with similar structure (height, width and number of branches) nearby. No plants with homopterous and other visible liquid-resource sources were considered when sampling plants without EFNs. All selected plants should be at least 10 m apart to minimize the possibility of collecting the same ant colony foraging on different plants. We used the entomological umbrella methodology instead of feeding baits to minimize biases towards ant species with efficient recruitment behavior and ants with particular feeding habits.

Plants and ants were identified to the lowest possible taxonomic level using taxonomic keys and with the help of specialists. Ant specimens were deposited in the Entomological Section of the Zoological Collection of Universidade Federal de Mato Grosso, Brazil (CEMT), and the plants were deposited in the Herbário Centro-Norte Mato-Grossense (CNMT).

2.3) *Network topology*

Mutualistic interactions networks can be viewed in two main ways: bipartite graphs or/and ordered matrices. In bipartite graphs, nodes represent different animal species and links are positive interactions among species. On the other hand, the visualization of interactions in ordered matrices indicates that each column represents one animal species and each row represents a plant species, and filled cells represent positive interactions among species. Thus, in order to describe the network topology of the ecological networks of the ants and plants with and without EFNs, we calculated the difference in the number of species (ants and plants) and the interaction frequency of the ecological networks between ants and plants with and without EFNs. In addition, we also calculated the following metrics: connectance, mean and variance of number of links per plant and ant species, modularity and nestedness in each of the 12 interaction networks between ants and plants with and without EFNs. The connectance (C) is the proportion of possible links that are actually made (Jordano 1987). Mean and variance of number of links per plant and ant species were obtained from the arithmetic mean of the number of interactions in which each species was involved.

We calculated modularity using the modularity index M (range 0 - 1), which estimates the degree in which groups of species (ants and plants) interact more among each other than with species in other groups in the network (Guimerà and Amaral 2005). The M index decreases when the fraction of between-module links increases in the total network. High values of M indicate that the ants and plants form modules that are semi-independent of other interactions within the network (Olesen et al. 2007). We tested the significance of index M for each network through 1.000 simulated networks generated by a null model, in order to assess whether the value of M observed in the empirical network is higher than expected for networks of equal size and with similar heterogeneity in interactions among species (Null Model II, Bascompte et al. 2003). We made the null model network through

a routine in MATLAB, and the M indices of all networks were calculated by the software Netcarto (Guimerà and Amaral 2005).

We used the NODF index (Nestedness metric based on Overlap and Decreasing Fill) to estimate the nestedness value of networks, using ANINHADO software (Guimarães and Guimarães 2006). This metric is a much better nestedness metric than others and less prone to type-I statistical error, since it is based on the nestedness of all pairs of columns and rows in the matrix (Almeida-Neto et al. 2008). To assess if the nestedness value observed was higher than expected by random patterns of interaction, we tested the nestedness degree of each network with 1,000 networks generated by Null Model CE (Null Model II). In addition, we also calculated the nestedness value standardizing the difference of richness, connectance and heterogeneity of interactions among the networks using the Z-Score metric, which is defined as: $Z_{\text{nodf}} = (x - \mu) / \sigma$, where x = NODF value observed, μ = NODF mean value of randomized matrices, and σ = is the standard deviation of the randomized matrices (Almeida-Neto et al. 2008, Ulrich et al. 2009). To test the difference of all the metrics described above for the ecological networks of ants and plants with and without EFNs, we used the paired T-Test (paired per plot) using R-Project software (R Development Core Team 2010).

2.4) Overrepresentation of species interactions

To determine if the matrices of interaction networks of ants and plants with EFNs have their structure determined by abundance, we developed a simple mathematical model that allowed us to differentiate in which regions of the observed matrices the ant-plant interactions occurred more than expected by the abundance. In this model we used the ant records of neutral interactions as a measure of ant abundance in the foliage. As plants without EFNs offer no predictable resource to the ants, the frequency of ants in these plants

reflects the spatial abundance of these species in the vegetation without the aggregation caused by the resource (Blüthgen et al. 2000).

Initially, we built an interaction matrix between ants and plants, where a_{ij} = number of interactions between plant species i and ant species j . The theoretical matrix of abundance was determined by $b_{ij} = \left(\frac{F_i}{F_p} \cdot \frac{F_j}{F_a} \right)$, where F_i is the absolute frequency of a given plant with EFNs in the plot, F_p is the total frequency of plants with EFNs found in the plot, F_j is the absolute frequency of a given ant collected in plants without EFNs in the plot, F_a is the total frequency of ants collected in plants without EFNs found in the plot. The probability of any particular interaction between ants and plants occurs was

determined as $P_{(aij)} = \frac{a_{ij}}{\sum_n^p \sum_m^a a_{nm}}$, where a_{ij} is the number of interaction events of the

matrix and $\sum_n^p \sum_m^a a_{nm}$ is the total number of interaction events between ants and plants with EFNs. The probability of an interaction occurring is determined by the abundance of

interacting partners as $P_{(bij)} = \frac{b_{ij}}{\sum_o^p \sum_g^a a_{og}}$, where b_{ij} is the number of interaction events of

the matrix and $\sum_o^p \sum_g^a a_{og}$ is the total number of interaction events in plants without EFNs.

Finally, we computed the differences between the actual probability of a given interaction occurring and the probability derived from the species abundances, $C_{ij} = (A' - B)$. We

standardized C_{ij} to $C_{ij}' = \frac{C_{ij}}{|\max(C_{ij})|}$, where $|\max(C_{ij})|$ was the maximum value of C_{ij} of

matrix to allow across-plot comparisons. $C_{ij} < 0$ are cases in which plant i and ant j interact less than expected by their abundances; $C_{ij} > 0$ are cases in which plant i and ant j interact

more than expected from their abundances; and values equal to 0 representing the cases where ant species collected in plants without EFNs was not collected in plants with EFNs.

2.5) Additive partitioning of diversity

To evaluate the turnover of ant species composition (β -diversity) between the networks of ants and plants with and without EFNs, we calculated the additive partitioning of diversity as proposed by Veech et al. (2002). From the total ant richness found in each plot (γ -diversity), we calculated the α -diversity, defined as: $\alpha_{\text{mean}} = (\alpha_1 + \alpha_2) / 2$; where α_1 = ant richness of plant networks with EFNs found in plot; α_2 = ant richness of plant networks without EFNs found in plot. Then, we calculated the β -diversity, defined as: $\beta = \gamma - \alpha_{\text{mean}}$. In addition, to summarize the composition of the ant community in plants with and without EFN, we ordered the similarity between points using Non-Metric Multidimensional Scaling (NMDS) and tested the difference in the ant species composition through a permutation test (10.000 permutations) based on an analysis of similarities (ANOSIM) (Supplementary information). Additionally, to exclude the effect of the presence of EFNs and evaluate if the presence of EFNs was a factor which caused an increase in ant species richness, we randomized the plants classification (presence or absence of EFNs) while keeping fixed the plant richness from sampling points (γ -diversity). We performed this randomization 1.000 times. In the end of each set of randomization, we counted the ant network richness of plants with EFNs (α_1) and plants without EFNs (α_2), in order to assess whether the ant network richness of plants with EFNs is higher than expected by ant network richness of plants without EFNs. The average of these values (α_{mean}) was subtracted from the γ -diversity to obtain the β -diversity. We did this randomization using MATLAB.

3. RESULTS

In this study, we recorded 238 plant species (72 with EFNs) and 149 ant species. The number of plant species with EFNs was lower (Mean \pm SD: 21.4 ± 3.77) than plants without EFNs (27.2 ± 3.97 , $t = -3.093$, $df = 11$, $P = 0.011$). However, the number of ant species on plants with EFNs (23.2 ± 5.85) was not different from plants without EFNs in the sampling points studied (23.3 ± 4.11) ($t = -0.0647$, $df = 11$, $P = 0.949$). Likewise, the interaction frequency was also equal between networks of ants and plants with EFNs (77.91 ± 12.58) and plants without EFNs (75.58 ± 10.46) ($t = 1.239$, $df = 11$, $P = 0.241$). The mean and variance as to the number of links per plant with EFNs (Mean \pm Variance: 3.46 ± 0.28) were higher than plants without EFNs (Mean \pm Variance: 2.69 ± 0.48) ($t = -3.876$, $df = 11$, $P = 0.003$, Figure 1A). For ants, the mean and variance of the number of links per ant did not differ between networks of plants with EFNs (Mean \pm Variance: 2.67 ± 0.14) and without EFNs (Mean \pm Variance: 2.64 ± 0.10) ($t = 0.258$, $df = 11$, $P = 0.801$) (Figure 1B).

The network connectance of ants and plants with EFNs were higher (Mean \pm SD: 0.140 ± 0.03) than the network connectance of ants and plants without EFNs (0.109 ± 0.02) ($t = -3.528$, $df = 11$, $P = 0.005$) (Figure 1C). In none of our sampling points did we observe significantly higher modularity than expected by the heterogeneity of interactions ($P > 0.05$). However, there is a tendency that the modularity index was lower in networks of ants and plants with EFNs (0.459 ± 0.059) than in the networks of ants and plants without EFNs (0.519 ± 0.061) ($t = 3.552$; $df = 11$; $P = 0.005$) (Figure 1D).

The nestedness was higher in networks of ants and plants with EFNs (Mean \pm SD: 21.01 ± 4.46) than the nestedness in networks of ants and plants without EFNs (15.75 ± 3.33) ($t = -3.427$; $df = 11$; $P = 0.006$) (Figure 1E), as well as their standard normal deviate (Z-Score metric): networks of ants and plants with EFNs (3.63 ± 1.51) and networks of ants and plants without EFNs (2.34 ± 1.35) ($t = -2.270$; $df = 11$; $P = 0.04$) (Figure 1F).

The model developed in this study showed that the species of ants and plants that form the generalist core region interact more among themselves than expected by their abundances (Figure 2). Moreover, the species on the periphery of the network interact less than expected by their abundances. This same pattern was repeated in the 12 networks evaluated in this study.

The β -diversity turnover of ant and plant networks with and without EFNs involves approximately one third of the total diversity, indicating limited turnover of ant composition among the networks. In addition, when we randomized the plant classification (presence or absence of EFNs), there was no difference between the observed and expected values for β -diversity. In other words, the presence of EFNs did not propitiate an increase in the number of ant species ($t= 1.448$, $df= 11$, $P= 0.176$) and the species turnover does not change in intensity in the networks. Thus, the ant species present in plants with EFNs were similar to those in plants without EFNs (Figure 3). Additional NMDS ordination of the ant community followed by ANOSIM showed similar results (see supplementary information).

4. DISCUSSION

Recently, some studies have shown that the relative species abundance is an important factor structuring mutualistic networks (Krishna et al. 2008, Vázquez et al. 2007, 2009b, Verdú and Valiente-Banuet 2011). Such studies show that the asymmetric interaction among species results from their abundances and that species abundance explains almost a third of the nested pattern in mutualistic networks. Here we showed that the topological structure of networks generated through deterministic process (plants with EFNs) is different than that of the networks generated by the natural abundance of ants on vegetation. Moreover, the core of generalist species in the ant and plant networks with EFNs interact more among themselves than expected by their abundance.

Some studies have suggested that, within a biological community, the difference in species abundances and sampling techniques can generate nested patterns (Fischer and Lindenmayer 2002, Lewinsohn et al. 2006, Blüthgen 2010), including simulated neutral networks (Blüthgen et al. 2008). This possibly explains the nested pattern on our ant and plant networks without EFN, as less abundant species tend to be subsets of the more abundant species (Vázquez et al. 2009b). When comparing the ecological networks in ant-plant interactions, however, we showed that the ant and plant networks with EFNs are more connected and nested than the ant and plant networks without EFNs, possibly due to difference in networks size. However, in this study nestedness cannot be explained by different patterns of richness, connectance and heterogeneity of interactions, since we controlled these variables through the Z-Score metric. Also the nestedness found is not explained by richness and mean number and nor is its variance of links per ant species, since they are similar in both ecological networks and controlled by the Z-Score metric. Thus, this points out that the structure of mutualistic ant-plant interactions found in this study is fundamentally different from what we should expect from connectances, species richness and abundances alone.

According to Bastolla et al. (2009) the nested pattern in mutualistic networks between plants and their animal pollinators or seed dispersers reduces interspecific competition enhancing the number of coexisting species. This is due to the small number of shared partners when compared with fully connected and compartmentalized networks (Bastolla et al. 2009). In the interaction ant and plant networks with EFNs studied here, we showed that within the matrices there is a region, or a "hard-core", where generalist plant and ant species interact more among themselves than expected by their abundances. In this study, similar to other mutualism networks, we assumed that all resources offered are nutritionally equal; however, we know that the quantity and quality of nectar may vary among species and plant individuals (Schupp and Feener 1991, Heil 2000). Maybe the core

of generalist plant species has better resources and only ant species competitively superior (i.e. more recruitment or aggressive) monopolize the resources (Blüthgen and Fiedler 2004ab, Heil and McKey 2003) independent of its spatial abundance. Thus, we hypothesize that the hard-core region might be formed by competitively superior ant species that can exclude others. This hard-core region can be generated by ants that have ecophysiological and behavioral traits that enable them to find, dominate, and use the resource efficiently and over a longer time period.

In interactions between ants and plants with EFNs, only a few ant species of the subfamilies Myrmicinae, Formicinae and Dolichoderinae are known to have physiological adaptations for the acquisition, storage and digestion of liquid resources secreted by EFNs (Fowler et al. 1991, Oliveira and Brandão 1991, Davidson et al. 2003, 2004). Although there is an adaptive filter for the ants that feed on the liquid resources secreted by EFNs, we found a low turnover of ant species foraging on plants with and without EFNs. The absence of a particular ant species composition foraging on plants with EFNs was also observed by Schoereder et al. (2010). Therefore, the nested pattern observed in both ecological networks also studied here can not be explained by the ant species composition, as the presence of EFNs does not influence a particular ant species composition associated with this resource. Some ant genera that do not have adaptations for feeding on liquids resource can carry liquids externally, and these genera strongly associate with EFN bearing plants (e.g. *Acanthoponera*, *Ectatomma*, *Heteroponera*, *Pachycondyla*, *Paraponera*, *Pseudomyrmex*) (Oliveira and Brandao 1991, Almeida and Figueredo 2003, Davidson et al. 2003, 2004). But, in this study, the ant species composing the "hard-core" compartment show adaptations for liquid diet, such as: *Azteca*, *Brachymyrmex*, *Camponotus*, *Crematogaster*, *Dolichoderus*.

Here we showed that abundance is important in the nested pattern (Vázquez and Aizen 2006, Blüthgen et al. 2008, Vázquez et al. 2009b, Verdú and Valiente-Banuet 2011),

but the core of generalist species interact more than expected by their abundance. Such core conformation implies that the generalist core can act as a coevolutionary vortex of more specialized interactions, where the convergence of traits occurs on both sides of the interaction, increasing over evolutionary time (Bascompte et al. 2003, Thompson 2005, Guimarães et al. 2007). Furthermore, it is possible that the hard-core is less variable over space and time, and can affect the ecological and evolutionary dynamics of these interactions. For instance, a stable core will have unknown implications to the geographic mosaic of coevolution (Thompson 2005). Finally, we suggest studies evaluating the consequences of spatial and temporal variations in ant-plant and other types of mutualistic networks as being the next step in the analysis of stability in core interactions.

ACKNOWLEDGEMENTS

We would like to thank Jéssica Falcão for your help during the fieldwork and for comments on earlier versions of this manuscript. We also thank Office National des Forêts Brazil for logistical support and Brazilian PPBio project for financial support. WD was granted with a master fellowship from the Coordination for the Improvement Higher Education Personnel (CAPES-Brazil). PRG and FMDM acknowledge the financial support of FAPESP. This is publication XXX in the NEBAM technical series.

LITERATURE CITED

- Almeida-Neto, M., P. R. Guimarães Jr, P. Guimarães, R. D. Loyola, and W. Ulrich. 2008. A consistent metric for nestedness analysis in ecological systems: reconciling concept and measurement. *Oikos* 117:1227–1239.

- Almeida, A. M., and F. A. Figueiredo. 2003. Ants visit nectaries of *Epidendrum denticulatum* (Orchidaceae) in a Brazilian rainforest: effects on herbivory and pollination. *Brazilian Journal of Biology* 63:551–558.
- Andersen, A. N. 1990. The use of ant communities to evaluate change in Australian terrestrial ecosystems: a review and a recipe. *Proceedings of the Entomological Society of Australia* 16:347–357.
- Baker, H. G., P. A. Opler, and I. Baker. 1978. A comparison of the amino acid complements of floral and extrafloral nectars. *Botanical Gazette* 139:322–332.
- Bascompte, J., P. Jordano, C. J. Melián, and J. M. Olesen. 2003. The nested assembly of plant–animal mutualistic networks. *Proceedings of the National Academy of Sciences USA* 100:9383–87.
- Bascompte, J., P. Jordano, and J. M. Olesen. 2006. Asymmetric coevolutionary networks facilitate biodiversity maintenance. *Science* 312:431–433.
- Bastolla, U., M. A. Fortuna, A. Pascual–García, A. Ferrera, B. Luque, and J. Bascompte. 2009. The architecture of mutualistic networks minimizes competition and increases biodiversity. *Nature* 458:1018–1021.
- Bestelmeyer, B. T., D. Agosti, L. E. Alonso, C. R. F. Brandão, W. L. Brown, J. H. C. Delabie, and R. Silvestre. 2000. Field techniques for the study of ground–dwelling ants: An overview, description and evaluation. Pages 122–144 in D. Agosti, J. D. Majer, L. E. Alonso, and T. R. Schultz, editors. *Ants: Standard methods for measuring and monitoring biodiversity*. Smithsonian Institution Press, Washington, USA.
- Blüthgen, N., M. Verhaagh, W. Goitía, K. Jaffé, W. Morawetz, and W. Barthlott. 2000. How plants shape the ant community in the Amazonian rainforest canopy: the key role of extrafloral nectaries and homopteran honeydew. *Oecologia* 125:229–240.

- Blüthgen, N., and K. Fiedler. 2004a. Competition for composition: lessons from nectar-feeding ant communities. *Ecology* 85:1479–1485.
- Blüthgen, N., and K. Fiedler. 2004b. Preferences for sugars and amino acids and their conditionality in a diverse nectar-feeding ant community. *Journal of Animal Ecology* 73:155–166.
- Blüthgen, N., G. Gottsberger, and K. Fiedler. 2004. Sugar and amino acid composition of ant-attended nectar and honeydew sources from an Australian rainforest. *Austral Ecology* 29:418–429.
- Blüthgen, N., F. Menzel, T. Hovestadt, B. Fiala, and N. Blüthgen. 2007. Specialization, constraints, and conflicting interests in mutualistic networks. *Current Biology* 17:341–346.
- Blüthgen, N., J. Fründ, D. P. Vázquez, and F. Menzel. 2008. What do interaction network metrics tell us about specialization and biological traits? *Ecology* 89:3387–3399.
- Blüthgen, N. 2010. Why network analysis is often disconnected from community ecology: A critique and an ecologist's guide. *Basic and Applied Ecology* 11:185–195.
- Burgos, E., H. Ceva, L. Hernández, and R. P. J. Perazzo, R.P.J. 2009. Understanding and characterizing nestedness in mutualistic bipartite networks. *Computer Physics Communications* 180:532–535.
- Burkle, L. A., and R. Alarcón. 2011. The future of plant-pollinator diversity: understanding interaction networks across time, space, and global change. *American Journal of Botany* 98:1–11.
- Camargo, F. F., R. B. Costa, M. D. V. Resende, R. A. R. Roa, N. B. Rodrigues, L. V. Santos, A. C. A. Freitas. 2010. Variabilidade genética para caracteres morfométricos de matrizes de castanha-do-brasil da Amazônia Mato-grossense. *Acta Amazonica* 40:705–710.

- Chamberlain, S. A.; J. R. Kilpatrick, and J. N. Holland. 2010. Do extrafloral nectar resources, species abundances, and body sizes contribute to the structure of ant–plant mutualistic networks? *Oecologia* 164:741–750.
- Davidson, D. W., S. C. Cook, R. R. Snelling, and T. H. Chua. 2003. Explaining the abundance of ants in lowland tropical rainforest canopies. *Science* 300:969–972.
- Davidson, D.W., S. C. Cook, and R. R. Snelling. 2004. Liquid eeding performances of ants (Formicidae): ecological and evolutionary implications. *Oecologia* 139:255–66.
- Del–Claro, K., V. Berto, and W. Réu. 1996. Effect of herbivore deterrence by ants increase fruit set in an extrafloral nectary plant *Qualea multiflora* (Vochysiaceae). *Journal of Tropical Ecology* 12:887–892.
- Dormann, C. F., J. Fründ, N. Blüthgen, and B. Gruber. 2009. Indices, graphs and null models: analysing bipartite ecological networks. *The Open Ecology Journal* 2:7–24.
- Dreisig, H. 2000. Defense by exploitation in the Florida carpenter and, *Camponotus floridanus*, at an extrafloral nectar resource. *Behavioral Ecology Sociobiology* 47:274–279.
- Dunne, J. A., R. J. Williams, and N. D. Martinez. 2002. Food–web structure and network theory: the role of connectance and size. *Proceedings of the National Academy of Sciences USA* 99:12917–12922.
- Fischer, J., and D. Lindenmayer. 2002. Treating the nestedness temperature calculator as a black box can lead to false conclusions. *Oikos* 99:193–199.
- Fortuna, M. A., D. B. Stouffer, J. M. Olesen, P. Jordano, D. Mouillot, B. R. Krasnov, R. Poulin, and J. Bascompte. 2010. Nestedness versus modularity in ecological networks: two sides of the same coin? *Journal of Animal Ecology* 79:811–817.

- Fourcassié, V., C. Bredard, K. Volpatti, and G. Theraulaz G. 2003. Dispersion movements in ants: spatial structuring and density-dependent effects. *Behavioural Processes* 63:33–43.
- Fowler, H. G., L. C. Forti, C. R. F. Brandão, J. H. C. Delabie, and H. L. Vasconcelos. 1991. Ecologia nutricional de formigas. Pages 131–223 *in* A. R. Panizzi, and J. R. P. Parra, editors. *Ecologia nutricional de insetos e suas implicações no manejo de pragas*. Editora Manole, São Paulo, BR.
- Guimarães-Jr, P. R., and P. R. Guimarães. 2006. Improving the analyses of nestedness for large sets of matrices. *Environmental Modelling and Software* 21:1512–1513.
- Guimarães, P. R., V. Rico-Gray, S. F. Reis, and J. N. Thompson. 2006. Asymmetries in specialization in ant–plant mutualistic networks. *Proceedings of the Royal Society, B, Biological Sciences* 273:2041–2047
- Guimarães, P. R., C. Sazima, S. F. Reis, and I. Sazima. 2007. The nested structure of marine cleaning symbiosis: is it like flowers and bees? *Biology Letters* 3:51–54.
- Guimarães, P. R., P. Jordano, and J. N. Thompson. 2011. Evolution and coevolution in mutualistic networks. *Ecology Letters* 14: 877–88.
- Guimerà, R., and L. A. N. Amaral. 2005. Functional cartography of complex metabolic networks. *Nature* 433:895–900.
- Heil, M., B. Fiala, B. Baumann, and K. E. Linsenmair. 2000. Temporal, spatial and biotic variations in extrafloral nectar secretion by *Macaranga tanarius*. *Functional Ecology* 14:749–757.
- Heil, M., and D. McKey. 2003. Protective ant–plant interactions as model systems in ecological and evolutionary research. *Annual Review of Ecology, Evolution and Systematics* 34:425– 453.

- Heil, M., A. Hilpert, R. Krüger, and K. E. Linsenmair. 2004. Competition among visitors to extrafloral nectaries as a source of ecological costs of an indirect defence. *Journal of Tropical Ecology* 20:201–208.
- Hölldobler, B., and E. O. Wilson. 1990. *The ants*. Harvard University Press, Cambridge, Massachusetts, USA.
- Jordano, P. 1987. Patterns of mutualistic interactions in pollination and seed dispersal: connectance, dependence, and coevolution. *American Naturalist* 129:657–677.
- Koptur, S., V. Rico-Gray, and M. Palacios-Rios. 1998. Ant protection of the nectaried fern *Polypodium plebeium* in central Mexico. *American Journal of Botany* 85:736–739.
- Krishna, A., P. R. Guimarães, P. Jordano, and J. Bascompte. 2008. A neutral–niche theory of nestedness in mutualistic networks. *Oikos* 117:1609–1618.
- Levin, S. A. 1992. The problem of pattern and scale in ecology. *Ecology* 73:1943–1967.
- Lewinsohn, T. M., P. I. Prado, P. Jordano, J. Bascompte, and J. M. Olesen. 2006. Structure in plant–animal interaction assemblages. *Oikos* 113:174–84.
- Magnusson, W. E., A. P. Lima, R. Luizão, F. Luizão, F. R. C. Costa, C. V. Castilho, and V. P. Kinupp. 2005. RAPELD: a modification of the Gentry method for biodiversity surveys in long-term ecological research sites. *Biota Neotropica* 5:19–24.
- Majer, J. D., and O. Camer-Pesci. 1991. Ants species in tropical australian tree crops and native ecosystems – is there a mosaic? *Biotropica* 23:173–181.
- Maurer, B. A., and B. J. McGill. 2004. Neutral and non-neutral macroecology. *Basic and Applied Ecology* 5:413–422.
- Medan, D., R. P. J. Perazzo, M. Devoto, E. Burgos, M. G. Zimmermann, H. Ceva, and A. M. Delbue. 2007. Analysis and assembling of network structure in mutualistic systems. *Journal of Theoretical Biology* 246:510–521.

- Montoya, J. M., S. L. Pimm, and R. V. Solé. 2006. Ecological networks and their fragility. *Nature* 442:259–264.
- Morales, J. M., and D. P. Vázquez. 2008. The effect of space in plant–animal mutualistic networks: insights from a simulation study. *Oikos* 117:1362–1370.
- Newman, M. E., and M. Girvan. 2004. Finding and evaluating community structure in networks. *Physical Review E* 69:025103.
- Nielsen, A., and J. Bascompte. 2007. Ecological networks, nestedness and sampling effort. *Journal of Ecology* 95:1134–1141.
- Olesen, J. M., J. Bascompte, Y. L. Dupont, and P. Jordano. 2007. The modularity of pollination networks. *Proceedings of the National Academy of Sciences USA* 104:19891–19896.
- Oliveira, P. S., and C. R. F. Brandão. 1991. The ant community associated with extrafloral nectaries in the Brazilian cerrado. Pages 198–212 *in* D. F. Cutler, and C. R. Huxley, editors. *Ant– Plant Interactions*. Oxford University Press, Oxford, UK.
- Oliveira, P. S., V. Rico–Gray, C. Díaz–Castelazo, and C. Castilho–Guevara. 1999. Interaction between ants, extrafloral nectaries and insect herbivores in Neotropical coastal and dunes: herbivore deterrence by visiting ants increases fruit set in *Opuntia stricta* (Cactacea). *Functional Ecology* 13:623–631.
- R Development Core Team. 2010. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rezende, E. L., P. Jordano, and J. Bascompte. 2007. Effects of phenotypic complementarity and phylogeny on the nested structure of mutualistic networks. *Oikos* 116:1919–1929.
- Rico–Gray, V., J. G. Garcia–Franco, M. Palacios–Rios, C. Díaz–Castelazo, V. Parra–Tabla, and J. A. Navarro. 1998. Geographical and seasonal variation in the richness of ant – plant interactions in Mexico. *Biotropica* 30:190–200.

- Rico-Gray, V., and P. S. Oliveira. 2007. The ecology and evolution of ant-plant interactions. University of Chicago Press, Chicago, Illinois, USA.
- Schoereder, J. H., T. G. Sobrinho, M. S. Madureira, C. R. Ribas, and P. S. Oliveira. 2010. The arboreal ant community visiting extrafloral nectaries in the Neotropical cerrado savanna. *Terrestrial Arthropod Reviews* 3:3–27.
- Schupp, E. W., and D. H. D.H. 1991. Phylogeny, lifeform, and habitat dependence of ant-defended plants in a Panamanian forest. Pages 175–197 *in* D. F. Cutler, and C. R. Huxley, editors. *Ant-Plant Interactions*. Oxford University Press, Oxford, UK.
- Stang, M., P. G. L. Klinkhamer, N. M. Waser, I. Stang, and E. Van-der-Meijden. 2009. Size-specific interaction patterns and size matching in a plant-pollinator interaction web. *Annals of Botany* 103:1459–1469.
- Sugiura, S. 2010. Species interactions-area relationships: Biological invasions and network structure in relation to island area. *Proceedings of the Royal Society, B, Biological Sciences* 277:1807–1815.
- Thompson, J. N. 2005. The geographic mosaic of coevolution. University of Chicago Press, Chicago, Illinois, USA.
- Tobin, J. E. 1995. Ecology and diversity of tropical forest canopy ants. Pages 129–147 *in* M. D. Lowman, and N. M. Nadkarni, editors. *Forest canopies*. San Diego Academic Press, San Diego, USA.
- Turner, M. G. 1990. Spatial and temporal analysis of landscape patterns. *Landscape Ecology* 4:21–30.
- Ulrich, W., M. Almeida-Neto, and N. J. Gotelli. 2009. A consumer's guide to nestedness analysis. *Oikos* 118:3–17.
- Vázquez, D. P., and M. A. Aizen. 2004. Asymmetric specialization: a pervasive feature of plant-pollinator interactions. *Ecology* 85:1251–1257.

- Vázquez, D. P., and M. A. Aizen. 2006. Community-wide patterns of specialization in plant–pollinator interactions revealed by null models. Pages 201–219 *in* N. M. Waser, and J. Ollerton, editors. *Plant–Pollinator Interactions: from Specialization to Generalization*. The University of Chicago Press, Chicago, USA.
- Vázquez, D. P.; C. J. Melián, N. M. Williams, N. Blüthgen, B. R. Krasnov, and R. Poulin. 2007. Species abundance and asymmetric interaction strength in ecological networks. *Oikos* 116:1120–1127.
- Vázquez, D. P., N. Blüthgen, L. Cagnolo, and N. P. Chacoff. 2009a. Uniting pattern and process in plant–animal mutualistic networks: a review. *Annals of Botany* 103:1445–1457.
- Vázquez, D. P., N. P. Chacoff, and L. Cagnolo. 2009b. Evaluating multiple determinants of the structure of mutualistic networks. *Ecology* 90:2039–2046.
- Veech, J. A., K. S. Summerville, T. O. Crist, and J. C. Gering. 2002. The additive partitioning of species diversity: recent revival of an old idea. *Oikos* 99:3–9.
- Veloso, H. P., A. L. R. Rangel-Filho, and J. C. A. Lima. 1991. *Classificação da vegetação brasileira, adaptada a um sistema universal*. Editora IBGE, Rio de Janeiro, BR.
- Verdú, M., and A. Valiente-Banuet. 2011. The relative contribution of abundance and phylogeny to the structure of plant facilitation networks. *Oikos* 120:1351–1356.
- Zhou, S., and D. Zhang. 2008. Neutral theory in community ecology. *Frontiers of Biology in China* 3:1–8.
- Williams, R. J., and N. D. Martinez. 2008. Success and its limits among structural models of complex food webs. *Journal of Animal Ecology* 77:512–519.


Figure 1

Figure 1. Relationship for the different metrics used in both ecological networks of ants and plants with and without extrafloral nectaries (EFNs): A) mean number of links for plant specie; B) mean number of links for ant specie; C) connectance; D) modularity*; E) nestedness (NODF metric); F) nestedness (Z-Score metric). Each line represents one of the

12 paired plots. It is also shown the boxplots with the distribution of the data set based on their descriptive parameters. Only the mean number of links for ant specie (B) was not significant (paired t-test: $t = 0.258$, $df = 11$, $P = 0.801$).

* In any of the plots studied were observed modularity significantly higher than expected by the heterogeneity of interactions.


Figure 2

Figure 2. Average percentage of additive partitioning of total diversity of ants collected on plants with and without extrafloral nectaries in 12 plots of an *terra-firme* dense rainforest in the Brazilian Meridional Amazon. α -diversity represents the mean of ants richness collected on plants with and without extra-floral nectaries (EFNs). β -diversity represent the turnover of ant species composition collected on plants with and without EFNs in the plots. The total ant richness in both networks was determined by γ -diversity. The values of the additive partitioning of diversity expected were calculated excluding the effect of the presence of EFNs through 1,000 randomizations of plants classification (presence or absence of EFNs).


Figure 3

Figure 3. Graphical representation of one of the 12 interaction matrices among ants with extrafloral nectaries (EFNs) plants studied. This matrix shows the cases of an interaction between ants and plants mediated by the relative species abundance of both plants and ants. The red cells represents that the interaction occurs less than expected by the abundance, blue cells represents the interaction occurs more than expected by the abundance, white cells represents ant species collected in plants without EFNs was not collected in plants with EFNs.

Supplementary Information

To summarize the composition of the ant community in plants with and without EFNs, we ordered the similarity between points using Non-Metric Multidimensional Scaling. The ordinations analyses were performed from a distance matrix calculated from the Sorensen's dissimilarity index (qualitative data) and Bray-Curtis's dissimilarity index (quantitative data: frequency occurrence of ants on plants). Additionally, we tested the difference in the ant species composition through a permutation test (10,000 permutations) based on analysis of similarities (ANOSIM) (Clarke 1993). Both the ordination and analysis of similarities were made through the software R Development Core Team (version 2.13.1).




Figure 4. Non-metric multidimensional scaling of ants collected in plants with (triangles) and without extrafloral nectaries (squares) in 12 plots of an *terra-firme* dense rainforest in the Brazilian Meridional Amazon. This ordination analysis was calculated from the (A) Sorensen's dissimilarity index (Stress= 0.351 ; Axis 1 + Axis 2= 34.9% of explanation) and (B) Bray-Curtis's dissimilarity index (Stress= 0.318 ; Axis 1 + Axis 2= 43.7% of explanation).

CHAPTER 3

Spatial structure of ant-plant mutualistic networks*

** This manuscript will be submitted to Oikos*

Submitted to Oikos

Running head: Effect of space in ant-plant networks

Spatial structure of ant-plant mutualistic networks

Wesley Dáttilo ^{1*}; Thiago J. Izzo ¹ and Paulo R. Guimarães Jr ²

¹ Department of Ecology and Botany, Community Ecology Lab., Universidade Federal de Mato Grosso, Cuiabá, MT, CEP 78060-900, Brazil.

² Department of Ecology, Instituto de Biociências, Universidade de São Paulo, São Paulo, SP, CEP 05508-900, Brazil.

** Corresponding author:*

Wesley Dáttilo

UFMT-IB-PPGECB

Avenida Fernando Correa/ SN

Cuiabá, Mato Grosso State, 78068-900

Brazil

E-mail wdattilo@hotmail.com

ABSTRACT:

The topological structure of mutualism networks provides insights into ecological and evolutionary dynamics of interacting species. However, only recently has the spatial effect been incorporated as a mechanism structuring mutualistic networks. Here we used the ant-plant mutualistic networks to evaluate for the first time how the topological structure and species turnover of mutualistic networks varies over spatial gradient. Even the fact that ant and plant composition of networks changes over space, the generalist core species and the topological structure of networks remain unaltered. This finding indicates that independently of local and landscape environmental factors the nonrandom pattern of community organization is not changed. In short, contributing thus to our understanding of the maintenance of biodiversity and coevolutionary processes.

Keywords: beta diversity, coevolution, generalist core, nestedness, spatial turnover.

1. Introduction

The study of mutualistic networks has provided important insights into the mechanisms that contribute to the structural organization of plant-animal interactions. (Medan et al. 2007, Morales and Vázquez 2008, Nielsen and Bascompte 2007, Rezende et al. 2007, Stang et al. 2007, Vázquez et al. 2007, 2009). Using measures of graph theory to characterize the network topology, several studies have found non-random patterns of interaction on a wide range of ecosystems (Bascompte et al. 2003, Lewinsohn et al. 2006, Guimarães et al. 2007, Vázquez 2009a). However, little is known as such metrics vary over space (Morales and Vázquez 2008, Vázquez et al. 2009, Burkle and Alarcón 2011). In addition, to understand the consequences of spatial variation in mutualistic networks it is essential to determine how the space modulates the dynamics of interacting species (Burkle and Alarcón, 2011).

Some theoretical and empirical studies have shown that when new species are introduced into a network, they can influence the ecological dynamics of the entire network, mainly because of the dominance hierarchy and species abundance of interactions can be modified (Solé and Montoya 2001, Olesen et al. 2002, Memmott et al. 2007, Aizen et al. 2008, Olesen et al. 2008, Díaz-Castelazo et al. 2004). Such studies have focused on evaluating how the entry of invasive alien species affects the structural organization of the network. However, very little is still known on how the species turnover along space affects the structure of networks.

When we compared this with other systems, eg. plant-pollinator, ant-plant mutualistic networks have been studied on a small scale (Guimarães et al. 2006, 2007, Chamberlain et al. 2010, Sugiura 2010) and the knowledge of space effect on these networks is null. Thus, we used the interactions between ants and plants with extrafloral nectaries (EFNs), to evaluate for the first time how the topological structure of mutualistic networks varies over the spatial gradient. In ant-plant mutualistic networks, plants produce

nutritious liquid for ants (Baker et al. 1978, Rico-Gray and Oliveira 2007), which in exchange for the food provided, the ants defend the plants against potential herbivores (Del-Claro et al. 1996, Oliveira et al. 1999, Rico-Gray and Oliveira 2007).

On the other hand, different parameters can change the nature of ant-plant interactions, such as: competition, abundance and quality of resources, seasonality of nectar production, and other biotic and several abiotic factors (Blüthgen and Fiedler 2004, Díaz-Castelazo et al. 2004, Rico-Gray and Oliveira 2007, Rico-Gray et al. 2011). All these factors can be affected by the spatial distributions of ants on plants (Bentley 1976, Barton 1986, Heil 2000, Apple and Feener 2001, Cogni et al. 2003). In fact, in tropical forests, it is known that several plant species are spatially aggregated (Newbery et al. 1986, Condit et al. 2000, Köhler 2000), and that the foraging and dispersal of ants is limited to small spatial scales (Soares and Schoereder 2001, Fourcassié et al. 2003). So, according to neutral theory the compositional similarity among plant communities will decrease as the distance between two points increases, due to the limited dispersal of organisms and environmental gradients (Hubbel 2001, Chave and Leigh 2002, Gilbert and Lechowicz 2004).

Here we predicted that due to high spatial aggregation of plants in tropical regions and low ant mobility there is a mosaic of interactions with different partners over a relatively small geographic space, and this could generate differences in the topological structure of these networks (Thompson 2005, Morales and Vázquez 2008, Burkle and Alarcón, 2011). In order to test this hypothesis, we collected twelve ant-plant mutualistic networks in a *terra-firme* forest in Brazilian Meridional Amazon and analyzed their network topology. Subsequently, we calculated the dissimilarity of network topology over geographic distance among sampling points in order to examine whether: (1) species turnover over spatial gradient influences the topological structure of ant-plant mutualistic

networks, and (2) the core of generalist species remains stable on a geographic scale of up to 5.099 meters.

2. Material and Methods

2.1) Study area

We conducted this study at São Nicolau Farm (9°48'S e 58°15'W, elev. 254m), located in the municipality of Cotriguaçu, north of Mato Grosso State, Brazil. According to the Köppen classification, the climate is tropical humid (Am) with average annual temperature: 24 °C, humidity 85 %, and 2.300 mm of precipitation (Camargo et al. 2010, Dáttilo et al. 2012). It has two well defined seasons, a rainy season between November and April and dry season between May and October. The area is characterized as a *terra-firme* dense rainforest inserted in the Brazilian Meridional Amazon (Veloso et al. 1991, Camargo et al. 2010). The reserve area covers 7.000 ha of continuous forest, surrounded by a much larger area of intact forest. The terrain is undulating with altitudinal variation of 50 m between the plateaus and the riverside terrains. Canopy trees range from 30-40 m high with some emergent trees which reach 50 m in height. The understory is relatively open, with high frequency of *Orbignya phalerata* Mart. (Arecaceae). In the Brazilian Amazon, it is usual to find between 18 and 53% of plant species in different physiognomies having EFNs, reaching up to 50% coverage of these plants in a given physiognomy (Morellato and Oliveira, 1991, Rico-Gray and Oliveira 2007).

2.2) Data Collection

We collect data in a site (module) managed by the Brazilian Research Program in Biodiversity (PPBio) (PPBio: <http://ppbio.inpa.gov.br>) inserted at São Nicolau Farm. The module consists of six parallel trails in the North-South and two parallel trails East-West.

Every 1 Km trail is one permanent plot with 250 m x 25 m (6.250 m²), total of 12 sampling points in the entire module.

We collected ants and plants between December 2010 and January 2011. In each of the 12 sampling points, we looked for EFN plants reaching from 0.5 m to 3 m height. This size was used because of its easily accessible size to researchers without disturbance. In each plant, we recorded all occurrences of ants collecting liquids in EFN. The plants should be at least 10 meters away from each other, in order to minimize the possibility of collecting ants from the same colony foraging on different plants. Plants and ants were identified to the lowest possible taxonomic level using taxonomic keys and with the help of specialists. Ant specimens were deposited in the Setor de Entomologia of Coleção Zoológica of Universidade Federal de Mato Grosso, Brazil (CEMT).

2.3) *Network topology*

To evaluate how topological properties of ant-plant mutualistic networks vary over space, we calculated the dissimilarity among the 12 sampling points of the following metrics: connectance, network specialization, and nestedness. The connectance (C) is the proportion of possible links that are actually realized (Jordano 1987). We calculated the level of specialization networks using the specialization index (H_2') (ranges from zero (extreme generalization) to one (extreme specialization) through R-Project software version 2.13.1 (bipartite packpage, R Development Core Team 2005). This index is mathematically derived from the Shannon entropy, and based on the deviation from the expected probability distribution of the interactions (Blüthgen et al. 2006). In addition, the index is robust to changes in sampling intensity and the number of interacting species (see more details of this index in Blüthgen et al. 2006, 2007).

We calculated the NODF metric (Nestedness metric based on Overlap and Decreasing Fill) (Almeida-Neto et al. 2008) to estimate the nestedness value of networks,

using ANINHADO software (Guimarães and Guimarães 2006). We tested the nestedness observed for each network with 1.000 networks generated by Null Model II, in order to assess if the nestedness value observed was higher than expected by random patterns of interaction. In this null model, the probability of an interaction occur is proportional to the level of generalization (degree) of plant and animal species (Bascompte et al. 2003). We also calculated the nestedness value standardizing the difference in richness, connectance and heterogeneity of interactions among the sampling points studied using the Z-Score metric to allow cross network comparisons. Z-Score is defined as: $Z_{\text{nodf}} = (x - \mu) / \sigma$, where x = NODF value observed, μ = NODF value of randomized matrices, and σ = is the standard deviation of the randomized matrices (Ulrich et al 2009).

2.4) Spatial turnover of species composition

We calculated the additive partitioning of diversity in ant-plant networks as suggested by Veech et al. (2002) to access the spatial turnover among the sampling points studied on plant and ant species composition (β -diversity). From the total richness of the same trophic level found in two sampling points (γ -diversity), we calculated the α -diversity, defined as: $\alpha = (\alpha_1 + \alpha_2) / 2$, where α_1 = species richness of same trophic level of plot 1, α_2 = species richness of the same trophic level of plot 2). Then, we calculated the β -diversity, defined as: $\beta = (\gamma - \alpha)$. Moreover, we calculated which species belongs to the

generalist core through: $Cg = \left(\frac{x - y}{z} \right) > 1$, where x = mean number of links for given

plant/ant species, y = mean number of links for all plant/ant species in network, z = standard deviation of the number of links for plant/ant species. Thus, we also calculated the turnover of β -diversity (as described above) just for plants and ants inserted on generalist core species.

2.5) Statistical analysis

We used Mantel tests to determine the existence of a relationship between the turnover on different network metrics described above and the matrices of geographic distances among all the sampling points studied. We conducted these tests using the *vegan* package (Oksanen et al. 2007) in the R-Project software version 2.13.1 (R Development Core Team 2005) using Euclidean distance to calculate the dissimilarity in the metrics and geographic distances among sampling points. In these analyses, we also tested the correlation coefficient (r) using this analysis. We made all graphics using the software GraphPad Prism version 5.0 (Motulsky, 1999).

3. Results

In this study, we recorded 72 plant species (or morphospecies) with EFNs, belonging to 24 genera and 16 families. The Bignoniaceae family corresponded to 26.3% of plant species, followed by 22.8% of Mimosaceae and 10.5% of Caesalpiniaceae. The plant species richness per sampling points was 21.41 ± 3.77 (Mean \pm SD). For ants, we recorded 149 species, inserted into 23 genera and eight subfamilies. The subfamily Myrmicinae corresponded to 42.28% of ant species, followed by 26.1% for Formicinae and 14.9% for Dolichoderinae. The ant species richness per sampling points was 23.16 ± 5.85 . The mean and standard deviation of metrics used in this study were: Connectance: (0.140 ± 0.035), Network specialization: (0.088 ± 0.049), NODF: (21.01 ± 4.406), nestedness Z-Scores (3.63 ± 1.50).

In the spatial scale studied, we did not find significant correlation of dissimilarity of geographic distance with the dissimilarity of connectance (Mantel statistic $r = 0.044$, $P = 0.374$) and network specialization (H_2') (Mantel statistic $r = 0.004$, $P = 0.457$) (Figure 1A-B). When analyzed we had different results in the two metrics that describe the nested

pattern in ant-plant interactions. We did not observe significant correlation of dissimilarity of nestedness calculated by NODF metric with geographic distance (Mantel statistic $r = 0.078$, $P = 0.283$). However, we observed significant correlation of dissimilarity of geographic distance with the dissimilarity of nestedness calculated by Z-Score metric (Mantel statistic $r = 0.315$, $P < 0.01$) (Figure 1C-D).

We observed a turnover of species composition (β -diversity), for both plants and ants, along the geographic distance (Plants: Mantel statistic $r = 0.401$, $P < 0.01$, Ants: Mantel statistic $r = 0.307$, $P = 0.013$) (Figure 2A-B). However, we did not observe a turnover on core species composition, for both plants and ants, along the geographic distance (Plants: Mantel statistic $r = 0.007$, $P = 0.437$, Ants: Mantel statistic $r = -0.088$, $P = 0.734$) (Figure 2C-D). In addition, the number of ant species present in the generalist core of networks ranged between one and three species, and the species *Azteca* sp2, *Brachymyrmex* sp1 and *Crematogaster* sp8 were present in the core in more than 58 % of sampling points. For plants, the number of species present in the generalist core of networks ranged between one and two species, and the species *Inga* sp12, *Mabea* sp2, *Protium* sp1, *Stryphnodendron* sp1 were present in the core in more than 66 % of sampling points.

4. Discussion

We know from previous studies that mutualistic networks of free-living species show a nested and asymmetrical pattern in different habitats and ecosystems such as plant-pollinator, fruit-frugivore, ant-plant, clownfish-anemone and marine fish cleaning symbioses (Bascompte et al. 2003, Guimarães et al. 2006, Guimarães et al. 2007, Ollerton et al. 2007, Mello et al. 2011). However, how the topological structure of these networks varies over spatial gradients has often been ignored. The few studies that evaluated the role

of spatial variations in mutualistic networks, show us that local and landscape environmental factors, beyond spatial aggregation and animal mobility are important factors that structure the plant-animal interactions (Morales and Vázquez 2008, Burkle and Alarcón 2011). In this study, standardizing the network metrics, collection effort and habitat, we showed that in ant-plant mutualistic networks, the topological structure remains unaltered in the spatial scale studied. In addition, even the fact that ant and plant composition of networks changes over space, the generalist core species remain stable along the 5.099 m on a *terra-firme* forest in Brazilian Meridional Amazon.

In tropical rainforests, the main factors that explain the distribution and diversity of ants are competitive interactions, habitat complexity (abundance of food and nesting sites), climate stability and natural barriers that prevent ant queen dispersal (Benson and Harada 1988, Hölldobler and Wilson 1990, Reyes-Lopes et al. 2003). Thus, along a spatial scale, different biotic and abiotic factors can influence the richness and diversity of interactions between ants and plants differently (Brühl et al. 1999, Díaz-Castelazo et al. 2004, Rico-Gray and Oliveira 2007, Rico-Gray et al. 2011). In this study, there was a wide variation in the values of connectance among sampling points, influencing the values of nestedness using NODF metric. In fact, when we calculated the dissimilarity among these metrics, there was no relation with geographic distance. On the other hand, when we controlled the connectance effects, we found different patterns and nestedness values using Z-Score metric, since the dissimilarity of nestedness increased with geographic distance. So why are closer sampling points more similar? This can be explained because closer sampling points have lower turnover of plant and ant composition, as we observed. Therefore, it was expected that the number of ant and plant species, beyond number of interactions would be more similar in closer sampling points. Additionally, the metric used to calculate the network specialization is also robust with respect to networks with different connectance values (Blüthgen et al. 2006, 2007, 2010), which indicated us that the low specialization

found here is a non-random pattern of ant-plant mutualistic networks. Thus, we showed that the topological structure of ant-plant mutualistic networks is stable and predictable over a spatial scale of up to 5.099 meters, independent of variations in biotic and abiotic factors of sampling points studied.

Interestingly, we showed that the core of generalist species remained stable over the spatial scale studied. One factor that could explain the stability of the core over space is the species abundance, where abundant species could interact most frequently with each other and with other less abundant species (Vázquez et al. 2007, 2009b). However, in a previous paper by Dáttilo et al. *in preparation*, showed that in ant-plant mutualistic networks, the generalist core interact with each other more than expected by their abundances. This finding indicates that ants of the core possibly have mechanisms which maintain such species in the core over space. Due to the fact that competition for resource is an important factor in the structuring of ant communities in tropical regions (Djieto-Lordon & Dejean. 1999, Delabie et al 2000), we believe that the core of generalist ant species are competitively superior (i.e. more recruitment orientated or aggressive) and monopolize resources. In addition, we propose that this generalist core is less variable in time and space than the network periphery as originally proposed by Bascompte et al. (2003). Moreover, the interaction strength among interacting partners of generalist core species is symmetrical (Bascompte et al. 2006, Vázquez et al. 2007, Guimarães et al. 2011). This implies that the generalist core can act as a coevolutionary vortex for more specialized interactions, where the convergence of traits on both sides of the interaction increases over evolutionary time (Bascompte et al. 2003; Thompson 2005; Guimarães et al. 2007). The existence of a geographic stable hard-core enhances the coevolutionary importance of such a vortex, once several individuals of ants and plants that belong to the core, in a given region, will have an increased chance of interacting with each-other. Additionally, Díaz-Castelazo et al. (2010), showed that the proportion of generalist core species in ant-plant

interaction did not change in 10 years on a local scale. Thus, the generalist core being stable over space and time at different spatial scales, can generate the geographic mosaic of coevolution, where the number of interacting species increases over evolutionary time (Thompson 2005).

In conclusion, our study shows that in general, the topological structure of ant-plant mutualistic networks is stable over a relatively small geographic space inserted in a *terra-firme* forest in Brazilian Meridional Amazon. Moreover, even that a turnover on interacting ant and plant composition changes along this spatial scale, the generalist core remains stable. This finding indicates that independently of local and landscape environmental factors the nonrandom pattern of community organization is not changed. In short, contributing thus to our understanding as to the maintenance of the biodiversity and coevolutionary processes. However, it is not known what the consequences of temporal variations and resilience of the generalist core in larger spatial scales is. There is still much to do to better understand patterns and processes related to spatiotemporal variation in ant-plant systems. Finally, other empirical studies that assess the spatial effect in different mutualistic networks are essential to understanding the current view of coevolutionary processes in mutualistic interactions.

Acknowledgements

We would like to thank Jéssica Falcão for your help during the fieldwork and for their comments on earlier versions this manuscript. We also thank Office National des Forêts Brazil by logistical support and Brazilian Program for Biodiversity Research by financial support. WD is granted by a Coordenação de Aperfeiçoamento de Pessoal de Nível Superior master's fellowship. PRG acknowledge the financial support of Fundação

de Amparo à Pesquisa do Estado de São Paulo. This is publication XXX in the Núcleo de Estudos da Biodiversidade da Amazônia Mato-Grossense technical series.

References

- Aizen, M. et al. 2008. Invasive mutualists erode native pollination webs. – PLoS Biol. 6: e31.
- Almeida-Neto, M. et al. 2008. A consistent metric for nestedness analysis in ecological systems: reconciling concept and measurement. – Oikos 117:1227–1239.
- Apple, J. L. and Feener-Jr, D. H. 2001. Ant visitation of extrafloral nectaries of *Passiflora*: the effects of nectary attributes and ant behavior on patterns in facultative ant–plant mutualisms. – Oecologia 127: 409–416.
- Baker, H. G. et al. 1978. A comparison of the amino acid complements of floral and extrafloral nectars. – Bot. Gaz. 139: 322–332.
- Barton, A. M. 1986. Spatial variation in the effect of ants on an extrafloral nectary plant. – Ecology 67: 495–504.
- Bentley, B. L. 1976. Plants bearing extrafloral nectaries and the associated ant community: inter-habitat differences in the reduction of herbivore damage. – Ecology 57: 815–820.
- Bascompte, J. et al. 2003. The nested assembly of plant–animal mutualistic networks. – Proc. Natl. Acad. Sci. 100: 9383–9387.
- Bascompte, J. et al. 2006. Asymmetric coevolutionary networks facilitate biodiversity maintenance. – Science 312: 431–433.
- Benson, W. and Harada, A. Y. 1988. Local diversity of tropical and temperate ant faunas (Hymenoptera: Formicidae). – Acta Amaz. 18: 275–289.

- Blüthgen, N. and Fiedler, K. 2004 Competition for composition: lessons from nectar-feeding ant communities. – *Ecology* 85: 1479–1485.
- Blüthgen, N. et al. 2006. Measuring specialization in species interaction networks. – *BMC Ecol.* 6: 9–12.
- Blüthgen, N. et al. 2007. Specialization, constraints, and conflicting interests in mutualistic networks. – *Cur. Biol.* 17: 341–346.
- Blüthgen, N. 2010. Why network analysis is often disconnected from community ecology: A critique and an ecologist's guide. – *Basic Appl. Ecol.* 11: 185–195.
- Burkle, L. A. and Irwin, R. 2009. The importance of interannual variation and bottom-up nitrogen enrichment for plant–pollinator networks. – *Oikos* 118: 1816–1829.
- Burkle, L. A. and Alarcón, R. 2011. The future of plant–pollinator diversity: understanding interaction networks across time, space, and global change. – *Am. J. Bot.* 98:1–11.
- Brühl, C. A. et al. 1999. Altitudinal distribution of leaf litter ants along a transect in primary forests on Mount Kinabalu, Sabah, Malaysia. – *J. Trop. Ecol.* 15: 265–277.
- Camargo, F. F. et al. 2010. Variabilidade genética para caracteres morfométricos de matrizes de castanha-do-brasil da Amazônia Mato-grossense. – *Acta Amaz.* 40: 705–710.
- Carlo, T. A. et al. 2007. Plant–frugivore interactions as spatially explicit networks: integrating frugivore foraging with fruiting plant spatial patterns. – In: Dennis, A. J. et al. (eds), *Seed dispersal: theory and its application in a changing world*. Cabi Publishing, pp 369–390.
- Chamberlain, S. A. et al. 2010. Do extrafloral nectar resources, species abundances, and body sizes contribute to the structure of ant–plant mutualistic networks? – *Oecologia* 164: 741–750.

- Chave, J. and Leigh, E. G. 2002. A spatially explicit neutral model of [beta]–diversity in tropical forests. – *Theor. Pop. Biol.* 62: 153–168.
- Condit, R. et al. 2000. Spatial patterns in the distribution of tropical tree species. – *Science* 288: 1414–1418.
- Cogni, R. et al. 2003. Interhabitat differences in ant activity on plant foliage: ants at extrafloral nectaries of *Hibiscus pernambucensis* in sandy and mangrove forests. – *Ent. Exp. App.* 107: 125–131.
- Del–Claro, K. et al. 1996. Effect of herbivore deterrence by ants increase fruit set in an extrafloral nectary plant, *Qualea multiflora* (Vochysiaceae). – *J. Trop. Ecol.* 12: 887–892.
- Díaz–Castelazo, C. et al. 2004. Extrafloral nectary–mediated ant–plant interactions in the coastal vegetation of Veracruz, México: Richness, occurrence, seasonality, and ant foraging patterns. – *Ecoscience* 11: 472–482.
- Fourcassié, V. et al. 2003. Dispersion movements in ants: spatial structuring and density–dependent effects. – *Beh. Proc.* 63: 33–43.
- Gilbert, B. and Lechowicz, M. J. 2004. Neutrality, niches, and dispersal in a temperate forest understory. – *Proc. Natl. Acad. Sci.* 101: 7651–7656.
- Guimarães, P. R. Jr. and Guimarães, P. R. 2006. Improving the analyses of nestedness for large sets of matrices. – *Environ. Model. Soft.* 21: 1512–1513.
- Guimarães, P. R. Jr. et al. 2006. Asymmetries in specialization in ant–plant mutualistic networks. – *Proc. R. Soc. B.* 273: 2041–2047.
- Guimarães, P. R. Jr. et al. 2007. The nested structure of marine cleaning symbiosis: is it like flowers and bees? – *Biol. Lett.* 3: 51–54.
- Guimarães, P. R. Jr. et al. 2011. Evolution and coevolution in mutualistic networks. – *Ecol. Lett.* 14: 877–88.

- Heil, M. et al. 2000. Temporal, spatial and biotic variations in extrafloral nectar secretion by *Macaranga tanarius*. – *Func. Ecol.* 14:749–757.
- Hölldobler, B. and Wilson, E. O. 1990. The ants. – Harvard Univ. Press.
- Hubbell, S. P. 2001. The Unified Theory of Biodiversity and Biogeography. – Princeton Univ. Press.
- Jordano, P. 1987. Patterns of mutualistic interactions in pollination and seed dispersal: connectance, dependence, and coevolution. *Am. Nat.* 129: 657–677.
- Köhler, P. et al. 2000. Concepts for the aggregation of tropical tree species into functional types and the application to Sabah's lowland rain forest. – *J. Trop. Ecol.* 16: 591–602.
- Longino, J. T. et al. 2002. The ant fauna of a tropical rain forest: estimating species richness in three different ways. – *Ecology* 83: 689–702.
- Medan, D. et al. 2007. Analysis and assembling of network structure in mutualistic systems. – *J. Theor. Biol.* 246: 510–521.
- Mello, M. A. R. et al. 2011. The modularity of seed dispersal: differences in structure and robustness between bat- and bird-fruit networks. – *Oecologia* 167: 131–140.
- Morales, J. M. and Vázquez, D. P. 2008. The effect of space in plant-animal mutualistic networks: insights from a simulation study. – *Oikos* 117: 1362–1370.
- Memmott, J. et al. 2007. Global warming and the disruption of plant-pollinator interactions. – *Ecol. Lett.* 10: 710–717.
- Morellato, L. P. C. and Oliveira, P. S. 1991. Distribution of extrafloral nectaries in different vegetation types of Amazonian Brazil. – *Flora* 185: 33–38.
- Motulsky, H. 1999. Analyzing data with graph pad prism software. – GraphPad Soft.
- Newbery, D. M. C. et al. 1986, Spatial pattern of trees in Kerangas forests, Sarawak. – *Vegetatio* 65: 77–89.

- Nielsen, A. and Bascompte, J. 2007. Ecological networks, nestedness and sampling effort. – J. Ecol. 95: 1134–1141.
- Oksanen, J. et al. 2007. Vegan: community ecology package. R package version 1.8–5. <<http://cran.r-project.org>>
- Olesen, J. M. and Jordano, P. 2002. Geographic patterns in plant–pollinator mutualistic networks. – Ecology 83: 2416–2424.
- Olesen, J. M. et al. 2002. Invasion of pollination networks on oceanic islands: Importance of invader complexes and endemic super generalists. – Div. & Dist. 8: 181–192.
- Olesen, J. M. et al. 2008. Temporal dynamics of a pollination network. – Ecology 89: 1573–1582.
- Oliveira, P. S. et al. 1999. Interaction between ants, extrafloral nectaries and insect herbivores in Neotropical coastal and dunes: herbivore deterrence by visiting ants increases fruit set in *Opuntia stricta* (Cactaceae). – Func. Ecol. 13: 623–631.
- Ollerton, J. et al. 2007. Pollination niche overlap between a parasitic plant and its host. – Oecologia 151: 473–485.
- Pemberton, P. W. 1998. The occurrence and abundance of plants with extrafloral nectaries, the basis for antiherbivore defensive mutualisms, along a latitudinal gradient in east Asia. – J. Biog. 25: 661–668.
- R Development Core Team 2005. R: A language and environment for statistical computing. Vienna, R Foundation for Statistical Computing. <<http://www.r-project.org>>
- Rezende, E. L. et al. 2007. Effects of phenotypic complementarity and phylogeny on the nested structure of mutualistic networks. – Oikos 116:1919–1929.
- Reyes–Lopes, J. et al. 2003. Community structure of ground–ants: the role of single trees in a Mediterranean pastureland. – Oecologia 24: 195–202.

- Rico-Gray, V. and Oliveira, P. S. 2007. The ecology and evolution of ant-plant interactions. – Univ. of Chicago Press.
- Rico-Gray, V. et al. 2011. Abiotic factors shape temporal variation in the structure of an ant-plant network. – *Arth-Plant. Interac.* (in press)
- Soares, S. M. and Schoereder, J. H. 2001. Ant-nest distribution in a remnant of tropical rainforest in southeastern Brazil. – *Insect. Soc.* 48: 280–286.
- Solé, R. V. and Montoya, J. M. 2001. Complexity and fragility in ecological networks. – *Proc. R. Soc. B.* 268: 2039–2045.
- Stang, M. et al. 2007. Asymmetric specialization and extinction risk in plant-flower visitor webs: a matter of morphology or abundance? – *Oecologia* 151: 442–453.
- Sugiura, S. 2010. Species interactions–area relationships: Biological invasions and network structure in relation to island area. – *Proc. R. Soc. B.* 277: 1807–1815.
- Thompson, J. N. 2005. The geographic mosaic of coevolution. – Univ. of Chicago Press.
- Ulrich, W. et al. 2009. A consumer's guide to nestedness analysis. – *Oikos* 118: 3–17.
- Vázquez, D. P. et al. 2007. Species abundance and asymmetric interaction strength in ecological networks. – *Oikos* 116: 1120–1127.
- Vázquez, D. P. et al. 2009. Uniting pattern and process in plant-animal mutualistic networks: a review. – *Ann. Bot.* 103: 1445–1457.
- Veech, J. A. et al. 2002. The additive partitioning of species diversity: recent revival of an old idea. – *Oikos* 99: 3–9.
- Veloso, H.P. et al. 1991. Classificação da vegetação brasileira, adaptada a um sistema universal. – Edit. IBGE.

Figures




Figure 1. Relationship among the dissimilarity of: a) connectance, b) network specialization, c) nestedness by NODF metric and d) nestedness by Z-Score metric, with dissimilarity of geographic distance of 12 plots collected in the São Nicolau Farm, Mato Grosso State, Brazilian Meridional Amazon. Correlation coefficient (r) and significance (P computed using Mantel tests) also are shown. ($n = 66$ points in each of the metrics calculated).




Figure 2. Relationship among the dissimilarity of: a) plants species composition, b) ants species composition, c) plants core composition and b) ants core composition, with dissimilarity of geographic distance of 12 plots collected in the São Nicolau Farm, Mato Grosso State, Brazilian Meridional Amazon. Correlation coefficient (r) and significance (P computed using Mantel tests) also are shown. ($n = 66$ points in each of the metrics calculated).

FINAL CONCLUSION

Here I showed that the topological structure of networks generated through deterministic process (plants with EFNs) is different than that of the networks generated by the natural abundance of ants on vegetation. Moreover, the core of generalist species in the ant and plant networks with EFNs interact more among themselves than expected by their abundance. Moreover, standardizing the network metrics, collection effort and habitat, I showed that in ant-plant mutualistic networks, the topological structure remains unaltered in the spatial scale studied. In addition, even the fact that ant and plant composition of networks changes over space, the generalist core species remain stable along the 5.099 m on a *terra-firme* forest in Brazilian Meridional Amazon.

APPENDIX 1

List of 238 plant species with and without extrafloral nectaries (EFNs) collected in 12 plots (250 x 25 m) in an terra-firme forest in the Brazilian Meridional Amazon located in the municipality of Cotriguaçu, north of Mato Grosso State, Brazil. The plants were collected between between December 2010 and January 2011.

| | with EFNs | without EFNss |
|-------------------------|-----------|---------------|
| Anacardiaceae | | |
| <i>Astronium</i> sp1 | | X |
| <i>Tapirira</i> sp1 | | X |
| <i>Thyrsodium</i> sp1 | | X |
| Anonaceae | | |
| <i>Annona</i> sp1 | | X |
| <i>Annona</i> sp2 | | X |
| <i>Duguetia</i> sp1 | | X |
| <i>Xylopia</i> sp1 | | X |
| <i>Xylopia</i> sp2 | | X |
| <i>Xylopia</i> sp3 | | X |
| Apocynaceae | | |
| <i>Aspidosperma</i> sp1 | | X |
| <i>Aspidosperma</i> sp2 | | X |
| <i>Aspidosperma</i> sp3 | | X |
| Bignoniaceae | | |
| <i>Jacaranda</i> sp1 | X | |
| <i>Jacaranda</i> sp2 | | X |
| Unidentified sp1 | X | |
| Unidentified sp2 | X | |
| Unidentified sp3 | X | |
| Unidentified sp4 | | X |
| Unidentified sp5 | X | |
| Unidentified sp6 | X | |
| Unidentified sp7 | X | |
| Unidentified sp8 | X | |

| | | |
|--------------------------|---|---|
| Unidentified sp9 | | X |
| Unidentified sp10 | | X |
| Unidentified sp11 | X | |
| Unidentified sp12 | X | |
| Unidentified sp13 | X | |
| Unidentified sp14 | | X |
| Unidentified sp15 | X | |
| Unidentified sp16 | X | |
| Unidentified sp17 | X | |
| Unidentified sp18 | | X |
| Unidentified sp19 | X | |
| Unidentified sp20 | | X |
| Bixaceae | | |
| <i>Bixa</i> sp1 | | X |
| Bombacaceae | | |
| <i>Eriotheca</i> sp1 | | X |
| Burseraceae | | |
| <i>Protium pilosum</i> | X | |
| <i>Protium</i> sp1 | X | |
| <i>Trattinnickia</i> sp1 | | X |
| <i>Trattinnickia</i> sp2 | | X |
| Caesalpinaceae | | |
| <i>Bauhinia</i> sp1 | X | |
| <i>Bauhinia</i> sp2 | X | |
| <i>Bauhinia</i> sp3 | X | |
| <i>Bauhinia</i> sp4 | X | |
| <i>Bauhinia</i> sp5 | X | |
| <i>Bauhinia</i> sp6 | | X |
| <i>Hymenaea</i> sp1 | | X |
| <i>Unidentified</i> sp1 | X | |
| <i>Unidentified</i> sp2 | | X |
| <i>Tachigali</i> sp1 | | X |
| <i>Tachigali</i> sp2 | | X |
| <i>Tachigali venusta</i> | | X |

Celastraceae

| | | |
|--------------------------|---|--|
| <i>Cheiloclinium</i> sp1 | X | |
|--------------------------|---|--|

| | | |
|------------------------|--|---|
| <i>Hypocrateae</i> sp1 | | X |
|------------------------|--|---|

Chrysobalanaceae

| | | |
|---------------------|--|---|
| <i>Hirtella</i> sp1 | | X |
|---------------------|--|---|

| | | |
|---------------------|--|---|
| <i>Hirtella</i> sp2 | | X |
|---------------------|--|---|

| | | |
|--------------------|--|---|
| <i>Licania</i> sp1 | | X |
|--------------------|--|---|

Combretaceae

| | | |
|----------------------|---|--|
| <i>Combretum</i> sp1 | X | |
|----------------------|---|--|

Costaceae

| | | |
|-------------------|---|--|
| <i>Costus</i> sp1 | X | |
|-------------------|---|--|

| | | |
|-------------------|---|--|
| <i>Costus</i> sp2 | X | |
|-------------------|---|--|

| | | |
|-------------------|---|--|
| <i>Costus</i> sp3 | X | |
|-------------------|---|--|

Cucurbitaceae

| | | |
|------------------|---|--|
| Unidentified sp1 | X | |
|------------------|---|--|

Elaeocarpaceae

| | | |
|--------------------|--|---|
| <i>Sloanea</i> sp1 | | X |
|--------------------|--|---|

| | | |
|--------------------|--|---|
| <i>Sloanea</i> sp2 | | X |
|--------------------|--|---|

| | | |
|--------------------|--|---|
| <i>Sloanea</i> sp3 | | X |
|--------------------|--|---|

Euphorbiaceae

| | | |
|------------------|--|---|
| <i>Hevea</i> sp1 | | X |
|------------------|--|---|

| | | |
|------------------|---|--|
| <i>Mabea</i> sp1 | X | |
|------------------|---|--|

| | | |
|------------------|--|---|
| <i>Mabea</i> sp3 | | X |
|------------------|--|---|

| | | |
|------------------|---|--|
| Unidentified sp1 | X | |
|------------------|---|--|

| | | |
|------------------|---|--|
| Unidentified sp1 | X | |
|------------------|---|--|

Fabaceae

| | | |
|---------------------|--|---|
| <i>Dipteryx</i> sp1 | | X |
|---------------------|--|---|

| | | |
|----------------------|---|--|
| <i>Erythrina</i> sp2 | X | |
|----------------------|---|--|

| | | |
|-----------------------|---|--|
| <i>Machaerium</i> sp1 | X | |
|-----------------------|---|--|

| | | |
|-----------------------|---|--|
| <i>Machaerium</i> sp2 | X | |
|-----------------------|---|--|

| | | |
|-----------------------|---|--|
| <i>Machaerium</i> sp3 | X | |
|-----------------------|---|--|

| | | |
|-----------------------|--|---|
| <i>Machaerium</i> sp4 | | X |
|-----------------------|--|---|

| | | |
|------------------|--|---|
| Unidentified sp1 | | X |
|------------------|--|---|

| | | |
|--------------------|--|---|
| <i>Ormosia</i> sp1 | | X |
|--------------------|--|---|

| | | |
|-------------------------|---|---|
| <i>Platymiscium</i> sp1 | | X |
| <i>Platymiscium</i> sp2 | | X |
| <i>Platymiscium</i> sp3 | | X |
| <i>Vatairea</i> sp1 | X | |
| Flacourtiaceae | | |
| <i>Casearia</i> sp1 | | X |
| <i>Casearia</i> sp3 | | X |
| <i>Casearia</i> sp3 | X | |
| <i>Laetia</i> sp1 | | X |
| <i>Laetia</i> sp2 | | X |
| <i>Laetia</i> sp3 | | X |
| Heliconiaceae | | |
| <i>Heliconia</i> sp1 | | X |
| Humiriaceae | | |
| <i>Vantanea</i> sp1 | | X |
| Lacistemataceae | | |
| <i>Lacistema</i> sp1 | | X |
| Lauraceae | | |
| <i>Aniba</i> sp1 | | X |
| <i>Unidentified</i> sp1 | | X |
| <i>Unidentified</i> sp2 | | X |
| <i>Ocotea</i> sp1 | | X |
| <i>Ocotea</i> sp2 | | X |
| <i>Lecytidaceae</i> | | |
| <i>Couratari</i> sp1 | X | |
| Malpighiaceae | | |
| <i>Byrsonima</i> sp1 | | X |
| Marantaceae | | |
| <i>Ischnosiphon</i> sp1 | | X |
| Melastomataceae | | |
| <i>Bellucia</i> sp1 | | X |
| <i>Miconia</i> sp1 | | X |
| Meliaceae | | |
| <i>Guarea</i> sp1 | | X |

| | | |
|----------------------------|---|---|
| <i>Guarea</i> sp2 | | X |
| <i>Guarea</i> sp3 | | X |
| <i>Guarea</i> sp4 | | X |
| <i>Trichilia micrantha</i> | | X |
| <i>Trichilia pallida</i> | | X |
| <i>Trichilia</i> sp1 | | X |
| <i>Trichilia</i> sp2 | | X |
| Memecylaceae | | |
| <i>Mouriri</i> sp1 | | X |
| Menispermaceae | | |
| <i>Abuta grandifolia</i> | | X |
| <i>Abuta</i> sp1 | | X |
| Mimosaceae | | |
| <i>Abarema</i> sp1 | X | |
| <i>Abarema</i> sp1 | X | |
| <i>Enterolobium</i> sp1 | X | |
| <i>Inga</i> sp1 | X | |
| <i>Inga</i> sp2 | X | |
| <i>Inga</i> sp3 | X | |
| <i>Inga</i> sp4 | X | |
| <i>Inga</i> sp5 | X | |
| <i>Inga</i> sp6 | X | |
| <i>Inga</i> sp7 | X | |
| <i>Inga</i> sp8 | X | |
| <i>Inga</i> sp9 | X | |
| <i>Inga</i> sp10 | X | |
| <i>Inga</i> sp11 | X | |
| <i>Inga</i> sp12 | X | |
| <i>Inga</i> sp13 | X | |
| <i>Inga</i> sp14 | X | |
| <i>Inga</i> sp15 | X | |
| <i>Inga</i> sp16 | | X |
| <i>Inga</i> sp17 | | X |
| <i>Mimosa</i> sp1 | X | |

| | | |
|--------------------------------------------|---|---|
| <i>Mimosa</i> sp2 | X | |
| <i>Parkia</i> sp1 | | X |
| <i>Parkia</i> sp2 | | X |
| <i>Samanea</i> sp1 | X | |
| <i>Senegalia</i> sp1 | X | |
| <i>Stryphnodendron</i> sp1 | X | |
| Moraceae | | |
| <i>Brosimum</i> sp1 | | X |
| <i>Brosimum</i> sp2 | | X |
| <i>Brosimum</i> sp3 | | X |
| <i>Ficus</i> sp1 | | X |
| <i>Ficus</i> sp2 | | X |
| <i>Ficus</i> sp3 | | X |
| <i>Ficus</i> sp4 | | X |
| <i>Ficus</i> sp5 | | X |
| <i>Pseudolmedia</i> cf <i>laevigata</i> | | X |
| <i>Pseudolmedia</i> sp1 | | X |
| <i>Pseudolmedia</i> sp2 | | X |
| <i>Sorocea</i> sp1 | | X |
| Myristicaceae | | |
| <i>Cybianthus</i> sp1 | | X |
| <i>Virola</i> sp1 | | X |
| <i>Virola</i> sp2 | | X |
| Nyctaginaceae | | |
| <i>Guapira</i> sp1 | | X |
| Ochnaceae | | |
| <i>Cespedezia</i> sp1 | | X |
| Olacaceae | | |
| <i>Dulacia</i> sp1 | | X |
| Piperaceae | | |
| <i>Piper</i> sp1 | | X |
| <i>Piper</i> sp2 | | X |
| <i>Piper</i> sp3 | | X |

| | | |
|---------------------------|---|---|
| <i>Piper</i> sp4 | | X |
| Polygonaceae | | |
| <i>Coccoloba</i> sp1 | | X |
| Quiinaceae | | |
| <i>Quiina pteridofila</i> | | X |
| Rubiaceae | | |
| <i>Capirona</i> sp1 | | X |
| <i>Duroia</i> sp1 | | X |
| Unidentified sp1 | | X |
| <i>Palicourea</i> sp1 | X | |
| <i>Palicourea</i> sp2 | | X |
| <i>Palicourea</i> sp3 | | X |
| <i>Palicourea</i> sp4 | | X |
| <i>Psychotria</i> sp1 | | X |
| <i>Remijia amazonica</i> | | X |
| <i>Remijia</i> sp1 | | X |
| <i>Remijia</i> sp2 | | X |
| <i>Uncaria</i> sp1 | X | |
| Rutaceae | | |
| <i>Esenbeckia</i> sp1 | | X |
| <i>Metrodorea</i> sp1 | | X |
| <i>Metrodorea</i> sp2 | | X |
| <i>Zanthoxylum</i> sp1 | | X |
| Sapindaceae | | |
| <i>Paullinia</i> sp1 | | X |
| <i>Serjania</i> sp1 | | X |
| <i>Thalisia</i> sp1 | | X |
| <i>Thalisia</i> sp2 | | X |
| <i>Thalisia</i> sp3 | X | |
| Sapotaceae | | |
| <i>Ecclinusa</i> sp1 | | X |
| <i>Ecclinusa</i> sp2 | | X |
| <i>Ecclinusa</i> sp3 | | X |
| <i>Ecclinusa</i> sp4 | | X |

| | | |
|---------------------------|---|---|
| <i>Manilkara hulberia</i> | | X |
| <i>Manilkara</i> sp1 | | X |
| <i>Manilkara</i> sp2 | | X |
| <i>Pouteria</i> sp1 | | X |
| <i>Pouteria</i> sp2 | | X |
| <i>Pouteria</i> sp3 | | X |
| <i>Pouteria</i> sp4 | | X |
| Simaroubaceae | | |
| <i>Simarouba amara</i> | | X |
| Siparunaceae | | |
| <i>Siparuna</i> sp1 | | X |
| <i>Siparuna</i> sp2 | | X |
| <i>Siparuna</i> sp3 | | X |
| Solanaceae | | |
| Unidentified sp1 | | X |
| Unidentified sp2 | | X |
| Sterculiaceae | | |
| <i>Theobroma</i> sp1 | | X |
| <i>Theobroma</i> sp2 | | X |
| <i>Theobroma</i> sp3 | | X |
| <i>Theobroma</i> sp4 | | X |
| Theophrastaceae | | |
| <i>Clavija</i> sp1 | | X |
| Tiliaceae | | |
| <i>Luehea</i> sp1 | | X |
| Ulmaceae | | |
| <i>Celtis</i> sp1 | | X |
| <i>Trema micrantha</i> | X | |
| Urticaceae | | |
| <i>Urera</i> sp1 | X | |
| <i>Urera</i> sp1 | | X |
| Vouchysiaceae | | |
| <i>Vouchysia</i> sp1 | | X |

| | | |
|-------------------|---|---|
| Unidentified sp1 | | X |
| Unidentified sp2 | | X |
| Unidentified sp3 | | X |
| Unidentified sp4 | | X |
| Unidentified sp5 | | X |
| Unidentified sp6 | | X |
| Unidentified sp7 | | X |
| Unidentified sp8 | | X |
| Unidentified sp9 | X | |
| Unidentified sp10 | | X |
| Unidentified sp11 | | X |
| Unidentified sp12 | | X |
| Unidentified sp13 | X | |
| Unidentified sp14 | | X |
| Unidentified sp15 | | X |
| Unidentified sp16 | X | |
| Unidentified sp17 | X | |
| Unidentified sp18 | | X |
| Unidentified sp19 | | X |
| Unidentified sp20 | | X |
| Unidentified sp21 | | X |
| Unidentified sp22 | X | |
| Unidentified sp23 | | X |
| Unidentified sp24 | | X |
| Unidentified sp25 | | X |
| Unidentified sp26 | | X |
| Unidentified sp27 | | X |

APPENDIX 2

List of 149 ant species collected on plants with and without extrafloral nectaries (EFNs) collected in 12 plots (250 x 25 m) in an terra-firme forest in the Brazilian Meridional Amazon located in the municipality of Cotriguaçu, north of Mato Grosso State, Brazil. The plants were collected between between December 2010 and January 2011.

| FAMLIY FORMICIDAE | Occurrence on plants | |
|---------------------------------|----------------------|--------------|
| | With EFNs | Without EFNs |
| SUBFAMILY DOLICHODERINAE | | |
| Tribe Dolichoderini | | |
| <i>Azteca</i> sp1 | X | X |
| <i>Azteca</i> sp2 | X | X |
| <i>Azteca</i> sp3 | X | X |
| <i>Azteca</i> sp4 | X | X |
| <i>Dolichoderus</i> sp1 | X | X |
| <i>Dolichoderus</i> sp2 | X | X |
| <i>Dolichoderus</i> sp3 | X | X |
| <i>Dolichoderus</i> sp4 | X | X |
| <i>Dolichoderus</i> sp5 | X | X |
| <i>Dolichoderus</i> sp6 | X | X |
| <i>Dolichoderus</i> sp7 | X | X |
| <i>Dolichoderus</i> sp8 | X | X |
| <i>Dolichoderus</i> sp9 | X | X |
| <i>Dolichoderus</i> sp10 | X | X |
| <i>Dolichoderus</i> sp11 | X | X |
| <i>Dolichoderus</i> sp12 | X | X |
| <i>Dolichoderus</i> sp13 | X | X |
| <i>Dolichoderus</i> sp14 | X | X |
| <i>Tapinoma</i> sp1 | X | X |
| <i>Tapinoma</i> sp2 | X | X |
| <i>Tapinoma</i> sp3 | X | X |
| SUBFAMILY ECITOTINAE | | |

Tribe Ecitonini

| | | |
|-------------------|--|---|
| <i>Eciton</i> sp1 | | X |
|-------------------|--|---|

SUBFAMILY ECTATOMMINAE**Tribe Ectatommini**

| | | |
|-------------------------------|---|---|
| <i>Ectatomma tuberculatum</i> | X | X |
|-------------------------------|---|---|

SUBFAMILY FORMICINAE**Tribe Camponotini**

| | | |
|------------------------------|---|---|
| <i>Camponotus latangulus</i> | X | X |
| <i>Camponotus</i> sp1 | X | X |
| <i>Camponotus</i> sp2 | X | X |
| <i>Camponotus</i> sp3 | X | X |
| <i>Camponotus</i> sp4 | X | X |
| <i>Camponotus</i> sp5 | X | X |
| <i>Camponotus</i> sp6 | X | X |
| <i>Camponotus</i> sp7 | X | X |
| <i>Camponotus</i> sp8 | X | X |
| <i>Camponotus</i> sp9 | X | X |
| <i>Camponotus</i> sp10 | X | X |
| <i>Camponotus</i> sp11 | X | X |
| <i>Camponotus</i> sp12 | X | X |
| <i>Camponotus</i> sp13 | X | X |
| <i>Camponotus</i> sp14 | X | X |
| <i>Camponotus</i> sp15 | X | X |
| <i>Camponotus</i> sp16 | X | X |
| <i>Camponotus</i> sp17 | X | X |
| <i>Camponotus</i> sp18 | X | X |
| <i>Camponotus</i> sp19 | X | X |
| <i>Camponotus</i> sp20 | X | X |
| <i>Camponotus</i> sp21 | X | X |
| <i>Camponotus</i> sp22 | X | X |
| <i>Camponotus</i> sp23 | X | X |
| <i>Camponotus</i> sp24 | X | X |

| | | |
|-------------------------------|---|---|
| <i>Camponotus</i> sp25 | X | X |
| <i>Camponotus</i> sp26 | X | X |
| Tribe Plagiolepidini | | |
| <i>Brachymyrmex</i> sp1 | X | X |
| <i>Brachymyrmex</i> sp2 | X | X |
| <i>Brachymyrmex</i> sp3 | X | X |
| <i>Brachymyrmex</i> sp4 | X | X |
| <i>Brachymyrmex</i> sp5 | X | X |
| <i>Brachymyrmex</i> sp6 | X | X |
| <i>Nylanderia</i> sp1 | X | |
| <i>Nylanderia</i> sp2 | X | X |
| <i>Nylanderia</i> sp3 | X | X |
| <i>Nylanderia</i> sp4 | X | X |
| <i>Nylanderia</i> sp5 | X | |
| <i>Nylanderia</i> sp6 | X | |
| SUBFAMILY MYRMICINAE | | |
| Tribe Attini | | |
| <i>Sericomyrmex</i> sp1 | | X |
| <i>Trachymyrmex</i> sp1 | X | |
| Tribe Blepharidattini | | |
| <i>Wasmannia auropunctata</i> | X | X |
| Tribe Cephalotini | | |
| <i>Cephalotes atratus</i> | X | X |
| <i>Cephalotes</i> sp1 | X | X |
| <i>Cephalotes</i> sp2 | X | X |
| <i>Cephalotes</i> sp3 | X | X |
| <i>Cephalotes</i> sp4 | X | X |
| <i>Cephalotes</i> sp5 | X | X |
| <i>Cephalotes</i> sp6 | X | X |
| <i>Cephalotes</i> sp7 | X | X |
| <i>Cephalotes</i> sp8 | X | X |
| Tribe Crematogastrini | | |
| <i>Crematogaster</i> sp1 | X | X |

| | | |
|---------------------------------|---|---|
| <i>Crematogaster</i> sp2 | X | X |
| <i>Crematogaster</i> sp3 | X | X |
| <i>Crematogaster</i> sp4 | X | X |
| <i>Crematogaster</i> sp5 | X | X |
| <i>Crematogaster</i> sp6 | X | X |
| <i>Crematogaster</i> sp7 | X | X |
| <i>Crematogaster</i> sp8 | X | X |
| <i>Crematogaster</i> sp9 | X | X |
| <i>Crematogaster</i> sp10 | X | X |
| <i>Crematogaster</i> sp11 | X | X |
| <i>Crematogaster</i> sp12 | X | X |
| Tribe Formicoxenini | | |
| <i>Nesomyrmex</i> sp1 | X | |
| <i>Nesomyrmex</i> sp2 | X | X |
| <i>Nesomyrmex</i> sp3 | X | |
| Tribe Ochetomyrmecini | | |
| <i>Ochetomyrmex neopolitus</i> | X | X |
| <i>Ochetomyrmex semipolitus</i> | X | X |
| Tribe Pheidolini | | |
| <i>Pheidole</i> sp1 | X | X |
| <i>Pheidole</i> sp2 | X | X |
| <i>Pheidole</i> sp3 | X | X |
| <i>Pheidole</i> sp4 | X | X |
| <i>Pheidole</i> sp5 | X | X |
| <i>Pheidole</i> sp6 | X | X |
| <i>Pheidole</i> sp7 | X | X |
| <i>Pheidole</i> sp8 | X | X |
| <i>Pheidole</i> sp9 | X | X |
| <i>Pheidole</i> sp10 | X | X |
| <i>Pheidole</i> sp11 | X | X |
| <i>Pheidole</i> sp12 | X | X |
| <i>Pheidole</i> sp13 | X | X |
| <i>Pheidole</i> sp14 | X | X |
| <i>Pheidole</i> sp15 | X | X |

| | | |
|----------------------|---|---|
| <i>Pheidole</i> sp16 | X | X |
| <i>Pheidole</i> sp17 | X | X |
| <i>Pheidole</i> sp18 | X | X |
| <i>Pheidole</i> sp19 | X | X |
| <i>Pheidole</i> sp20 | X | X |
| <i>Pheidole</i> sp21 | X | X |
| <i>Pheidole</i> sp22 | X | X |
| <i>Pheidole</i> sp23 | X | X |
| <i>Pheidole</i> sp24 | X | X |

Tribe Solenopsidini

| | | |
|-------------------------|---|---|
| <i>Megalomyrmex</i> sp1 | X | |
| <i>Solenopsis</i> sp1 | X | X |
| <i>Solenopsis</i> sp2 | X | X |
| <i>Solenopsis</i> sp3 | X | X |
| <i>Solenopsis</i> sp4 | X | X |
| <i>Solenopsis</i> sp5 | X | X |
| <i>Solenopsis</i> sp6 | X | X |
| <i>Solenopsis</i> sp7 | X | X |
| <i>Solenopsis</i> sp8 | X | X |
| <i>Solenopsis</i> sp9 | X | X |

SUBFAMILY PARAPONERINAE

Tribe Paraponerini

| | | |
|---------------------------|---|---|
| <i>Paraponera clavata</i> | X | X |
|---------------------------|---|---|

SUBFAMILY PONERINAE

Tribe Ponerini

| | | |
|-------------------------|---|---|
| <i>Odontomachus</i> sp1 | X | X |
| <i>Pachycondila</i> sp1 | X | X |
| <i>Pachycondila</i> sp2 | X | X |
| <i>Pachycondila</i> sp3 | X | X |
| <i>Pachycondila</i> sp4 | X | X |
| <i>Pachycondila</i> sp5 | X | X |
| <i>Pachycondila</i> sp6 | X | X |

| | | |
|--------------------------|---|---|
| <i>Pachycondila</i> sp7 | X | X |
| <i>Pachycondila</i> sp8 | X | X |
| <i>Pachycondila</i> sp9 | X | X |
| <i>Pachycondila</i> sp10 | X | X |
| <i>Pachycondila</i> sp11 | X | X |

SUBFAMILY PSEUDOMYRMICINAE

Tribe Pseudomyrmecini

| | | |
|--------------------------|---|---|
| <i>Pseudomyrmex</i> sp1 | X | X |
| <i>Pseudomyrmex</i> sp2 | X | X |
| <i>Pseudomyrmex</i> sp3 | X | X |
| <i>Pseudomyrmex</i> sp4 | X | X |
| <i>Pseudomyrmex</i> sp5 | X | X |
| <i>Pseudomyrmex</i> sp6 | X | X |
| <i>Pseudomyrmex</i> sp7 | X | X |
| <i>Pseudomyrmex</i> sp8 | X | X |
| <i>Pseudomyrmex</i> sp9 | X | X |
| <i>Pseudomyrmex</i> sp10 | X | X |
| <i>Pseudomyrmex</i> sp11 | X | X |

APPENDIX 3

Instructions for Authors – Ecology

Articles. While a Report is a concise scientific statement on a single simple topic, an Article tells a more complicated story with distinct components. The greater length of Articles relative to Reports must be justified by their greater complexity. We are asking authors to submit shorter, better-organized pieces that make use of Ecological Archives for digital publication of appendices and supplements. The target length for Articles is 20-30 manuscript pages (double-spaced, 12-point font, including everything from Title Page through the last figure). Longer Articles (those between 30 and 50 manuscript pages) should be accompanied by a detailed justification for the length in the cover letter at the time of submission. The abstract can have a maximum of 350 words. Manuscripts longer than 50 pages may be considered for Ecological Monographs, at the editor's discretion.

Requirements for submission

Original submission. Provide information describing the extent to which data or text in the manuscript have been used in other works that are published, in press, submitted, or soon to be submitted elsewhere.

Resubmission policy. If the manuscript (or a previous version of the manuscript) has been previously submitted to the same or another ESA journal, provide the previous manuscript number; explain how the current version differs from the previously submitted version and why it should be considered now for this journal. There are no guarantees it will be reviewed by the newly targeted journal.

ESA Code of Ethics. Authors must adhere to the ESA Code of Ethics.

Data Policy. The editors and publisher expect authors to make the data underlying published articles available. Authors must disclose software and statistical procedures used in the manuscript and provide any novel computer code used for models, simulations, or statistical analyses.

English. Authors whose native language is not English are encouraged to enlist the aid of a native English-speaking colleague to go over the manuscript for correct usage and clarity prior to submission. You may also search the ESA Author Help Directory to find a volunteer. No guarantees are associated with the use of this volunteer database.

Agreement to pay page charges. Authors must agree to pay page charges, or must obtain an ESA page-charge grant. Page charges will only be incurred for manuscripts accepted for publication.

Formatting your manuscript: Minimum formatting requirements

Consult recent issues for examples of journal style. For purposes of review, submitted manuscripts need not adhere to journal style in every detail; however, preparation of final revisions of manuscripts accepted for publication will be easier if ESA style is followed from the outset. But be sure to abide by the following minimum formatting requirements for submitted manuscripts:

- * The entire manuscript must be double-spaced (text, quotations, figure legends, literature cited) at three lines per inch (12 lines/10 cm) with a 12-point font, Times New Roman. Choose the "double-spacing" option for line spacing. Leave a 1 inch (2.4-cm) margin on all sides of each page. Page size should be Letter 8 ½" by 11". Do not justify the right margin.

- * Assemble the parts of the manuscript in this order: title page, abstract, key words, text, acknowledgments, literature cited, tables (one table per page), figure legends (on separate page preceding the first figure), figures (one figure per page; label each figure, i.e., Figure 1, Figure 2, etc.). Appendices for Ecological Archives should be in a separate file.

- * Number all pages (including tables, and figures), starting with the title page.

- * All pages of text should have line numbers as well.

Allowable file formats:

Manuscript files in Word (.doc or .docx), WordPerfect (.wpd), Rich-text format (.rtf) or LaTeX (.TEX) format. (See ESA-LaTeX for some tips on getting your TEX document to conform to ESA style.)

Tables in doc, xls, tds, or csv format (or Tables may be included in the manuscript file)

Figures/Images in doc, jpeg, tif, gif, eps, ps, or ppt format (or Figures may be included in the manuscript file)

Appendices for Ecological Archives in doc or html format. Video appendices in mpeg format.

Supplements for Ecological Archives can include, but are not limited to, original and derived data sets, source code for simulation models, and details of and software for unusual statistical analyses.

Appendices and Supplements for Ecological Archives should be in files separate from the article (and not merged with the article file). Zipped files can be submitted for appendices and supplements if necessary. In that case, choose Zipped File as the file type, so that the system does not attempt a pdf conversion.

Tables and figures may be in a separate file or in one file together with the manuscript text. If figures are in a separate file, please provide a separate file with all the figure legends (or include it in the manuscript file). It is not necessary to provide a figure caption on the screen when uploading your figures. (Please be aware that a lower resolution figure may look fine on a computer screen, but that does not mean it will look good if a reviewer or editor prints it out.)

APPENDIX 4

Instructions for Authors – *Oikos*

Author Guidelines

General editorial practice

Submit your manuscript electronically using a single PDF file at www.oikos.ekol.lu.se/oikosgateway

You will receive a receipt with a tracking code. Please refer to this code in all correspondence with the Editorial Office.

We as well as reviewers have problems in handling LaTeX files, please avoid this format.

Authors submitting a manuscript do so on the understanding that the work has not been published before, is not being considered for publication elsewhere **and has been read and approved by all authors**. The submission of the manuscript by the authors means that the authors automatically agree to assign exclusive copyright to the journal if and when the manuscript is accepted for publication.

This work shall not be published elsewhere in any language without the written consent of the journal. The articles published in this journal are protected by copyright, which covers translation rights and the exclusive right to reproduce and distribute all of the articles printed in the journal. No material published in the journal may be stored on microfilm or videocassettes or in electronic databases and the like or reproduced photographically without the prior written permission of the journal.

Manuscripts are submitted to reviewers for evaluation of their significance and soundness. Authors will generally be notified of acceptance, rejection, or need for revision within three months. Decisions of the editor are final.

Manuscripts are edited to improve communication between author and reader.

Authors will receive electronic proofs via e-mail, together with the Exclusive License Form (ELF)). Correct **only** printer's mistakes. You will be charged for excessive corrections beyond such mistakes.

Exclusive Licence Form. Authors will be required to sign the Exclusive Licence Form (ELF) for all papers accepted for publication. Signature of the ELF is a condition of publication and papers will not be passed to the publisher for production unless a signed form has been received. Please note that signature of the Exclusive Licence Form does not affect ownership of copyright in the material. (Government employees need to complete the Author Warranty sections, although copyright in such cases does not need to be assigned). After publication authors will retain the right to publish their paper in various media/circumstances (please see the form for further details). To assist authors an ELF can be supplied by the Editorial Office. Alternatively, authors may like to download a copy of the form.

No offprints will be supplied. Instead corresponding authors will receive a locked PDF file to the use at their discretion.

There are no page charges. We do, however, charge for colour printing.

Authorship. Statement on authorship. Papers should conform to recommendations for authorship provided by the International Committee of Medical Journal Editors (the Vancouver Group; see <http://www.icmje.org>). That is, authorship of a paper carries with it responsibility as well as credit. All those whose names appear as authors should have played a significant role in designing or carrying out the research, writing the manuscript, or providing extensive guidance to the execution of the project. They should be able to present and defend the work in a public forum. Honorary authorship is to be avoided. All authors must be in agreement on both the submission and full content of any article carrying their name. Any violation of these conditions represents academic misconduct and will be dealt with accordingly.

Manuscripts

Manuscripts should be submitted as one single pdf-file with tables and figures included. Upon acceptance the manuscript should be provided in Microsoft Word, Rich Text Format

or Post Script format with high resolution figures included. Oikos does not yet accept manuscripts in Microsoft Word 2007 format.

Language. Manuscripts should be in English. Linguistic usage should be correct. Avoid the use of the passive voice. Avoid extensive reviews in the Introduction and Discussion. Cite only essential sources of a theory or opinion.

Title. The title should be brief and contain words useful for indexing and information retrieval.

Text. The first page should contain only the title and the author's name, address, fax and email-address. Page two contains the abstract, in which the main results of the work should be summarized. The abstract should not contain more than 300 words. Begin the introduction on page three. Avoid right margin justification and hyphenation. Double-check the contents of your manuscript before submitting. Only printer's mistakes in proofs will be changed free of charge. Oikos do not print symbols or formulas in italics.

Illustrations. Tables and legends of illustrations should be written double-spaced on separate sheets. Do not incorporate the legend in the figure itself. Tables and illustrations should be comprehensible without reference to the text. Do not use italic lettering.

Figures should be planned to appear with a maximum final width of 8 cm (single-column), 12.5 cm (1.5 column) or 16.6 cm (double-column). The font used in figures should be either Helvetica or Arial. Letters, numbers and symbols must appear clearly but not oversized. A suitable final size for lettering is 1-2 mm at reproduction size. One uniform size throughout is generally recommended. Avoid complicated symbols or patterns. Use open and closed circles, squares and triangles; open, striped and closed bars in histograms. Each figure should be boxed in and scale marks (turning inwards) provided. Lines should be clear, but not thick and heavy. Plan your illustrations for the smallest size possible (one column). Be sure that the lettering is clear and readable, even if the figure is de-sized.

Colour plates may be included at the author's expense, €300 per paper.

Units. Use SI units as far as possible.

Nomenclature. Binomial Latin names should be used in accordance with International Rules of Nomenclature.

References. In the list of references (double-spaced), the following usage should be conformed to:

Journal

Haila, Y. and Järvinen, O. 1983. Land bird communities on a Finnish island: species impoverishment and abundance patterns. - *Oikos* 41: 255-273.

If more than two authors: Lindsay, A. et al. 2000. Are plant populations seed-limited? A review of seed sowing experiments. – *Oikos* 88: 225–238.

Book

Mayr, E. 1963. *Animal species and evolution*. - Harvard Univ. Press.

Chapter

Goodall, D. W. 1972. Building and testing ecosystem models. - In: Jeffers, J. N. R. (ed.), *Mathematical models in ecology*. Blackwell, pp. 173-194.

In the text references are given: Mayr (1963) or, at the end of a sentence, (Mayr 1963).

Titles of journals should be abbreviated following *Biological Abstracts*. If in doubt, give the title in full. Do not refer to unpublished material.

The list of references should be arranged alphabetically on authors' names and chronologically per author. If the author's name is also mentioned with co-authors the following order should be used: publications of the single author, arranged chronologically - publications of the same author with one co-author, arranged chronologically - publications of the author with more than one co-author, arranged chronologically. Publications by the same author(s) in the same year should be listed as 2004a, 2004b, etc. Reference lists not conforming to this format will be returned for revision.

Excessive use of references causes unnecessary long articles. To avoid excessive use of references, use only the most relevant. As a rule, avoid using more than 50 references in a regular research paper.

Acknowledgements. Keep them short.

Appendices: Supplementary material may be posted as electronic appendices on the journal's appendix site.

Read important instructions on how we handle supplementary material [here](#).

Authors are recommended to follow the guidelines set out in: O'Connor, M. 1991. Writing successfully in science. - Harper Collins Academic, London, **and to examine the latest issues of Oikos**. Manuscripts not conforming to the requirements will be returned for revision.

Online Open

OnlineOpen is available to authors of primary research articles who wish to make their article available to non-subscribers on publication, or whose funding agency requires grantees to archive the final version of their article. With OnlineOpen, the author, the author's funding agency, or the author's institution pays a fee to ensure that the article is made available to non-subscribers upon publication via Wiley Online Library, as well as deposited in the funding agency's preferred archive. For the full list of terms and conditions, see http://wileyonlinelibrary.com/onlineopen#OnlineOpen_Terms

Any authors wishing to send their paper OnlineOpen will be required to complete the payment form available from our website at: <https://onlinelibrary.wiley.com/onlineOpenOrder>

Prior to acceptance there is no requirement to inform an Editorial Office that you intend to publish your paper OnlineOpen if you do not wish to. All OnlineOpen articles are treated in the same way as any other article. They go through the journal's standard peer-review process and will be accepted or rejected based on their own merit.