

## Stronger together: Combining automated classifiers with manual post-validation optimizes the workload vs reliability trade-off of species identification in bat acoustic surveys

Adrià López-Baucells<sup>a,b,c,e,\*</sup>, Laura Torrent<sup>b</sup>, Ricardo Rocha<sup>a,c,d</sup>, Paulo E.D. Bobrowiec<sup>c</sup>, Jorge M. Palmeirim<sup>a</sup>, Christoph F.J. Meyer<sup>a,c,d</sup>

<sup>a</sup> University of Lisbon, Centre for Ecology, Evolution and Environmental Changes (cE3c), Lisbon 1749-016, Portugal

<sup>b</sup> Granollers Museum of Natural Sciences, Granollers 08402, Catalonia, Spain

<sup>c</sup> Biological Dynamics of Forest Fragments Project (BDFFP), National Institute for Amazonian Research, Manaus 69011-970, Brazil

<sup>d</sup> University of Cambridge, Department of Zoology, Conservation Science Group, Cambridge CB23EJ, United Kingdom

<sup>e</sup> University of Salford, School of Environment and Life Sciences, Salford M54WT, United Kingdom

### ARTICLE INFO

#### Keywords:

Amazon  
Bioacoustics  
Chiroptera  
Echolocation  
Machine-learning algorithms

### ABSTRACT

Owing to major technological advances, bioacoustics has become a burgeoning field in ecological research worldwide. Autonomous passive acoustic recorders are becoming widely used to monitor aerial insectivorous bats, and automatic classifiers have emerged to aid researchers in the daunting task of analysing the resulting massive acoustic datasets. However, the scarcity of comprehensive reference call libraries still hampers their wider application in highly diverse tropical assemblages. Capitalizing on a unique acoustic dataset of > 650,000 bat call sequences collected over a 3-year period in the Brazilian Amazon, the aims of this study were (a) to assess how pre-identified recordings of free-flying and hand-released bats could be used to train an automatic classification algorithm (random forest), and (b) to optimize acoustic analysis protocols by combining automatic classification with visual post-validation, whereby we evaluated the proportion of sound files to be post-validated for different thresholds of classification accuracy. Classifiers were trained at species or sonotype (group of species with similar calls) level. Random forest models confirmed the reliability of using calls of both free-flying and hand-released bats to train custom-built automatic classifiers. To achieve a general classification accuracy of ~85%, random forest had to be trained with at least 500 pulses per species/sonotype. For seven out of 20 sonotypes, the most abundant in our dataset, we obtained high classification accuracy (> 90%). Adopting a desired accuracy probability threshold of 95% for the random forest classifier, we found that the percentage of sound files required for manual post-validation could be reduced by up to 75%, a significant saving in terms of workload. Combining automatic classification with manual ID through fully customizable classifiers implemented in open-source software as demonstrated here shows great potential to help overcome the acknowledged risks and biases associated with the sole reliance on automatic classification.

### 1. Introduction

Bioacoustics is a rapidly expanding field and of increasing importance for informing conservation projects. This is largely due to recent technological advances and the rising number of long-term monitoring programs which are being established for a number of taxa (Dickinson et al., 2010; Kerstenbaum et al., 2014), including birds (Gregory et al., 2005), reptiles (Sewell et al., 2012), arthropods (Penone et al., 2013) and bats (Barlow et al., 2015). Interest in bat monitoring has increased over the last decades since bats have been acknowledged

to provide important ecosystem services such as pest control (Boyles et al., 2013; Puig-Montserrat et al., 2015) and have been identified as good bioindicators of ecosystem health (Cunton and Bernard, 2012; Jones et al., 2009).

Autonomous ultrasound detectors have proven essential for optimizing surveys of aerial insectivorous bats worldwide (Law et al., 2015; Murray et al., 1999). In the Neotropics, in contrast to phyllostomid bats, aerial insectivores are rarely captured in mist-nets (MacSwiney et al., 2008) and therefore, although they represent a high proportion of Neotropical bat diversity (Jung and Kalko, 2011), the ecology of many

\* Corresponding author at: University of Lisbon, Centre for Ecology, Evolution and Environmental Changes (cE3c), Lisbon 1749-016, Portugal.

E-mail address: [adria.baucells@gmail.com](mailto:adria.baucells@gmail.com) (A. López-Baucells).

species remains elusive and their echolocation calls poorly described (e.g. López-Baucells et al., 2014; López-Baucells et al., 2017a). In fact, despite enormous recent advances in recording technology and equipment, comprehensive regional bat reference call libraries are currently lacking for much of the tropics (Madhukumar Menon et al., 2018; Walters et al., 2013).

Reference call libraries containing echolocation calls from a wide range of locations and habitats are crucial to reliably identify bat species acoustically. Although many species have distinctive echolocation calls, those of others can be very ambiguous due to producing very similar calls with overlapping characteristics (Russo and Voigt, 2016). Moreover, weather conditions (e.g. Lawrence and Simmons, 1982), geographical location (e.g. López-Baucells et al., 2017b), sex (e.g. Puechmaille et al., 2014), body condition (e.g. Puechmaille et al., 2014), age (e.g. Jones and Kokurewicz, 1994), reproductive status (e.g. Jones and Ransome, 1993) or habitat structure (e.g. Pedro and Simonetti, 2014) are all factors that contribute to substantial variation in call structure within species.

Different algorithms such as discriminant function analysis and random forest have already been used to automatically classify bat pulses (Armitage and Ober, 2010; MacSwiney et al., 2008; Russo and Jones, 2002; Walters et al., 2012; Zamora-Gutiérrez et al., 2016). However, substantial controversy still exists around the trade-off between the use of automatic classifiers versus manual species identification (Kershenbaum et al., 2014; Russo and Voigt, 2016). While the former allows for the rapid analysis of a large number of recordings using an objective and repeatable protocol, manual identification provides more accurate, yet highly subjective and non-reproducible results, apart from being considerably more time-consuming (Kershenbaum et al., 2014). Unfortunately, even though considered a vital analysis step when relying on automated classifiers, posterior visual cross-checking by an expert is all too often neglected (Russo and Voigt, 2016). Moreover, no study so far has quantified the potential time savings from using automatic classifiers as a function of the classification accuracy threshold adopted. Automatic classifiers have been criticised because of the inability to distinguish among species with similar calls, and because their algorithms are typically trained with calls from hand-released bats (Russo and Voigt, 2016). It has been suggested that the use of hand-release calls can compromise the reliability of species identifications since these calls might be strongly affected by handling-related stress of the animal (Szewczak, 2000).

Given that automatic classifiers are now widely available, there is a substantial risk that beginners solely rely on automated species identification without proper manual post-validation, which can result in incorrect identifications and thus wrong management decisions and negative conservation outcomes (Russo and Voigt, 2016). Automatic classifiers were first applied to bat species identification in temperate areas as a direct consequence of the massive acoustic datasets that are now typically accumulated using passive bat recorders (Russo and Voigt, 2016). However, the scarcity of suitable reference call libraries and the controversy around automatic vs. manual classification still hamper their wider application, especially in mega-diverse tropical regions.

The present study is the first to attempt to test the suitability of combining automatic classifiers trained with pre-identified recordings of free-flying bats obtained in the study area (which are much easier to obtain than reference calls from hand-released bats) with posterior manual validation (Fig. 1). This approach addresses the aforementioned issues of geographic variability, only classifies calls to the taxonomic level that the researcher can visually confirm with certainty and gives the user full control and flexibility concerning implementation of the algorithm. Capitalizing on a unique acoustic dataset collected over a 3-year period in the Central Amazon, here we use random forest, a machine learning algorithm that has performed well in previous bat acoustic studies (Bas et al., 2017; Zamora-Gutiérrez et al., 2016), to automatically classify aerial insectivorous bats. We evaluated the

discriminative ability of the classifier by training it with a) previously identified calls from free-flying bats and those of hand-released bats; and b) datasets of different sizes of reference echolocation calls. To effectively combine the advantages of an automatic classifier with those of manual identification requires establishing a “correct classification probability” threshold below which a recording will need to be visually post-validated. Thus, to evaluate how acoustic studies could be optimized in terms of time commitment for the analyses, we also calculated, for different thresholds, the percentage of sound files from the full dataset that would need to be visually post-validated.

## 2. Material and methods

### 2.1. Study site

The study was conducted at the Biological Dynamics of Forest Fragments Project (BDFFP), a large-scale fragmentation experiment located ~80 km north of Manaus (Brazil) in the Central Amazon (2°20'S, 60°6'W), aimed at assessing the impacts of fragmentation on tropical forest communities (Laurance et al., 2011). Beginning in 1979, the BDFFP established 11 experimental forest fragments, which at the time of isolation were separated from continuous forest by distances of 80–650 m. Nowadays the fragments are surrounded by a matrix of secondary forest at varying successional stages (Laurance et al., 2017). The area is currently composed of a mosaic of unflooded lowland forest (80–160 m a.s.l.), pastures and secondary regrowth forest. Primary forest reaches 30–37 m in mean canopy height, with isolated trees up to 55 m tall (Laurance et al., 2011). Annual rainfall varies between 1900 and 3500 mm per year, with a rainy season between November and June and a dry season from July to November (Ferreira et al., 2017), while mean annual temperature usually oscillates between 26 and 30 °C (de Oliveira and Mori, 1999).


### 2.2. Mist-netting and hand-release recordings

Intensive bat sampling was carried out in the context of a larger project assessing fragmentation effects on bats in the BDFFP landscape over a period of four years (2011–2014), using both ground- and canopy-level mist-netting. Sampling covered various types of rainforest habitats including continuous primary forest, forest fragments and secondary regrowth (Farneda et al., 2015; Rocha et al., 2017a; Rocha et al., 2017b). Sporadic sampling was also done over temporary lakes, small ponds and streams, as well as campsites, roads, and pastures (Torrent et al., 2018). Mist-netting was usually conducted from 18:00 to 00:00, except for some lakes where high capture rates sometimes required closing the nets earlier. Captured bats were identified using different keys (Gardner, 2007; Lim and Engstrom, 2001).

Echolocation call recordings of captured aerial insectivorous bats were made with a Pettersson D1000 bat detector (Pettersson Elektronik, Sweden), using 384 kHz sampling frequency in full spectrum (16-bit resolution) and no triggers or filters. Release calls were obtained after hand release of bats in either clearings or open areas within the forest ( $N = 722$  individuals). The detector was placed 5–10 m from the point of release (depending on the species) and once the individual was in flight, the microphone was pointed towards it to record as many search pulses as possible. For analysis, all pulses recorded immediately after release were discarded, as were overloaded calls, those too faint (for which it was impossible to distinguish the shape from the background noise), social or stress calls, calls emitted in passive hunting mode and feeding buzzes.

### 2.3. Acoustic monitoring dataset

A total of 50 sites across the BDFFP landscape were acoustically surveyed 2012–2014, including the same sites used for mist-netting as described in Rocha et al. (2017a, 2017b). These comprised different-



**Fig. 1.** Diagram illustrating the classification process from automatic classification to posterior visual validation. Squares represent the datasets and selections of recordings; arrows and red text represent the analytical processes. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

sized forest fragments ( $N = 8$ ), continuous forest ( $N = 9$ ), forest edges ( $N = 11$ ), secondary forest ( $N = 11$ ) and forest clearings ( $N = 11$ ). At each recording point, an automatic SM2Bat detector with an omnidirectional ultrasonic SMX-US microphone (Wildlife Acoustics, Inc., USA) was placed ca. 1.5 m above the ground. Acoustic surveys covered both dry and wet seasons and were conducted twice per season. Detectors were set to automatically record bats from 18:00 to 06:00 in real time with a full spectrum resolution of 16 bit, a high-pass filter set at fs/32 (12 kHz), an adaptive trigger level relative to noise floor of 18 SNR, and for periods of five consecutive nights per site. All recordings were split into five-second long sequences. Within such a five-second sound file, a bat pass was defined as a sequence with a minimum of two recognizable echolocation pulses per species (Appel et al., 2017; Millon et al., 2015; Torrent et al., 2018). This unit was used as a measure of activity levels. A total of 1,088,940 sound files were acquired during the study period in which ~650,000 bat passes were identified.

#### 2.4. Echolocation call analysis

Kaleidoscope v.4.0.4 software (Wildlife Acoustics Inc., USA) was used to visualize and manually classify all bat passes from the acoustic monitoring dataset. Call sequences were manually identified to species/sonotype level as in previous studies (Silva and Bernard, 2017; Torrent et al., 2018). For the purpose of this study, a sonotype was defined as a category that grouped species with similar calls when it was not possible to clearly assign a call to a particular species (Table S1). Call identification was based on a series of acoustic features and standard measurements - call shape (CS), frequency of maximum energy (FME), start (SF), end (EF), maximum (MaxF) and minimum (MinF) frequency and duration (Dur) - and followed the echolocation key in López-Baucells et al. (2016). Moreover, recordings were also compared with a local reference call library compiled for the same study area over the course of the whole 3-year sampling period. Call sequences or pulses that were too faint for reliable identification (< 10 dB difference in power between background noise and FME of the echolocation pulses) were discarded from the analysis.

In addition to this manual identification, the same recordings were also subjected to an automatic identification process whereby pulse measurements were automatically extracted (~4,788,000 pulses) using SCAN'R (Snapshot Characterization and Analysis Routine) v1.7.4. (Binary Acoustic Technology, USA). Settings were adjusted as specified in Table S2 to minimize the confusion between noise and bat calls. The following measurements were extracted for all pulses: Duration (Dur,

ms), Maximum frequency (Fmax, kHz), Minimum frequency (Fmin, kHz), total bandwidth (BW, ms), Frequency at strongest sound pressure level (Fdom, kHz; equivalent to FME or *Frequency of maximum energy*), percentage in duration of Fdom (Ldom, %), High end of characteristic frequency (HiFc, kHz; equivalent of the knee frequency), Low end of characteristic frequency (LowFc, kHz), global slope of the call (Slope, kHz/ms), curvature (Curv) (SCAN'R, 2009). After extraction, a Principal Component Analysis (PCA) was performed, separately for each bat family, in order to visualize how different species/sonotypes clustered based on the similarity of their acoustic parameters.

#### 2.5. Supervised machine learning

Supervised classification based on a machine learning algorithm (random forest, RF) was conducted using the R package “caret” (Classification and Regression Training) (Kuhn, 2008). Random forest has performed well in several bat studies and is currently the preferred machine learning algorithm for the classification of bat echolocation calls (e.g. Bas et al., 2017; Zamora-Gutiérrez et al., 2016). Random forest models are built by comparing and averaging decision tree classifiers that are designed by bootstrapping random samples of the training dataset (Breiman, 2001). Among its advantages, random forest is not affected by heteroscedasticity, is not strongly affected by outliers or low-informative variables, and is relatively easy to use computationally (Olden et al., 2008), which makes it the method of choice for large acoustic datasets. In our case we selected three separate 10-fold cross-validations to tune the training model, with a final value of *mtry* of 2 (chosen for their highest accuracy) (Breiman, 2001).

##### 2.5.1. Data preparation

All pulse measurements were centred and scaled (Kuhn, 2008; Mukherjee and Manna, 2006) to make them comparable. The global dataset (~4,178,000 pulses) was split into different training and testing subsets. Training datasets were composed of 50, 100, 500, 1000 and 2000 reference pulses per species/sonotype, which were randomly selected from all recordings (except for *Rhynchonycteris naso* and *Furipterus horrens*, for which we only had data from 12 and 1000 pulses respectively).

##### 2.5.2. Data classification

Using the 1000-pulse training dataset, we evaluated classification accuracy and predictive power of the RF algorithm. Evaluation of performance of the training algorithm on the testing datasets was based

on the performance metrics accuracy and kappa. Kappa measures inter-rater agreement for qualitative items (usually considered to be more robust than other measures as it also takes into account the agreement occurring by chance) (Viera and Garrett, 2005). The same metrics were then additionally assessed for the different-sized training datasets, ranging from 50 to 2000 pulses/sonotype. Variable (feature) importance scores were also obtained using the R package caret (Kuhn, 2008). The contribution of each variable is measured as follows: For each tree, the prediction accuracy is recorded removing each predictor variable. The average of the differences between all accuracies is normalized by the standard error.

Classification success for each species/sonotype was evaluated using 1) a RF model trained with the 2000-pulse dataset based on calls of free-flying bats and 2) a RF model trained with the complete reference call library based on hand-release calls compiled during the whole 3-year study period. The latter unfortunately included < 2000 pulses for many species (Table S3) due to the inherent difficulty to capture enough individuals from which to obtain release calls. Both training datasets were classified using the same species/sonotype labels in order to make both classifications comparable. Among the whole set of metrics commonly used to evaluate classifiers, we selected *sensitivity* and *positive predictive value* (PPV) as the most conservative for evaluating the performance of the acoustic classification task because they highlight the true positives in the classification process (Jennings et al., 2008) (Fig. S1). While sensitivity is the proportion of calls correctly identified as one species/sonotype out of the total number of calls, positive predictive value is the proportion of calls correctly identified as such. Other metrics such as specificity or negative predictive value highlight the certainty of true negatives, which is quite unreliable in multicategory classifications (Fig. S1).

The estimation of the percentage of recordings that would need to be manually checked depending on several classification accuracy thresholds was also based on the 2000-pulse training dataset. Classification accuracy thresholds considered in the analyses ranged from 60 to 95%, in 5% increments.

### 3. Results

#### 3.1. Acoustic discrimination at family level

A total of 27 aerial insectivorous bat species from six different families were captured and recorded during the study period, representing 20 different species/sonotypes (Table S1). PCAs based on acoustic features showed that, for mormoopids, automatic parameter extraction often resulted in measurement values coming from different harmonics (Fig. 2). *Pteronotus alitonus* and *P. rubiginosus* clearly separated as distinct clusters and, although less evident, *P. personatus* and *P. gymnonotus* were also quite distinctly separated. Similarly, species with modulated calls such as vespertilionid or furipterid bats were split in rather well-defined bands. In contrast, except for *Saccopteryx bilineata* and *S. leptura*, emballonurid and molossid bats showed less defined limits between groups.

#### 3.2. Minimum training dataset size and variable importance

We found that, in order to achieve a minimum general accuracy of ~85%, a training dataset of > 500 pulses per species/sonotype was required (Fig. 3A). Classifications undertaken with training datasets based on only 50 pulses showed large variation in accuracy, reaching values below 75%. Classification performance was consistent between accuracy and kappa metrics. “High end of characteristic frequency” (equivalent to the frequency of the knee) was the most important variable in the RF model, followed by “Maximum frequency” and “Dominant frequency” (equivalent to the frequency of maximum energy). However, except for “Length of the dominant frequency”,

“Duration”, “Bandwidth” and “Curvature”, all the variables showed quite similar importance values (Fig. 3B).

#### 3.3. Classifier performance at species/sonotype level

Algorithm performance varied substantially among species/sonotypes (Table S4). Seven had values above 90% for both sensitivity and PPV (*P. alitonus*, *P. rubiginosus*, Vespertilionidae 1, *Myotis nigricans*, *Centronycteris maximiliani*, *Myotis riparius* and *S. bilineata*), indicating not only that most of the recordings were correctly assigned, but also that few other recordings were confused with these species (Table 1, Table S4). On the other hand, for other species such as *Furipterus horrens*, Emballonuridae 1, *P. gymnonotus*, Molossidae 3, *Promops* spp. and *P. personatus* there were considerable differences between metrics. For these, we found a low number of false negatives but a large number of false positives (low PPV). Molossidae 1 and 2 were the sonotypes with poorest levels of correct identifications, and *R. naso* (for which we had a very limited number of recordings) was the only species for which the classifier completely failed. Comparing the RF models trained with calls from free-flying vs. hand-released bats, the former nearly always outperformed the latter (Table 1). *Pteronotus alitonus* and *P. rubiginosus* obtained a similar proportion of correct identifications in both HR and FF algorithms, and Molossidae 3 was the only sonotype for which higher sensitivity scores were obtained using calls from hand-released bats, although it also had lower PPV.

#### 3.4. Combining automatic classification with manual post-validation

The total number of files to be visually confirmed after automatic classification ranged from ~20%, when the desired accuracy threshold was set to 60%, to ~30% when it was set to 95% (Fig. 4). Following the same pattern found for species/sonotype-specific predictive performance (Table 1), for some categories the number of files to be post-validated did not differ substantially for the different accuracy thresholds mentioned previously, while for others, this percentage varied up to 30%. Only in few cases was there marked variation depending on the chosen accuracy threshold (from 40 to 90% for Emballonuridae 1 and *P. personatus*).

### 4. Discussion

Our analyses suggest an inexpensive and relatively user-friendly approach (Fig. 1) to automatically classify large amounts of bat echolocation data, followed by visual post-validation of a reduced proportion of the original acoustic dataset. This approach overcomes the acknowledged risks and biases associated with the exclusive reliance on current automatic classifiers (Russo and Voigt, 2016). Using recordings obtained under real field conditions from a 3-year-long study in the Central Amazon, we confirmed the reliability of using locally-recorded echolocation calls from free-flying bats to train a custom-built classifier that automatically identifies the calls of a large subset of the species/sonotypes in the local assemblage with high accuracy (> 90%) and leaves the rest to be manually classified. This automatic pre-classification reduces the total number of recordings to be visually inspected, therefore optimizing the classification process. This equates to considerable time savings, especially in the case of projects that accumulate massive acoustic data. However, due to the customizable nature of this approach, the advantages of using recordings from free-flying bats hinge on manually pre-identifying a decent amount of calls from free-flying bats using release calls as references, literature and echolocation keys. This obviously entails the risk of including misidentified calls as a source for training the algorithm, a problem we overcame by restricting the classification of the recordings to easily distinguishable species/sonotypes, therefore avoiding misidentifications.

Being non-intrusive, automated recording systems and soundscape studies have recently become very popular, and have considerably






Fig. 2. Principal component analyses (PCA) based on measurements of a series of acoustic parameters (see Methods) that were automatically extracted with SCAN'R, and manually classified to species/sonotype level following López-Baucells et al. (2016).

improved our knowledge about the natural history of elusive bat species, anthropogenic impacts and wildlife conservation in habitats where sampling by traditional methods such as mist-netting would be inefficient or unfeasible (Kubista and Bruckner, 2017). However, in developing countries where funding is particularly limited, the widespread use of bioacoustics is still severely hampered by its elevated costs and cost-effective alternatives need to be found quickly. This has inspired a new trend towards developing user-friendly detectors and automatic classifiers which are fully customizable at reduced cost (Hill et al., 2018; Whytock et al., 2017). Reliance on self-built classification algorithms could greatly contribute to studies in regions for which no automatic classifier is available as part of commercially available software packages.

#### 4.1. Minimum training dataset size and variable importance


We identified the minimum number of pulses that should be used in the training dataset in order to achieve general accuracy levels between 75 and 95%. Our results show that training the algorithm with 500 pulses per species/sonotype results in average classifier performance > 85%. However, this reference value should be interpreted carefully as it depends on whether the species that are most frequently detected in a region are also those whose echolocation call characteristics are more clearly distinct and thus the species more easily

identifiable or not. In our case, the most common species (*P. rubiginosus*, *Myotis nigricans* and *M. riparius*), all easy to identify, might be positively biasing general accuracy, thus masking lower accuracies for the remaining categories (Biscardi et al., 2004).

The variables that contribute most to separating species may not be the same in all assemblages. For example, Monadjem et al. (2017) found that call duration represented one of the most relevant parameters to distinguish between species, while in our study, we obtained higher importance weights for several other variables. This probably reflects the diversity of pulse shapes and structures found in Neotropical aerial insectivorous bats. By including different variables in the algorithm, one can probably achieve better classification performance in such highly diverse areas (Walters et al., 2013).

#### 4.2. Classifier performance at species/sonotype level

Random forest performed very well with our dataset, confirming its great potential for analysing bat acoustic datasets. Among the available machine learning algorithms, random forest has already been successfully used in automatic species classification for bats (Armitage and Ober, 2010; Zamora-Gutiérrez et al., 2016) and other taxa such as birds (Briggs et al., 2009) and dolphins (Barkley et al., 2011). We obtained similar mean accuracies to those found in previous studies, although results varied among species and families (e.g. Britzke et al., 2011;



**Fig. 3.** A) Classifier performance, evaluated as general accuracy and kappa, for a random forest model built with different-sized training datasets (50 to 2000 pulses/sonotype). The x-axis has been scaled to allow better visualization. Dots are medians, boxes 25% and 75% quartiles and whiskers denote the range. B) Importance of each variable in the random forest model trained with 2000 reference pulses per species/sonotype.

MacSwiney et al., 2008; Pio et al., 2010; Zamora-Gutiérrez et al., 2016). For Neotropical bats, large variability in predictive power is found for Vespertilionidae and Molossidae, while Emballonuridae and Mormoopidae are usually more accurately identified (Zamora-Gutiérrez et al., 2016). Previous studies have evaluated the performance of automatic algorithms for classifying bat calls at species, genus, family or guild level (Vassilios et al., 2017; Zamora-Gutiérrez et al., 2016). However, it is now widely accepted in the scientific community that automatic classification must be used cautiously (Monadjem et al., 2017; Russo and Jones, 2002; Russo and Voigt, 2016). In this study, we aimed to optimize the classifiers not at species level but using sonotypes. Although classifying all calls to species level would be ideal, using sonotypes may be sufficient in most cases, obviously depending on a project's specific aims (Armitage and Ober, 2010; Redgwell et al., 2009).

For seven out of 20 species/sonotypes we obtained very high values (> 90%) for both sensitivity and PPV, proving that our random forest algorithm could be used with great confidence to detect and automatically classify them in our recordings. Very few false positives and false negatives were found, indicating that our classifier neither gets them wrong, nor ignores them when they are present (see Table 1). These species are also the most predominant in our dataset, which turns our classifier into a great tool due to its potential to greatly reduce the number of files to be manually analysed (Andreassen et al., 2014). One of the main reasons to explain the classification failure of some categories is the limited capacity of SCAN'R to detect and characterize pulses of different lengths (our SCAN'R pulse detection settings were

more suitable for long pulses). This will certainly improve soon with new technological advances, or alternatively, could be better implemented through R sound packages. Previous studies have exclusively used accuracy as a means of evaluating algorithm performance and predictive capacity (i.e. Wordley et al., 2014; Zamora-Gutiérrez et al., 2016). However, other more conservative metrics such as positive predictive value and sensitivity are often neglected. We encourage developers of algorithms and researchers to better scrutinize classifier performance by focusing on these more reliable metrics.

#### 4.3. Classifier trained with calls from free-flying versus hand-released bats

We compared the performance of the random forest classifier trained with calls from free-flying versus hand-released bats, using only data collected during the 3-year-period of the project. Classifier performance was substantially better using recordings from free-flying bats, probably due to the low number of recordings from hand-released bats for most of the species. In this regard it is important to mention that the effort required to compile complete reference call libraries of good quality using hand-released bats and which cover different environmental situations is titanic (O'Farrell et al., 1999). In fact, this has probably discouraged many researchers from developing their own classifiers so far.

Globally, echolocation call libraries are incomplete, especially in understudied regions such as most of the tropics (Aguilar, 2017). Due to species elusiveness, whispering behaviour or rarity, call libraries are usually only built with calls from a few hand-released individuals


**Table 1**

Performance of the random forest classifier for each species/sonotype based on calls from either free-flying (FF) or hand-released (HR) bats. Classification performance is ranked according to sensitivity and positive predictive value (see Methods for an explanation of the rationale underpinning this selection) as > 90% (dark green), 80–90% (olive green), and < 80% (light green).

Species/sonotypes	Acronym	Sensitivity		Positive Predictive Value (PPV)	
		FF	HR	FF	HR
<i>Pteronotus alitonus</i>	PA	0.96	0.92	0.99	0.80
<i>Pteronotus rubiginosus</i>	PR	0.96	0.95	0.99	0.73
Vespertilionidae 1	V1	0.95	0.76	0.96	0.51
<i>Myotis nigricans</i>	MN	0.93	0.32	0.96	0.70
<i>Furipterus horrens</i>	FH	0.93	0.88	0.00	0.00
<i>Centronycteris maximiliani</i>	CM	0.92	0.76	0.96	0.65
<i>Myotis riparius</i>	MR	0.91	0.83	0.98	0.97
<i>Saccopteryx bilineata</i>	SB	0.91	0.42	0.93	0.96
Emballonuridae 1	E1	0.90	0.00	0.21	0.02
<i>Cormura brevirostris</i>	CB	0.89	0.58	0.85	0.95
<i>Peropteryx kappleri</i>	PK	0.89		0.95	
<i>Peropteryx macrotis</i>	PM	0.88	0.27	0.60	0.02
<i>Saccopteryx leptura</i>	SL	0.88	0.76	0.95	0.89
<i>Pteronotus gymnonotus</i>	PG	0.86	0.17	0.17	0.05
Molossidae 3	M3	0.84	0.92	0.24	0.06
<i>Promops</i> sp.	P	0.82		0.11	
<i>Pteronotus personatus</i>	PP	0.80	0.01	0.09	0.81
Molossidae 1	M1	0.71	0.01	0.37	0.01
Molossidae 2	M2	0.59	0.27	0.44	0.17
<i>Rhynchonycteris naso</i>	RN	0.00	0.00	0.00	0.00

(Gager et al., 2016; Monadjem et al., 2017; Zamora-Gutiérrez et al., 2016). Although some studies have not found marked differences in automatic classifiers trained with data from distant regions (e.g. Zamora-Gutiérrez et al., 2016), other authors highlight the importance of taking these differences into consideration (Barclay et al., 1999; López-Baucells et al., 2017b; O’Farrell et al., 2000; Thomas et al.,

1987). Although we urge and support the compilation of comprehensive reference call libraries, our study suggests that training automatic classifiers with manually identified free-flying bats is a very valid option if it is cautiously used in conjunction with conservative classification criteria. As stressed by Jakobsen et al. (2013), it is of vital importance to record calls from naturally behaving bats in the wild and



**Fig. 4.** Percentage of files requiring visual post-validation as a function of the desired accuracy threshold for identification acceptance. Shaded area: Percentage of the whole dataset. Coloured lines: Percentage for each family; Species acronyms are as given in Table 1. Analysis was based on a random forest model, trained with 2000 pulses per species/sonotype (with the exception of *Rhynchonycteris naso* and *Furipterus horrens*, for which we had fewer recordings, see Methods).

use these recordings to improve classifier performances.

#### 4.4. Combining automatic classifiers with manual post-validation

No classifier has proved to provide 100% accuracy so far (Russo and Voigt, 2016). Therefore, some authors have recommended to manually validate all sound files (Kubista and Bruckner, 2017), which inevitably annihilates or at least greatly reduces the advantages of having automatic algorithms. In other cases, posterior cross-validation is completely neglected, which greatly affects the reliability of the study. According to our findings, even when aiming for an accuracy threshold of 95%, the remaining amount of data to be visually validated could be reduced by up to 75%. This represents a substantial saving in terms of workload.

Different acoustic analysis software with automatic classifiers has been released on the market in the last decades: batIdent (ecoObs, GmbH, Nürnberg, Germany), Kaleidoscope (Wildlife Acoustics, USA), Sonochiro (Sonochiro, France), Sonobat (Sonobat, USA), SCAN'R (Binary Acoustic Technology, USA) and more recently Tadarida (Bas et al., 2017), multiplying the options available to researchers to use technological advances to aid acoustic species identification. The best option for analysing the massive amounts of acoustic data generated by the latest recording devices without compromising the reliability of results, inevitably, lies in finding the right balance between automatic classification and manual cross-validation. This is especially true for threatened or rare species for which false positives will have greater conservation impact (Clement et al., 2014).

#### 4.5. Recommendations for effectively combining automatic and manual classification

Our approach, while highly versatile, requires that researchers must: A) have good knowledge about the bat fauna of the region (avoiding novice errors that result in misidentifications or passive acceptance of the results from any classifier and acknowledging regional and habitat variation), B) work together with experts on local call libraries and manual identifications, C) be skilled in programming in R or similar software packages, thus being able to adjust machine learning algorithms to particular situations, D) take into consideration both sensitivity and positive predictive values rather than global accuracies, E) define their own sonotypes conservatively (preventing classification to taxonomic levels that are not even visually distinguishable). We also recommend to base selection criteria on the PPV as the most conservative metric of performance (Armitage and Ober, 2010) since false negatives are always better than false positives.

## 5. Conclusions

Further research should focus on isolating and analysing individual call sequences instead of pulses, and analyse the whole sonogram rather than the pulses one by one (Damoulas et al., 2010; Kershenbaum et al., 2014; Ren et al., 2009). Our study shows how open-source statistical tools and software can be used to develop algorithms attaining similar levels of accuracy as commercial classifiers. However, their potential for wider application should be further explored with echolocation datasets from other regions. We also demonstrated that training algorithms with recordings from free-flying bats is possible and advisable if designed to classify recordings at sonotype level. This approach is not conceived to replace the use of calls from hand-released bats, but to aid in data management and classification with massive datasets. Combined with the availability of new low-cost automatic detectors and powerful supervised machine-learning algorithms, our analysis approach opens new opportunities for long-term monitoring programs to be undertaken by researchers in megadiverse regions where echolocation libraries are still scarce. In fact, in these regions, extended acoustic bat monitoring is urgently needed, and fortunately, the technical and

analytical tools are now at hand to do so.

## Acknowledgements

We would like to thank Oriol Massana, Diogo Ferreira, Marta Acácio, Fabio Farneda, Madalena Boto, Maria Mas, Gilberto Fernández and Eva Sánchez for fieldwork assistance and José Luis Camargo, Rosely Hipólito, Ary Jorge Ferreira, Luiz Quiroz, Leo and Josimar for logistical support. We thank two anonymous reviewers for helpful comments on an earlier version of the manuscript. Xavier Puig-Montserrat and Carles Flaquer also deserve special mention for their contribution to the discussion and continuous support. This work was supported by the Portuguese Foundation for Science and Technology under grants [PTDC/BIABIC/111184/2009] (CM), [SFRH/BD/80488/2011] (RR), [PD/BD/52597/2014] (ALB) and by the Foundation for Research Support of the State of Amazonas [FAPEAM 062.01173/2015] (PEDB). Additional funding was provided by a Bat Conservation International student research fellowship to ALB and RR. This is publication 749 in the BDFFP Technical Series.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ecoinf.2018.11.004>.

## References

- Aguilar, A.P.A., 2017. Morcegos insetívoros aéreos Neotropicals: identificação acústica e padrões de estruturação de assembleias. Universidade Federal do Rio Grande do Sul, Porto Alegre (135pp).
- Andreassen, T., Suriyke, A., Hallam, J., 2014. Semi-automatic long-term acoustic surveying: a case study with bats. *Ecol. Informa.* 21, 13–24. <https://doi.org/10.1016/j.ecoinf.2013.12.010>.
- Appel, G., López-Baucells, A., Magnusson, E., Bobrowiec, P.E.D., 2017. Aerial insectivorous bat activity in relation to moonlight intensity. *Mamm. Biol.* 85, 37–46. <https://doi.org/10.1016/j.mambio.2016.11.005>.
- Armitage, D.W., Ober, H.K., 2010. A comparison of supervised learning techniques in the classification of bat echolocation calls. *Ecol. Informa.* 5, 465–473. <https://doi.org/10.1016/j.ecoinf.2010.08.001>.
- Barclay, R.M.R., Fullard, J.H., Jacobs, D.S., 1999. Variation in the echolocation calls of the hoary bat (*Lasiurus cinereus*): influence of body size, habitat structure, and geographic location. *Can. J. Zool.* 77, 530–534. <https://doi.org/10.1139/z99-008>.
- Barkley, Y., Oswald, J.N., Carretta, J.V., Rankin, S., Rudd, A., Lammers, M.O., 2011. Comparison of Real-Time and Post-Cruise Acoustic Species Identification of Dolphin Whistles Using ROCCA (Real-Time Odontocete Call Classification Algorithm). Department of Commerce, California.
- Barlow, K.E., Briggs, P.A., Haysom, K.A., Hutson, A.M., Lechiara, N.L., Racey, P.A., et al., 2015. Citizen science reveals trends in bat populations: the National Bat Monitoring Programme in Great Britain. *Biol. Conserv.* 182, 14–26. <https://doi.org/10.1016/j.biocon.2014.11.022>.
- Bas, Y., Bas, D., Julien, J.-F., 2017. Tadarida: a toolbox for animal detection on acoustic recordings. *J. Open Res. Softw.* 5, 6. <https://doi.org/10.5334/jors.154>.
- Biscardi, S., Orprecio, J., Fenton, M.B., Tsoar, A., Ratcliffe, J.M., 2004. Data, sample sizes and statistics affect the recognition of species of bats by their echolocation calls. *Acta Chiropterol.* 6, 347–363. <https://doi.org/10.3161/001.006.0212>.
- Boyles, J.G., Sole, C.L., Cryan, P.M., McCracken, G.F., 2013. On estimating the economic value of insectivorous bats: prospects and priorities for biologists. In: *Bat Evolution, Ecology, and Conservation* New York, pp. 501–515.
- Breiman, L., 2001. Random forests. *Mach. Learn.* 45, 5–32.
- Briggs, F., Fern, X., Raich, R., 2009. Acoustic Classification of Bird Species from Syllables: An Empirical Study Oregon State University.
- Britzke, E.R., Duchamp, J.E., Murray, K.L., Swihart, R.K., Robbins, L.W., 2011. Acoustic identification of bats in the eastern United States: a comparison of parametric and nonparametric methods. *J. Wildl. Manag.* 75, 660–667. <https://doi.org/10.1002/jwmg.68>.
- Clement, M.J., Rodhouse, T.J., Ormsbee, P.C., Szewczak, J.M., Nichols, J.D., 2014. Accounting for false-positive acoustic detections of bats using occupancy models. *J. Appl. Ecol.* 51, 1460–1467. <https://doi.org/10.1111/1365-2664.12303>.
- Cunzio, G.C., Bernard, E., 2012. Neotropical bats as indicators of environmental disturbance: what is the emerging message? *Acta Chiropterol.* 14, 143–151. <https://doi.org/10.3161/150811012X654358>.
- Damoulas, T., Henry, S., Farnsworth, A., Lanzone, M., Gomes, C., 2010. Bayesian classification of flight calls with a novel dynamic time warping kernel. In: *Proceedings of the Ninth International Conference on Machine Learning and Applications*, pp. 424–429.
- de Oliveira, A.A., Mori, S.A., 1999. A central Amazonian terra firme forest. I. High tree species richness on poor soils. *Biodivers. Conserv.* 8, 1219–1244. <https://doi.org/10.1023/A:1008908615271>.
- Dickinson, J.L., Zuckerman, B., Bonter, D.N., 2010. Citizen science as an ecological research tool: challenges and benefits. *Annu. Rev. Ecol. Syst.* 41, 149–172. <https://doi.org/10.1146/annurev-ecolsys-102209-143302>.



- org/10.1146/annurev-ecolsys-102209-144636.
- Farneda, F.Z., Rocha, R., López-Baucells, A., Groenewald, M., Silva, I., Palmeirim, J.M., et al., 2015. Trait-related responses to habitat fragmentation in Amazonian bats. *J. Appl. Ecol.* 52, 1381–1391. <https://doi.org/10.1111/1365-2664.12490>.
- Ferreira, D.F., Rocha, R., López-Baucells, A., Farneda, F.Z., Carreiras, J.M.B., Palmeirim, J.M., et al., 2017. Season-modulated responses of Neotropical bats to forest fragmentation. *Ecol. Evol.* 7, 4059–4071. <https://doi.org/10.1002/ece3.3005>.
- Gager, Y., Tarland, E., Lieckfeldt, D., Menage, M., Botero-Castro, F., Rossiter, S.J., et al., 2016. The value of molecular vs. morphometric and acoustic information for species identification using sympatric molossid bats. *PLoS One* 11, e0150780. <https://doi.org/10.1371/journal.pone.0150780>.
- Gardner, A.L., 2007. *Mammals of South America 1: Marsupials, Xenarthrans, Shrews, and Bats*. University of Chicago Press, Chicago, pp. 690.
- Gregory, R.D., Van Strien, A., Vorisek, P., Meyling, A.W.G., Noble, D.G., Foppen, R.P., et al., 2005. Developing indicators for European birds. *Philos. Trans. R. Soc. Lond. Ser. B Biol. Sci.* 360, 269–288. <https://doi.org/10.1098/rstb.2004.1602>.
- Hill, A.P., Prince, P., Covarrubias, E.P., Doncaster, C.P., Snaddon, J.L., Rogers, A., 2018. AudioMoth: evaluation of a smart open acoustic device for monitoring biodiversity and the environment. *Meth. Ecol. Evol.* <https://doi.org/10.1111/2041-210X.12955>. (Online preview).
- Jakobsen, L., Brinklov, S., Surlykke, A., 2013. Intensity and directionality of bat echolocation signals. *Front. Physiol.* 4, 89. <https://doi.org/10.3389/fphys.2013.00089>.
- Jennings, N., Parsons, S., Pocock, M.J.O., 2008. Human vs. machine: identification of bat species from their echolocation calls by humans and by artificial neural networks. *Can. J. Zool.* 86, 371–377. <https://doi.org/10.1139/z08-009>.
- Jones, G., Kokurewicz, T., 1994. Sex and age variation in echolocation calls and flight morphology of Daubenton's bats *Myotis daubentonii*. *Mammalia* 58, 41–50. <https://doi.org/10.1515/mamm.1994.58.1.41>.
- Jones, G., Ransome, R.D., 1993. Echolocation calls of bats are influenced by maternal effects and change over lifetime. *Proc. R. Soc. Biol. Sci.* 252, 125–128. <https://doi.org/10.1098/rspb.1993.0055>.
- Jones, G., Jacobs, D.S., Kunz, T.H., Willig, M.R., Racey, P.A., 2009. Carpe noctem: the importance of bats as bioindicators. *Endanger. Species Res.* 8, 93–115. <https://doi.org/10.1035/esr00182>.
- Jung, K., Kalko, E.K.V., 2011. Adaptability and vulnerability of high flying Neotropical aerial insectivorous bats to urbanization. *Divers. Distrib.* 17, 262–274. <https://doi.org/10.1111/j.1472-4642.2010.00738.x>.
- Kershenbaum, A., Blumstein, D.T., Roch, M.A., Akcay, C., Backus, G., Bee, M.A., et al., 2014. Acoustic sequences in non-human animals: a tutorial review and prospectus. *Biol. Rev. Camb. Philos. Soc.* 91, 13–52. <https://doi.org/10.1111/brv.12160>.
- Kubista, C.E., Bruckner, A., 2017. Within-site variability of field recordings from stationary, passively working detectors. *Acta Chiropterol.* 19, 189–197. <https://doi.org/10.3161/15081109acc2017.19.1.015>.
- Kuhn, M., 2008. *Caret package*. *J. Stat. Softw.* 28, 1–26.
- Laurance, W.F., Camargo, J.L.C., Luizao, R.C.C., Laurance, S.G., Pimm, S.L., Bruna, E.M., et al., 2011. The fate of Amazonian forest fragments: a 32-year investigation. *Biol. Conserv.* 144, 56–67. <https://doi.org/10.1016/j.biocon.2010.09.021>.
- Laurance, W.F., Camargo, J.L.C., Fearnside, P.M., Lovejoy, T.E., Williamson, G.B., Mesquita, R.C.G., et al., 2017. An Amazonian rainforest and its fragments as a laboratory of global change. *Biol. Rev. Camb. Philos. Soc.* 93, 223–247. <https://doi.org/10.1111/brv.12343>.
- Law, B., Gonsalves, L., Tap, P., Penman, T., Chidel, M., 2015. Optimizing ultrasonic sampling effort for monitoring forest bats. *Aust. Ecol.* 40, 886–897. <https://doi.org/10.1111/aec.12269>.
- Lawrence, B.D., Simmons, J.A., 1982. Measurements of atmospheric attenuation at ultrasonic frequencies and the significance for echolocation by bats. *J. Acoust. Soc. Am.* 71, 585–590. <https://doi.org/10.1121/1.387529>.
- Lim, B.K., Engstrom, M.D., 2001. Bat community structure at Iwokrama Forest, Guyana. *J. Trop. Ecol.* 17, 647–665. <https://doi.org/10.1017/s0266467401001481>.
- López-Baucells, A., Rocha, R., Fernández-Arellano, G., Bobrowiec, P.E.D., Palmeirim, J.M., Meyer, C.F.J., 2014. Echolocation of the big red bat *Lasiurus egypticus* (Chiroptera: Vespertilionidae) and first record from the Central Brazilian Amazon. *Stud. Neotropical Fauna Environ.* 49, 18–25. <https://doi.org/10.1080/01650521.2014.907600>.
- López-Baucells, A., Rocha, R., Bobrowiec, P.E.D., Bernard, E., Palmeirim, J., Meyer, C., 2016. *Field Guide to Amazonian Bats*. INPA, Manaus, pp. 174.
- López-Baucells, A., Rocha, R., Tavares, V.C., Mora, L., Bobrowiec, P.E.D., Silva, S.E., et al., 2017a. Molecular, morphological and acoustic identification of *Eumops maurus* and *E. hansae* (Chiroptera: Molossidae) with new reports from Central Amazonia. *Trop. Zool.* <https://doi.org/10.1080/03946975.2017.1382284>.
- López-Baucells, A., Torrent, L., Rocha, R., Pavan, A.C., Bobrowiec, P.E.D., Meyer, C.F.J., 2017b. Geographical variation in the high-duty cycle echolocation of the cryptic common mustached bat *Pteronotus cf. rubiginosus* (Mormoopidae). *Bioacoustics* 1–17. <https://doi.org/10.1080/09524622.2017.1357145>.
- MacSwiney, G.C., Clarke, F.M., Racey, P.A., 2008. What you see is not what you get: the role of ultrasonic detectors in increasing inventory completeness in Neotropical bat assemblages. *J. Appl. Ecol.* 45, 1364–1371. <https://doi.org/10.1111/j.1365-2664.2008.01531.x>.
- Madhukumar Menon, A., João Ramos Pereira, M., Moura de Souza Aguiar, L., 2018. Are automated acoustic identification software reliable for bat surveys in the Neotropical region? *PeerJ*. <https://doi.org/10.7287/peerj.preprints.26712v1>.
- Millon, L., Julien, J.-F., Julliard, R., Kerbiriou, C., 2015. Bat activity in intensively farmed landscapes with wind turbines and offset measures. *Ecol. Eng.* 75, 250–257. <https://doi.org/10.1016/j.ecoleng.2014.11.050>.
- Monadjem, A., Shapiro, J.T., Mtsetfwa, F., Reside, A.E., McCreery, R.A., 2017. Acoustic call library and detection distances for bats of Swaziland. *Acta Chiropterol.* 19, 175–187. <https://doi.org/10.3161/15081109acc2017.19.1.014>.
- Mukherjee, G., Manna, S.S., 2006. Weighted scale-free networks in Euclidean space using local selection rule. *Phys. Rev. E* 74, 036111. <https://doi.org/10.1103/PhysRevE.74.036111>.
- Murray, K.L., Britzke, E.R., Hadley, B.M., Robbins, L.W., 1999. Surveying bat communities: a comparison between mist nets and the Anabat II bat detector system. *Acta Chiropterol.* 1, 105–112.
- O'Farrell, M.J., Miller, B.W., Gannon, W.L., 1999. Qualitative identification of free-flying bats using the Anabat detector. *J. Mammal.* 80, 11–23. <https://doi.org/10.2307/1383203>.
- O'Farrell, M.J., Corben, C., Gannon, W.L., 2000. Geographic variation in the echolocation calls of the hoary bat (*Lasiurus cinereus*). *Acta Chiropterol.* 2, 185–196.
- Olden, J.D., Lawler, J.J., Poff, N.L., 2008. Machine learning methods without tears: a primer for ecologists. *Q. Rev. Biol.* 83, 171–193. <https://doi.org/10.1086/587826>.
- Pedro, A.R.-S., Simonetti, J.A., 2014. Variation in search-phase calls of *Lasiurus varius* (Chiroptera: Vespertilionidae) in response to different foraging habitats. *J. Mammal.* 95, 1004–1010. <https://doi.org/10.1644/13-MammA-327>.
- Penone, C., Le Viol, I., Pellissier, V., Julien, J.-F., Bas, Y., Kerbiriou, C., 2013. Use of large-scale acoustic monitoring to assess anthropogenic pressures on orthoptera communities. *Conserv. Biol.* 27, 979–987. <https://doi.org/10.1111/cobi.12083>.
- Pio, D.V., Clarke, F.M., Mackie, I., Racey, P.A., 2010. Echolocation calls of the bats of Trinidad, West Indies: is guild membership reflected in echolocation signal design? *Acta Chiropterol.* 12, 217–229. <https://doi.org/10.3161/150811010X504716>.
- Puechmaille, S.J., Borissov, I.M., Zebok, S., Allegrini, B., Hizem, M., Kuenzel, S., et al., 2014. Female mate choice can drive the evolution of high frequency echolocation in bats: a case study with *Rhinolophus mehelyi*. *PLoS One* 9, e103452. <https://doi.org/10.1371/journal.pone.0103452>.
- Puig-Montserrat, X., Torre, I., López-Baucells, A., Guerrieri, E., Monti, M.M., Rafols-García, R., et al., 2015. Pest control service provided by bats in Mediterranean rice paddies: linking agroecosystems structure to ecological functions. *Mamm. Biol.* 80, 237–245. <https://doi.org/10.1016/j.mambio.2015.03.008>.
- Redgwell, R.D., Szewczak, J.M., Jones, G., Parsons, S., 2009. Classification of echolocation calls from 14 species of bat by support vector machines and ensembles of neural networks. *Algorithms* 2, 907–924. <https://doi.org/10.3390/a2030907>.
- Ren, Y., Johnson, M.T., Clemins, P.J., Darre, M., Glaeser, S.S., Osiejuk, T.S., et al., 2009. A framework for bioacoustic vocalization analysis using hidden markov models. *Algorithms* 2, 1410–1428. <https://doi.org/10.3390/a2041410>.
- Rocha, R., López-Baucells, A., Farneda, F.Z., Groenewald, M., Bobrowiec, P.E.D., Cabeza, M., et al., 2017a. Consequences of a large-scale fragmentation experiment for Neotropical bats: disentangling the relative importance of local and landscape-scale effects. *Landscape Ecol.* 32, 31–45. <https://doi.org/10.1007/s10980-016-0425-3>.
- Rocha, R., Ovaskainen, O., López-Baucells, A., Farneda, F.Z., Ferreira, D.F., Bobrowiec, P.E.D., et al., 2017b. Design matters: an evaluation of the impact of small man-made forest clearings on tropical bats using a before-after-control-impact design. *For. Ecol. Manag.* 401, 8–16. <https://doi.org/10.1016/j.foreco.2017.06.053>.
- Russo, D., Jones, G., 2002. Identification of twenty-two bat species (Mammalia: Chiroptera) from Italy by analysis of time-expanded recordings of echolocation calls. *J. Zool.* 258, 91–103. <https://doi.org/10.1017/S0952836902001231>.
- Russo, D., Voigt, C.C., 2016. The use of automated identification of bat echolocation calls in acoustic monitoring: a cautionary note for a sound analysis. *Ecol. Indic.* 66, 598–602. <https://doi.org/10.1016/j.ecolind.2016.02.036>.
- SCAN'R, 2009. *SCAN'R Snapshot Characterization and Analysis Software Manual*. Binary Acoustic Technology, pp. 35.
- Sewell, D., Guillera-Arroita, G., Griffiths, R.A., Beebe, T.J., 2012. When is a species declining? Optimizing survey effort to detect population changes in reptiles. *PLoS One* 7, e43387. <https://doi.org/10.1371/journal.pone.0043387>.
- Silva, C.R., Bernard, E., 2017. Bioacoustics as an important complementary tool in bat inventories in the Caatinga Drylands of Brazil. *Acta Chiropterol.* 19, 409–418. <https://doi.org/10.3161/15081109acc2017.19.2.017>.
- Szewczak, J., 2000. A tethered zip-line arrangement for reliably collecting bat echolocation reference calls. *Bat Res. News* 41, 142.
- Thomas, D.W., Bell, G.P., Fenton, M.B., 1987. Variation in echolocation call frequencies recorded from North American vespertilionid bats: a cautionary note. *J. Mammal.* 68, 842–847. <https://doi.org/10.2307/1381562>.
- Torrent, L., López-Baucells, A., Rocha, R., Bobrowiec, P.E.D., Meyer, C.F.J., 2018. The importance of lakes for bat conservation in Amazonian rainforests: an assessment using autonomous recorders. *Remote Sens. Ecol. Conserv.* <https://doi.org/10.1002/rse2.83>.
- Vassilios, S., Veronica, Z.-G., E.J.K., Mark, G., 2017. Bat echolocation call identification for biodiversity monitoring: a probabilistic approach. *J. R. Stat. Soc. Ser. C: Appl. Stat.* 67, 165–183. <https://doi.org/10.1111/rssc.12217>.
- Viera, A.J., Garrett, J.M., 2005. Understanding interobserver agreement: the Kappa statistic. *Res. Ser.* 37, 360.
- Walters, C.L., Freeman, R., Collen, A., Dietz, C., Fenton, M.B., Jones, G., et al., 2012. A continental-scale tool for acoustic identification of European bats. *J. Appl. Ecol.* 49, 1064–1074. <https://doi.org/10.1111/j.1365-2664.2012.02182.x>.
- Walters, C.L., Collen, A., Lucas, T., Mroz, K., Sayer, C.A., Jones, K.E., 2013. *Challenges of using bioacoustics to globally monitor bats*. In: *Bat Evolution, Ecology, and Conservation*. Springer, pp. 479–499.
- Whytock, R.C., Christie, J., Jones, K., 2017. Solo: an open source, customizable and inexpensive audio recorder for bioacoustic research. *Methods Ecol. Evol.* 8, 308–312. <https://doi.org/10.1111/2041-210X.12678>.
- Wordley, C.F.R., Foui, E.K., Mudappa, D., Sankaran, M., Altringham, J.D., 2014. Acoustic identification of bats in the Southern Western Ghats, India. *Acta Chiropterol.* 16, 213–222. <https://doi.org/10.3161/150811014x683408>.
- Zamora-Gutiérrez, V., López-González, C., MacSwiney González, M.C., Fenton, B., Jones, G., Kalko, E.K.V., et al., 2016. Acoustic identification of Mexican bats based on taxonomic and ecological constraints on call design. *Methods Ecol. Evol.* 7, 1082–1091. <https://doi.org/10.1111/2041-210X.12556>.