

Contents lists available at ScienceDirect

Journal of Thermal Biology

journal homepage: www.elsevier.com/locate/jtherbio

Extinction risks forced by climatic change and intraspecific variation in the thermal physiology of a tropical lizard



ournal of [HERMAL BIOLOG]

Emerson Pontes-da-Silva^{a,*}, William E. Magnusson^a, Barry Sinervo^b, Gabriel H. Caetano^b, Donald B. Miles^c, Guarino R. Colli^d, Luisa M. Diele-Viegas^e, Jessica Fenker^f, Juan C. Santos^{g,h}, Fernanda P. Werneck^a

^a Coordenação de Biodiversidade, Instituto Nacional de Pesquisas da Amazônia, Manaus, Brazil

^b Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, United States

^c Department of Biological Sciences, Ohio University, Athens, OH, United States

^d Department of Zoology, University of Brasília, Brasília, Brazil

^e Departamento de Ecologia, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

^f Division of Ecology and Evolution, Australian National University, Canberra, Australia

^g Biology Department, Brigham Young University, Utah, United States

h Department of Biological Sciences, St. John's University, Queens, NY, United States

ARTICLE INFO

Keywords: Amazon Cerrado Ecophysiology Global warming Preferred temperature Performance

ABSTRACT

Temperature increases can impact biodiversity and predicting their effects is one of the main challenges facing global climate-change research. Ectotherms are sensitive to temperature change and, although predictions indicate that tropical species are highly vulnerable to global warming, they remain one of the least studied groups with respect to the extent of physiological variation and local extinction risks. We model the extinction risks for a tropical heliothermic teiid lizard (Kentropyx calcarata) integrating previously obtained information on intraspecific phylogeographic structure, eco-physiological traits and contemporary species distributions in the Amazon rainforest and its ecotone to the Cerrado savannah. We also investigated how thermal-biology traits vary throughout the species' geographic range and the consequences of such variation for lineage vulnerability. We show substantial variation in thermal tolerance of individuals among thermally distinct sites. Thermal critical limits were highly correlated with operative environmental temperatures. Our physiological/climatic model predicted relative extinction risks for local populations within clades of K. calcarata for 2050 ranging between 26.1% and 70.8%, while for 2070, extinction risks ranged from 52.8% to 92.8%. Our results support the hypothesis that tropical-lizard taxa are at high risk of local extinction caused by increasing temperatures. However, the thermo-physiological differences found across the species' distribution suggest that local adaptation may allow persistence of this tropical ectotherm in global warming scenarios. These results will serve as basis to further research to investigate the strength of local adaptation to climate change. Persistence of Kentropyx calcarata also depends on forest preservation, but the Amazon rainforest is currently under high deforestation rates. We argue that higher conservation priority is necessary so the Amazon rainforest can fulfill its capacity to absorb the impacts of temperature increase on tropical ectotherms during climate change.

1. Introduction

Climate change is occurring at different spatial scales (IPCC, 2014), and responses to rising temperatures have included changes in phenology, shifts in elevation and local extinction of populations (Colwell et al., 2008; Sinervo et al., 2010; Winter et al., 2016). Several studies suggest that tropical ectotherms are more vulnerable to warming temperatures than species farther from the equator (e.g., Deutsch et al., 2008; Sinervo et al., 2010; Huey et al., 2012). Thus, climate change in tropical regions is expected to induce a decrease in biodiversity, because these regions tend to be the most affected by rapid increases in temperature as well as they support the highest numbers of species (Dillon et al., 2010).

Current species distribution models (SDMs) mainly describe the environmental niches of focal species through correlations between climatic factors and distributional data, without incorporating species-

* Corresponding author.

E-mail address: emersonpontesbio@gmail.com (E. Pontes-da-Silva).

https://doi.org/10.1016/j.jtherbio.2018.01.013 Received 9 September 2017; Received in revised form 11 January 2018; Accepted 31 January 2018

Available online 09 February 2018 0306-4565/ © 2018 Elsevier Ltd. All rights reserved. specific ecophysiological predictors (e.g., body and operative environmental temperatures) in the modeling (Kearney and Porter, 2009). However, such SDMs may be limited because they are not based on traits measured directly from the physiology of species. In contrast, mechanistic SDMs might overcome these drawbacks by using physiological data to represent the complexity of interrelationships between organismal physiology, microclimate, behavior, and distribution (Gunderson and Leal, 2012).

Climate change has been shown to have global impacts on reptile diversity (Winter et al., 2016) and to cause local declines in lizard diversity (Sinervo et al., 2010). Such climatic disturbances likely caused changes in thermal niches by reducing hours of activity (H_a) and increasing hours of restriction (i.e., H_r or the total number of daylight hours in which environmental temperatures are above or below critical limits that restrict lizard activity). Models estimating extinction risks on a global scale indicate that by 2070, > 39% of local populations of ~20% of lizard species could become extinct (Sinervo et al., 2010). These predictions are especially alarming for large and biodiverse areas in the Neotropics, such as the Amazon rainforest, the Cerrado savanna and their ecotone in South America, which are expected to become high extinction-risk regions for lizards (Sinervo et al., 2010).

Until recently, SDMs based on ecophysiology were mainly premised on extinction risk across lizard families. As they were not based on individual species, they need further refinement and validation with field data obtained in situ (Sinervo et al., 2010). Recent studies on lizards have refined estimates of extinction-risk at local scales, considering the interrelations between physiology and microclimate (e.g., Ceia-Hasse et al., 2014; Lara-Resendiz et al., 2015; Kubisch et al., 2015). However, few such analyses exist for lizard species inhabiting tropical rainforests (Sinervo et al., 2010).

Individuals of the same species at distinct locations are subject to variation in local environmental conditions, which is particularly true for species with broad geographic distributions (Hoffman and Sgro, 2011). Geographic variation in physiological traits between and within populations, which could result from plasticity or adaptation, indicates that climate sensitivity of tropical species may not be as rigid as previously believed (Huey et al., 2009; Llewelyn et al., 2016) and reveal the spatial structure of adaptive potential throughout species distributions (Moritz et al., 2012). For example, differences in physiological traits and sensitivity to temperature can confer differences in vulnerability of populations according to local habitat structure (Piantoni et al., 2016).

Differences in physiological traits have been revealed between species from temperate and tropical regions. For example, differences in locomotor performance at different body temperatures between different populations within a clade were detected in skinks in Australia (Moritz et al., 2012) and *Anolis sagrei* populations in the Bahamas (Logan et al., 2014). However, differences in performance were not found in *A. cristatellus* on Puerto Rico, which is a close relative of *A. sagrei* (Gunderson and Leal, 2012). Body temperature can vary with sex, age, and season, as well as among individuals at different locations (e.g.,Van Damme et al., 1986; Gunderson and Leal, 2012). Clusella-Trullas et al. (2011) suggested that differences in preferred temperature might be more related to other climatic variables, such as variation in environmental temperature and precipitation, than to mean temperature.

Critical thermal limits (CT_{min} and CT_{max}) are considered key predictors of ectotherm physiological sensitivity, and differences in both traits have been used to investigate macrophysiological patterns relative to temperature and precipitation (e.g., Clusella-Trullas et al., 2011), latitude (e.g., Addo-Bediako et al., 2000; Araujo et al., 2013), and habitat (e.g., terrestrial versus marine; Sunday et al., 2011). Likewise, critical thermal limits may vary intraspecifically (e.g., Moritz et al., 2012; Logan et al., 2014), but the extent of their variation remains underexplored (Losos, 2008). Uncovering intraspecific



Fig. 1. Kentropyx calcarata photographed in Ribeirão Cascalheira, Mato Grosso, Brazil by Guarino R. Colli.

differences may help clarify whether critical thermal limits are evolutionarily conserved or labile (Pearman et al., 2008). The upper critical temperature is known to vary little and tends to be conserved among populations (Moritz et al., 2012), and among ectotherm species (Addo-Bediako et al., 2000; Araujo et al., 2013). Thus, it is assumed that temperature increases will impose selective pressure over a wide range of species and populations (Sunday et al., 2011).

Regional diversification of some tropical lineages has been described and can be integrated with evolutionary-physiological studies (Moritz et al., 2012). Here, we use this approach using the lizard species Kentropyx calcarata (Spix 1825; Squamata: Teiidae) as a model system (Fig. 1). The species has a broad distribution in the northeastern part of South America including Brazil and the Guianas. Individuals are found in forest environments near streams, natural clearings or forest edges (Vitt, 1991). The species has a relatively well-known phylogeographic history with at least ten main intraspecific lineages previously identified (Werneck et al., 2009; Avila-Pires et al., 2012; Cronemberger, 2015). Given this species distribution and the assumed conservative nature of thermal traits typical of tropical ectotherms (Deutsch et al., 2008; Huey et al., 2009), K. calcarata is predicted to be sensitive to climate change. Most open-habitat heliotherms occupy heterogeneous thermal environments and are thermal generalists, which may explain their wide geographic distributions (Bonino et al., 2015). Kentropyx calcarata thermal-microhabitat use contrasts with those of heliotherms of open areas and forest nonheliotherms (Vitt et al., 1991; Vitt et al., 1997). Thus, K. calcarata belongs to an under-explored group in relation to its thermal biology.

Most teiid lizards are heliotherms that maintain body temperatures above air or ambient temperatures during activity and avoid heat stress by shuttling between sun and shade (e.g., Sartorius et al., 1999). Behavioral thermoregulation may buffer lizards from warming impacts. However, estimates of mean field or preferred temperature may not be suitable to estimate restriction in activity times for heliothermic lizards (Kearney and Porter, 2009). For heliothermic lizards, the voluntary thermal maximum (VTM) for activity is probably a better threshold to indicate activity limitation (Vitt et al., 1997).

Here, we estimate relative extinction risks for *K. calcarata* in large sub-regions within the Brazilian Amazon and the Cerrado-Amazon ecotone occupied by four of the main intraspecific lineages recognized by Cronemberger (2015), based on the calculation of thermally imposed hours of restriction, alone and in interaction with rainfall and vegetation cover. We also investigate how the thermal biology of *K. calcarata* is affected by different thermal environments to predict how differences in thermal biology might affect vulnerability of populations from different regions.



Fig. 2. Study area and locality records of *Kentropyx calcarata* used in extinction-risk modeling across four sub-regions in Dense forests (Amazon areas) and Open forests (Cerrado savannas areas). The geographic limits used to sub-select sampling points were based on the distribution of *Kentropyx calcarata* intraspecific lineages inferred from Cronemberger (2015) as follows: green circles refer to the Amazona sub-region that correspond to Amazonas State (AM) north of the Amazon river extending to Roraima (RR); yellow circles refer to the Amapá State (AP); white circles refer to the Tocantins sub-region that corresponds to sampling points in Tocantins State (TO) and extends to Maranhão (MA) and eastern Pará (PA); blue circles refer to the Pará sub-region that corresponds to sampling points in Pará State, interfluve of Xingu River (west), Araguaia River (east) and Amazon River (north). Red stars depict the sampling points for *K. calcarata* ecophysiological data, as follows: 1 – Floresta Nacional do Amapá, Amapá; 2 – Floresta Nacional de Caxiuanã, Pará; 3 – Reserva Florestal Adolpho Ducke, Amazonas; and 4 –Centro de Pesquisas Canguçu, Tocantins. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

2. Methods

2.1. Study area and sample method

Specimen sampling and experiments were carried in the Brazilian Amazon in September 2014 and between July and October 2015. These months correspond to the dry season in all sampling and experimental locations. The four sampling localities where chosen based on the regions occupied by four different intraspecific clades recovered by recent phylogeographic studies (Avila-Pires et al., 2012; Cronemberger, 2015): 1 – Floresta Nacional (FLONA) de Caxiuanã (1°44' S, 51°27' W; elevation 28 m), Melgaço/Portel, Pará State (PA); 2 – Reserva Florestal Adolpho Ducke (2°57' S, 59°55' W; elevation 68 m), Manaus, Amazonas State (AM); 3 – Floresta Nacional (FLONA) do Amapá (0°55' N, 51°36' W; elevation 14.4 m), Amapá State (AP); and the Cerrado-Amazon ecotone: 4 – Centro de Pesquisas Canguçu (9°56' S, 49°47' W; elevation 150 m), Pium, Tocantins State (TO) (Fig. 2).

The three Amazon sampling localities are covered by rainforests, have high mean annual precipitation (2200 mm, 2272 mm and 2300 mm for AM, PA and AP, respectively), but similar mean annual temperatures and small annual temperature range (approx. 2 °C difference between the coldest and hottest months). The Cerrado-Amazon ecotone sampling locality is characterized by transitional forest, where many trees are seasonally deciduous (Marimon et al., 2006), with mean annual rainfall of 1760 mm and a more variable climate (INMET, 2016).

At each sampling site, individuals of *K. calcarata* were captured by active search, nooses, pitfall traps and funnel traps. Each lizard was

housed individually in containers with base area of 22.5 cm \times 15.0 cm and height of 8.0 cm at room temperature (approximately 25–30 °C) and provided with water ad libitum until the experiments were performed on the same day of collection in a nearby field lab.

Collected individuals and tissue samples were deposited at institutions listed in Appendix II. All protocols involving live animals were approved by the Comissão de Ética no Uso de Animais na Pesquisa of the Instituto Nacional de Pesquisas da Amazônia (INPA) (Code: 029/ 2014).

We restricted extinction-risk modeling (see below) to distribution records from areas adjacent to where physiological data were collected and within areas occupied by the four *K. calcarata* intraspecific clades (Cronemberger, 2015). For the **Tocantins sub-region**, we considered geographic points east of the Araguaia River, except for disjunct records on the east coast; for the **Amazonas sub-region** we considered geographic records in the area around Manaus extending to Boa Vista (Roraima) and south of Pantepui; for the **Pará sub-region** we considered geographic records in the interfluves of the Xingu River (west), Araguaia (east) and Amazonas (north) Rivers; and for the **sub-region of Amapá** we considered geographic records from the entire Amapá State (Fig. 2).

Geographic (SDM-input) data were assembled that included a total of 2013 points distributed between sub-regions as follows: Pará (416), Amazonas (82), Amapá (449) and Tocantins (1066). When points were separated by 1 km or less from the next point (defined as one locality; Sinervo et al., 2010), we randomly selected one to be used in modeling. Geographic records were available in online databases of Scientific Collections and Museums records (Appendix I).

2.2. Operative temperatures, occupancy index and extinction risks

To explore the natural thermal environment (i.e., microclimate) of K. calcarata individuals in each location, we recorded local operative (Te) and air (Tair) temperatures. Laboratory data were collected simultaneously with operative-temperature recording in all locations. For this purpose, we used temperature data-loggers (HOBO [™] Onset Pro V2, approximately 25 by sampling locality) connected to PVC tubes matching adult K. calcarata in size and shape (3 cm in diameter, 14 cm long). The Te estimates using PVC pipe models has been validated against body temperatures for small ectotherms (Adolph, 1990; Sinervo et al., 2010; Lara-Reséndiz et al., 2014; Lara-Resendiz et al., 2015, Kubisch et al., 2015; Belasen et al., 2016; Kirchhof et al., 2017). We also used data-loggers unconnected to models to record air temperatures (30 by sampling locality). Each temperature data-logger had two probes, each inserted in a model tube. We placed temperature recorders in microhabitats most likely occupied by lizards (natural clearings, edges of the forest and stream borders) at heights up to two meters (approximated natural height used by K. calcarata), always with a probe shaded by leaves and the other placed in a position exposed to sunlight (Sinervo et al., 2010). Temperature was recorded every two minutes throughout the field period. We considered the temperature recorded by the probes to represent the operative temperature (Te) of K. calcarata in each sampling locality. We selected only temperature records within the time period when the species is active, based on the time of the first and the last sighting in the field (07:00 h and 16:00 h).

We fit a species-distribution model, using hours of activity, hours of restriction to activity, precipitation and vegetation surfaces, and interactions between those surfaces, as predictors for the distribution of *Kentropyx calcarata* across each sub-region. The hours of activity surfaces were compound from two previous surfaces, each one with different assumptions about the animals' physiology (heliotherm and nonheliotherm). Those surfaces were combined using a vegetation surface as a spatial criterion. This was a simple binary surface indicating where forests or open areas are predominant. Data from the nonheliotherm surface were assigned to forest areas and data from the heliotherm surface were assigned to open areas, creating the compound variable hours of activity surface. The same was done for hours of restriction.

We define hours of activity as the number of hours in a day in which the animal's body temperature was above lower thermal limit (T_{lower}), and hours of restriction as the number of hours above the upper thermal limit (T_{upper}), following Sinervo et al. (2010). Other than the choice of thermal threshold (T_{lower} or T_{upper}), all the calculations are identical for both surfaces. We calculated the extinction risks based on highest body temperatures recorded in the field instead of preferred body temperatures estimated in laboratory. T_{lower} is defined as the lowest air temperature at which the species was found active (26.9 °C) and T_{upper} as highest body temperature at which the species was found active (40.2 °C).

For the nonheliotherm surfaces, we simulated daily temperature variation across the projected area (See Fig. 2), by constructing a sinusoidal curve ranging from minimum to maximum temperature values for each cell, extracted respectively from WorldClim minimum and maximum temperature surfaces (Hijmans et al., 2005, 2.5 min resolution). Then we counted how many hours at each cell fell outside the established thermal threshold (T_{lower} or T_{upper}). Those values would then be used to create the nonheliotherm hours of activity or restriction surfaces, for each month of the year.

For the heliotherm surfaces, we used operative temperature data (T_e) collected from microhabitats used by the species at locations where the species was sampled. This data gave us actual measures of daily temperature variation, which we then used to count the number of hours outside thermal thresholds (T_{lower} or T_{upper}) on each day and in each microhabitat. However, since this data was representative only of the locations where *K. calcarata* was collected, we used a model to extrapolate those values across the projected area. This was done by fitting a generalized logistic model (Richard's curve) of hours above threshold to maximum air temperature, collected for the same location and days where the operative temperature measures were taken. This model was then applied to a WorldClim maximum temperature surface in order to predict hours of restriction and hours of activity across the projected area.

The heliotherm and nonheliotherm surfaces were then compounded using a forest surface, as described above, creating the final hours of activity and restriction surfaces for each month of the year. Each monthly variable (hours of restriction, activity and precipitation) was also averaged across the animal's breeding season and also across the months outside the breeding season. All the calculations were done in R. Those surfaces, together with the forest surface and the interactions between surfaces were used as predictors in species-distribution models ensembled from GLM, GAM and MaxEnt algorithms, using 1000 pseudo absence points per run, done with the package biomod2 (Thuiller et al., 2005) in program R. Then, model predictions were extrapolated to the years 2050 and 2070, using WorldClim climate projections under the RCP 4.5 scenario of the global climate model MPI-GSM-LR.

From each future projection, we extracted the occurrence probability or Occupancy Index of *Kentropyx calcarata* at each known current point of occurrence. We then subtracted those probabilities from the occurrence probabilities in the present for the same points, which gives us a probability of the species being extant at that point at that year, complementary to the probability of being extinct. Further details on the modeling and rationale can be found on Appendix III.

2.3. Thermal-trait estimates

We measured the thermal traits of the individuals of *K. calcarata* through the following experiments, with their respective sample sizes of

Table 1

Thermal physiology (preferred temperatures - Tpref and critical thermal limits - CTmin and CTmax) and operative temperatures (Te) of Kentropyx calcarata	at four
sampling locations. We present the sample sizes (n), mean estimates / variance (var), and the minimum and maximum values for each location in parent	theses.

	T _{pref} (°C)	CT _{min} (°C)	CT _{max} (°C)	T _e (°C)
Amapá	n = 17	n = 5	n = 5	24.6 ± 1.1
	29.9 / var = 5.2	7.8 / var = 27.9	39.8 / var = 12.8	(21.7 – 29.5)
	(24.5-33.1)	(3.5–16.5)	(34.9–43.7)	
Amazonas	n = 22	n = 7	n = 7	26.8 ± 1.0
	31.3 / var = 8.1	11.1 / var = 9.3	40.7 / var = 37.8	(23.5 - 31.3)
	(23.3-35.4)	(9.4–18.0)	(31.8–49.2)	
Pará	n = 8	n = 7	n = 7	31.1 ± 5.9
	30.4 / var = 7.2	14.0 / var = 41.6	41.4 / var = 11.0	(22.7 – 46.3)
	(24.8-33.7)	(5.1-23.0)	(37.9–47.1)	
Tocantins	n = 61	n = 10	n = 10	32.1 °C ± 7.8
	37.1 / var = 28.7	12.8 / var = 1.1	43.18 / var = 0.8	(15.4 – 64.3)
	(30.8-40.2)	(10.6–14.3)	(41.3-44.4)	

Table 2

Significance of coefficients obtained with Generalized Linear Models (GLM) for each model parameter: cumulative hours of restriction (Total_hr), hours of activity (Total_ha), and precipitation (Total_prec), and the quadratic terms: precipitation (Sqprec: Total_prec²), hours of restriction (Sqhr: Total_hr²), hours of activity (Sqha: Total_ha²) and the interactions between precipitation and hours of restriction (Prechr: Total_prec x Total_hr), precipitation and hours of activity (Precha: Total_prec x Total_ha) and vegetation cover (Forest Layer). Degrees of Freedom: 1.094 (total).

	Estimate	Std. Error	z value	p value
Intercept Total_h _r Total_ha Total_prec Sqprec Sqh _r Sqh _a Basilia	-1.796e + 02 4.195e + 01 1.076e + 01 1.232e - 02 1.461e - 06 1.864e - 03 1.909e - 03 0.909e - 02	3.765e + 01 5.830e + 00 1.364e + 00 1.745e - 03 5.859e - 07 4.567e - 04 5.927e - 04 1.012e - 02	- 4.771 7.195 7.891 7.068 2.494 4.082 3.221	 < 0.0001
Prechr Precha	– 9.328e – 08 2.287e – 07	1.913e – 08 2.346e – 08	4.877 9.747	< 0.0001 < 0.0001
Prechr Precha	– 9.328e – 08 2.287e – 07	1.913e - 08 2.346e - 08	4.877 9.747	< 0.0001 < 0.0001
Forest Layer	1.676e + 02	3.735e + 01	4.488	< 0.0001

the all sub-regions: Preferred temperature - T_{pref} (108), Critical thermal minimum - CT_{min} (29), Critical thermal maximum - CT_{max} (29) and Sensitivity of thermal performance (33). We used different individuals for the different experiments, except for CT_{min} and CT_{max} . The specific sample sizes of each region are detailed on Tables 1, 3.

2.4. Preferred temperature

The preferred temperature of individual lizards was estimated in a thermal gradient with base area of $2.0 \text{ m} \times 0.3 \text{ m}$ and 0.65 m high. The gradient was divided into lanes, where individuals were placed. A heat lamp was placed at one end of the lane to maintain a substrate temperature of 48 °C, simulating direct daylight, while gel ice packs and a fan kept the other end of the lane at an approximate temperature of 21 °C, which simulated a natural shaded microenvironment.

We measured the initial body temperature of individuals before placing them in the lane for the experiment. Body temperature was measured with an infrared laser thermometer (Digitech QM7221) at a distance of < 0.25 m or through an automated data logger with thermocouple probes attached to the belly of each individual. Both estimate instruments were calibrated against each other. We validated the use of body temperatures based on infrared thermometers with data on *Zootoca vivipara*, with high correlation between core and surface temperatures (0.85; n = 34, P < 0.001). This species is in similar size to most of the lizards in our data set. Before each round of estimates, lizards were left undisturbed in the thermal gradient for 10–15 min of acclimation. Regardless of the method, T_{pref} estimation lasted 2 h with body temperatures measured every 5 min, resulting in 25 estimates per individual. We used the mean temperature of the 25 estimates to represent each individual's preferred temperature, T_{pref}.

2.5. Sensitivity of thermal performance

We measured individual ability to sustain physical activity to simulate natural activity conditions where mobility is required for territorial defense, escape from predators or foraging (Robson and Miles, 2000). Lizards were stimulated to run in a circular track by gentle touches and only using data when lizards did not run to the sides (Clobert et al., 2000). The track was made of aluminum, with a diameter of the inner wall of 60 cm and of the outer wall of 150 cm. Lizards ran in a track with a width of 25 cm. We used the soil from each collection site as a substrate to mimic natural conditions.

We measured the endurance elapsed until exhaustion and motorresponse loss of individual locomotion. We estimated the endurance to account for effects of size and temperature on speed (distance) (Clobert et al., 2000).

Each individual was acclimated to run at seven different body temperatures (15 °C, 20 °C, 25 °C, 30 °C, 35 °C, 40 °C and 43 °C) in a randomized sequence. The test started when the individual's body temperature was within 0.5 °C of the target temperature. Lizards had a rest interval of 4 h between each race at a new temperature. Within individuals, running temperatures were given in a random sequence. The substrate temperature of the circular track was adjusted to each running body temperature through intercalation of the track between shadow and sun. The optimum performance temperature was estimated based on the body temperature at which each individual had the best performance (Huey and Stevenson, 1979).

2.6. Thermal tolerance: CT_{min} and CT_{max}

The thermal-tolerance range represents the physiological performance interval in which the activities of an organism are possible and is limited by the critical thermal minimum (CT_{min}) and maximum (CT_{max}) temperature (Huey and Stevenson, 1979). We estimated CT_{min} and CT_{max} of *K. calcarata* individuals at each study site using cold and heat sources to manipulate body temperature to a critical limit while monitoring at approximately 2-min intervals the loss of righting response, and individual body temperature was taken with a quick-reading cloacal thermometer (Miller and Weber 4K8893) when the individual lost motor function.

To estimate CT_{min} , lizards were isolated in Styrofoam boxes with gel ice packs. To estimate CT_{max} , lizards were directly exposed to sunlight or to bottles containing heated water placed in styrofoam boxes. To represent the critical thermal limit, we used the body temperature at which the individual did not move after being turned belly up (Clobert et al., 2000). We used the difference between CT_{max} and CT_{min} to represent the thermal-tolerance range of each individual (Amp_{tol}).

2.7. Data analysis

We compared the thermal biology among the four sampling localities using either an analysis of variance (ANOVA) or Kruskal-Wallis test. When significant (P < 0.05) differences were found in mean trait values among sites, we used the Tukey *post hoc* test to identify pairwise differences. The shapes of thermal performance curves (TPC's) of individuals from each site were estimated and plotted using generalized additive mixed models (GAMM), using the function "gamm" in the package mgcv (Woods, 2006) and ggplot2 (Wickham, 2009). Because size also may influence locomotor performance, we included snout-to-

Table 3

Results of generalized additive mixed models describing thermal performance curves for *Kentropyx calcarata* at the four sampling locations. Statistics are given for the effect of temperature and SVL on performance (temperature - Temp and snout-to-vent length - SVL, respectively), R^2 value, the optimum temperature for the performance estimates (T_{opt}), and the temperature range in which at least 80% of the maximum performance of the individual is possible.

	Temp	SVL	R ²	T _{opt} (°C)	B ₈₀ (°C)
Amazonas $(n = 10)$	$\begin{array}{l} F = 31.6; P = .0001 \\ F = 8.8; P = .0002 \\ F = 5.2; P = .0003 \\ F = 34.2; P = .0001 \end{array}$	P = .2	0.5	37.2	34.3–39.4
Pará $(n = 5)$		P = .7	0.2	29.1	24.2–33.9
Amapá $(n = 8)$		P = .4	- 0.08	25.8	21.6–29.4
Tocantins $(n = 10)$		P = .0001	0.5	35.9	33.3–38.1



Fig. 3. Kentropyx calcarata occupancy index for the studied sub-regions, based on the habitat suitability for the present (A), 2050 (B) and 2070 (C). The color scale indicates a gradient of favorable areas for species occurrence, ranging from blue (1.0 - suitable areas) to red (0.0 - unsuitable areas). Black dots refer to the current *K. calcarata* occurrence records in the four sub-regions used in partitioned modeling (models were estimated separately for each sub-region but plotted simultaneously). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

vent length (SVL) as a covariate. Sex of individuals not was used as a covariate. Gravid females and immature individuals were not used in the tests. To compute the curve, CT_{max} and CT_{min} mean values of individuals from each location were given a performance value of 0. The breadth of optimum performance was estimated as the range of body temperatures at which lizards attained 80% (B₈₀) of maximum performance, by multiplying the maximum performance speed at the best temperature (T_{opt}) by 0.8 (Huey and Stevenson, 1979). Statistical analyses were performed with R (R Development Core Team, 2014). Results are shown using mean \pm standard deviation and sample sizes (*n*).

3. Results

3.1. Operative temperatures, occupancy index and extinction risks

The estimated operative temperatures for *K. calcarata* differed significantly between sampling sites ($F_{3,13} = 91$, P = 0.0001). At the Tocantins site, models had higher mean T_e and an amplitude 93% higher than in other locations (Table 1). At Amapá and Amazonas sites, models had the lowest T_e (7.5 °C lower than TO) and were more thermally homogeneous. T_e at Pará site was higher and also had greater variability than in other Amazonian sites (Table 1).

The traits H_r and $H_{a,}$ as well as precipitation and historical Amazon forest cover, were significant explanatory variables in our SDM, along with most of their interaction effects and quadratic terms (Table 2).

The occupancy index in the present (Fig. 3a) for the Tocantins subregion was of 83.8%, 70.9% for the Amapá sub-region, 61.8% for the Amazonas sub-region and 74.3% for the Pará sub-region. The average extinction risk of local populations for 2050 (Fig. 3b) in the Pará subregion was 57.8%, 60.2% for the Tocantins sub-region, 26.1% for the Amazonas sub-region and 70.8% for the Amapá sub-region. The mean extinction risk of populations for 2070 (Fig. 3c) in the Pará subregion was of 85.2%, 86.6% for the Tocantins sub-region, 52.8% for the Amazonas sub-region and 92.8% for the Amapá sub-region.

3.2. Preferred temperatures and thermal tolerances

Individuals of *K. calcarata* from different locations exhibited significant differences in T_{pref} (F_{3,94} = 123, P < 0.0001; Appendix IV). Tocantins individuals selected temperatures 23% higher (37.1 ± 5.3 °C) than the mean T_{pref} of other locations (AM: 31.3 ± 2.8 °C; PA: 30.4 ± 2.6 °C and AP: 29.9 ± 2.2 °C). Lizards from the remaining three locations exhibited no significant difference in T_{pref} (Table 1).

No significant differences were found in mean thermal-tolerance range among the sampling locations, although there was a trend for differences in minimum critical temperature (CT_{min}) among locations



Fig. 4. Relationship between maximum operative temperature $(T_{e,max})$ and K. calcarata critical thermal maximum (CT_{max}) measured in the laboratory for the four studied locations.

(CT_{min}, P = 0.08; CT_{max}, P = 0.3; Amp_{tol}, P = 0.5). Amapá individuals had the lowest CT_{min} (6.2 °C difference from PA individuals, which had the highest CT_{min}). In contrast, variances of minimum and maximum critical limits were significantly different between locations (Levene test, CT_{min}, F_{3,26} = 3.2, P = 0.03; CT_{max}, F_{3,27} = 3.6, P = 0.02). Individuals from Pará (CT_{min}, 14.0 ± 6.4 °C; CT_{max}, 41.4 ± 3.3 °C), Amazonas (CT_{min}, 11.1 ± 3.0 °C; CT_{max}, 40.7 ± 6.1 °C), and Amapá (CT_{min}, 8 ± 5.2 °C; CT_{max}, 39.8 ± 3.5 °C) were similar to each other, showing greater variability in thermal-tolerance estimates than Tocantins individuals (CT_{min}, 12.8 ± 1.1 °C; CT_{max}, 43.1 ± 0.8 °C) (Table 1).

There was no significant relationship between mean operative temperature or maximum operative temperature ($T_{e,max}$) and T_{pref} , T_{opt} , or the upper temperature limit for 80% of maximum performance measured in the laboratory ($P \ge 0.3$ in all cases). However, there was a strong positive relationship between maximum operative temperatures ($T_{e,max}$) and critical thermal maximum (CT_{max}) measured in the laboratory ($CT_{max} = 37 \pm 0.1 * T_{e,max}$, $F_{(3,45)} = 37.1$, P = 0.02; Fig. 4).

3.3. Thermal sensitivity of locomotor performance

Body temperatures significantly affected individual performance at all sampling locations (AM: $F_{(3,38)} = 31.6$, P = < 0.0001, $R^2 = 0.5$; PA: $F_{(3,46)} = 8.8$, P = 0.0002, $R^2 = 0.2$; AP: $F_{(4,27)} = 5.2$, P = 0.0003, $R^2 = 0.08$ and TO : $F_{(5,91)} = 34.2$, P = 0.0001, $R^2 = 0.5$), with reduced performance at higher temperatures. We detected a significant effect (P < 0.0001) of SVL on locomotor performance only among Tocantins individuals (Table 3), with smaller individuals exhibiting lower locomotor performance than larger ones, generating a bimodal performance curve (Appendix V). We tested males and females from each location separately to evaluate possible sex effects on performance before calculating performance curves, but there were no statistically significant sexual differences in any sampling location (P > 0.05, in all cases) and, therefore, analyses were conducted without regard to sex.

The optimum-performance temperature was highest at the Amazonas site (37.2 °C) and lowest for Amapá individuals (25.8 °C), while Tocantins (35.9 °C) and Pará (29.1 °C) individuals exhibited better performance at intermediate temperatures (Table 3). Individuals from PA, AP and TO selected higher body temperatures than their optimum-performance temperatures, whereas the estimated T_{pref} for AM individuals was lower than the optimum-performance temperature (Table 1).

The optimum-performance temperature was higher at sampling sites where the 80% performance breadth (B_{80}) was narrower (TO and AM), while in places where AT-80% was wider (PA and PA), individuals showed better locomotor performance under a lower T_{opt} (Table 3), reflecting differences in the curve shapes (Appendix V).

4. Discussion

Here we present the first study with a tropical forest heliothermic lizard integrating previously obtained information on intraspecific phylogeographic structure with thermal biology traits and geographic distribution data to predict extinction risks due to climate change in a mega-diverse region. We also present an updated model of Sinervo et al. (2010) to calculate extinction risks based on highest body temperatures recorded in the field instead of preferred body temperatures estimated in laboratory, which can serve as a baseline for future studies with forest heliotherms.

While Sinervo et al. (2010) predicted an extinction risk for teilds of 13.6% by 2050 and 21% by 2080, our geographically more refined physiological models that included more habitat variables in the SDM for *K. calcarata* predicted higher relative extinction risks, with a mean across lineages of 53.7% for 2050 and 79.4% for 2070. A possible reason for the difference between the studies is that 30 out of the 36 heliothermic species Sinervo et al. (2010) considered for their

predictions inhabit open vegetation areas and deserts, and experience higher activity temperatures (mean body activity temperatures = 38.6° C, data from Sinervo et al., 2010) than the few inhabiting forest or aquatic environments (mean body activity temperatures = 34.8° C, data from Sinervo et al., 2010). However, even using other species, the model of Sinervo et al. (2010) should have had a greater relative extinction rate because they used a narrower thermal tolerance. Our modeling suggests that heliothermic lineages of *Kentropyx calcarata* living in densely forested environments (PA, AM, AP) have lower thermal preferences than the lineage living in the Amazon-Cerrado ecotone (Tocantins sub-region), and this elevates the risk of extinction. However, our interpretation of the differences between forested versus ecotone regions may also be because we included the effects of habitat change and not only thermal biology on extinction risk calculations.

In a similar approach, Kubisch et al. (2015) calculated the restriction hours for a Patagonian gecko thermoconformer (*Homonota darwinii*) and predicted that 20% of populations could become extinct by 2080, while Sinervo et al. (2010) predicted that the family as a whole would not suffer impacts due to climate change. As in our study, extinction-risk modeling by Kubisch et al. (2015) was conducted at the species level on a high latitude species of Phyllodactylid gecko, and thus, may represent a finer scale risk assessment.

Our SDM results suggest that ecophysiological traits, such as the thermal breadth for activity, H_r , H_a , and forest cover provide a useful explanation of the occurrence of *K. calcarata* in certain locations of the Amazon. In addition, the significant effect of precipitation (alone and in interaction with hours of restriction) suggests that a study of egg development and/or adult water loss (Muth, 1980), and the potential for climate change induced drought (Westphal et al., 2016), might prove to be a fruitful avenue of research at a deeper mechanistic level than we have done with our study of thermal ecophysiological traits across the four lineages of *K. calcarata*.

There was little relationship between mean or maximum operative temperatures and most of the thermal traits we measured. This might be expected, because the cost associated with different environments is mainly related to behavioral modifications that might not strongly impact fitness. However, passing the critical thermal maximum has a very strong immediate fitness cost (death) (Angilletta, 2009), because the lizard cannot escape from the immobilizing conditions, so it is not surprising that it is related to the maximum environmental temperature of the site.

Thermal traits appear to be, at least in part, determined by the environmental temperatures at each site, as evidenced by the strong relationship between critical thermal maximum and maximum operative temperatures. This indicates selection for different thermal traits at different sites (Gunderson and Leal, 2012; Moritz et al., 2012; Piantoni et al., 2016; Llewelyn et al., 2016). The variance in laboratory measures of thermal critical limits was greater in some sites than others, especially those sites that had low mean preferred temperatures. Therefore, adaptation to forested environments with lower environmental temperatures does not appear to have resulted in narrow amplitudes of thermal traits and loss of adaptive potential to climate changes. Huey et al. (2012) and Deutsch et al. (2008) predicted that presumed narrow amplitudes of thermal traits of tropical ectotherms would increase their vulnerability to climate change, but we show that individuals of Kentropyx calcarata, although living in thermally homogeneous environments, exhibit high variability in critical limits, which may favor individuals with increased tolerance to changes in temperature under a global-warming scenario.

When populations are polymorphic, variability tends to increase the niche amplitude in a way that thermal traits with greater variability may be favored by natural selection caused by climate change (Logan et al., 2014). Thus, we suggest that the potential adaptive amplitude will be greater in populations or areas whose individuals exhibit greater variation in thermal tolerance. Populations of *K. calcarata* in the Amazonian sites clearly show high flexibility in thermal critical limits

and, therefore, may have higher potential adaptive amplitude. In contrast, critical limits were less variable in the population at the ecotone. This could be a result of higher selection pressure at the ecotone, resulting in reduced variance in this trait. Despite the lower variability, under a scenario of climate warming, this could favor these populations as T_{pref} was higher at the ecotone. Further studies investigating the geographic and adaptive basis of genetic diversity variation and signals of selective pressure at the genomic level of *K. calcarata* are needed to test these hypotheses.

Most thermal biology studies consider the mean values of traits to evaluate differences between species, populations or locations (Hoffman and Sgro, 2011). We found geographic differences in thermal traits mean values, such as critical limits, but also in variability values. We suggest that future studies should pay closer attention to the variability of thermal traits. Variability in responses to environmental conditions can be considered a proxy for adaptive potentials of natural populations (e.g., Logan et al., 2014; Llewelyn et al., 2016) and, therefore, of persistence in response to climate change (Hoffman and Sgro, 2011).

Large areas of tropical forest have been predicted as potential refuges for biodiversity with increasing temperatures (Kearney et al., 2009, Clusella-Trullas et al., 2011). Our estimates are of increased extinction risks in most regions, mainly with the reduction of favorable areas for occurrence in lower-altitude Amazon forest regions that could contribute to local declines. However, even if the species goes extinct in some areas, colonization processes facilitated by large land connections that exist in the Amazon region may lead to species persistence in the long term.

As a forest heliotherm, persistence of *Kentropyx calcarata* also depends on forest preservation. Nonetheless, the Amazon rainforest is currently under high deforestation rates and some sub-regions are particularly impacted, such southern Pará state, which has the second-highest absolute deforestation rate of the Amazon (Fonseca et al., 2015). In addition to extinction risks estimated for the Tocantins sub-region, the Cerrado-Amazon ecotone largely coincides with the "Arc of Deforestation" of the Brazilian Amazon and is under intense anthropogenic change characterized by intense substitution of natural land-scapes for cattle ranching and agriculture (Costa and Pires, 2010). Combined effects of temperature increases on a global scale (Sinervo et al., 2010) and on a local scale (Lara-Resendiz et al., 2015) as a result of landscape conversion are factors that may further increase extinction risks of *K. calcarata* in these regions.

Our results show that Kentropyx calcarata thermal physiology varies geographically and thermal maxima appear to be conditional on the thermal environment of each surveyed locality. We used a novel approach to model extinctions risks for intraespecific lineages of a tropical heliothermic lizard that takes into account activity body temperatures measured in the field, vegetation cover and precipitation as environmental variables. Our results support the hypothesis that tropical lizard taxa are at higher risk of local extinction and this factor is geographically structured, with some regions within the species range predicted to become more or less favorable as the climatic conditions change over this century. Also, our laboratory experiments indicate significant differences in thermal traits between individuals from the same or distinct populations. We propose that this variability in thermal traits is the basis of local adaptation and might buffer some populations from extinction as the climate warms. Large areas of dense forest vegetation, thermal-tolerance variability and natural selection for heat tolerance are factors with considerable capacity to absorb the impacts of temperature increase on tropical ectotherms during climate change, and they should be included in future models of extinction risk.

Acknowledgements

We would like to thank Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) - FPW - 475559/2013-4; GRC -

456926/2014-3, Fundação de Amparo à Pesquisas do Estado do Amazonas-FAPEAM - FPW - 062.00665/2015 and 062.01110/2017 and Partnerships for Enhanced Engagement in Research from the US National Academy of Sciences and US Agency for International Development (PEER NAS/USAID PGA-2000005316) for funding this research. We thank Dra. Teresa Avila-Pires for support in data collection in Pará, the staff of INPA reserves for support in data collection in AM (Valdecira, Valdir and Rubenildo), and to all people who gave direct assistance for data collection in all locations: AM = G. Lima, L. Chrisley, J. Buckner, K. Bajer, O. R. Molnar, Sr. Compensa and Jesus; AP = A. Missassi, A. Nunes, J. Cosenza, and Sr. Maranhão: PA = C. Sette, A. Paula, Sr. Mór and Dorinha; and TO = C. Morais, S. Balbino, D. Tucker, M. Kucinick and S. Bee. FPW thanks financial support from L'Oréal-UNESCO-ABC Para Mulheres na Ciência - Brazil Werneck 2016 and L'Oréal-UNESCO For Women In Science awards - IRT Werneck 2017. GRC thanks financial suport from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) - GRC -88881.068430-2014-01 and Fundação de Apoio à Pesquisa do Distrito Federal (FAPDF) - Guarino Colli - 193.000.576/2009. B.S. and D. B. M. were supported by NSF EF-1241848. JCS thanks Jack W. Sites, Jr. (BYU) for his support as a postdoctoral fellow under NSF EF-1241885 and SJU for the start-up funds a new faculty. WEM was supported by the PPBio and CENBAM, and both contributed to the availability of a data repository. The authors have no conflict of interest to declare.

Authors' contributions

FPW, BS, DBM, WEM and GRC conceived the ideas and designed methodology; EPS, GHC, GRC, LMDV, JF and JCS collected the data; EPS, GHC, WEM and JCS analyzed the data; EPS led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Data accessibility

Data deposited in the KNB Data Repository. Identifier knb.1111.4 (https://knb.ecoinformatics.org/#view/knb.1111.4)

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at http://dx.doi.org/10.1016/j.jtherbio.2018.01.013.

References

- Addo-Bediako, A., Chown, S.L., Gaston, K.J., 2000. Thermal tolerance, climatic variability and latitude. Proc. R. Soc. Lond. B Biol. Sci. 267, 739–745.
- Adolph, S., 1990. Influence of behavioral thermoregulation on microhabitat use by two *Sceloporus* lizards. Ecology 7, 315–327.
- Angilletta Jr., M.J., 2009. Thermal Adaptation: a Theoretical and Empirical Synthesis. Oxford University Press, Oxford, UK.
- Araujo, M.B., Ferri-Yanez, F., Bozinovis, F., Marquet, P.A., Valladares, F., Chown, S.L., 2013. Heat freezes niche evolution. Ecol. Lett. 16, 1206–1219.
- Avila-Pires, T.C.S., Mulcahy, D.G., Werneck, F.P., Sites Jr., J.W., 2012. Phylogeography of the teiid lizard *Kentropyx calcarata* and the sphaerodactylid *Gonatodes humeralis* (Reptilia: squamata): Testing a geological scenario for the lower Amazon-Tocantins Basins, Amazonia, Brazil. Herpetologica 68, 272–287.
- Belasen, A., Brock, K., Li, B., Chremou, D., Valakos, E., Pafilis, P., Sinervo, B., Foufopoulos, J., 2016. Fine with heat, problems with water: microclimate alters water loss in a thermally adapted insular lizard. Oikos 126, 447–457.
- Bonino, M.F., Azócar, D.L.M., Schulte II, J.A., Abdala, C.S., Cruz, F.B., 2015. Thermal sensitivity of cold climate lizards and the importance of distributional ranges. Zoology 118, 281–290.
- Ceia-Hasse, A., Sinervo, B., Vicente, L., Periera, H.M., 2014. Integrating ecophysiological models into species distribution projections of European reptile range shifts in response to climate change. Ecography 7, 679–688.
- Clobert, J., Oppliger, A., Sorci, G., Ernande, B., Swallow, J.G., Garland, T., 2000. Tradeoffs in phenotypic traits: endurance at birth, growth, survival, predation and susceptibility to parasitism in a lizard. Lacerta vivipara. Funct. Ecol. 14, 675–684.
- Clusella-Trullas, S., Blackburn, T.M., Chown, S.L., 2011. Climatic predictors of temperature performance curve parameters in ectotherms imply complex responses to climate change. Am. Nat. 177, 738–751.

- Colwell, R.K., Brehm, G., Cardelus, C.L., Gilman, A.C., Longino, J.T., 2008. Global warming, elevational range shifts, and lowland biotic attrition in the wet tropics. Science 322, 258–261.
- Costa, M.H., Pires, G.F., 2010. Effects of Amazon and Central Brazil deforestation scenarios on the duration of the dry season in the arc of deforestation. Int. J. Climatol. 30, 1970–1979.
- Cronemberger, A.A., 2015. Filogeografia multi-locus de *Kentropyx calcarata* Spix 1825 (Reptilia: teiidae) na Amazônia Oriental (Master's thesis). Museu Paraense Emilio Goeldi, Pará.
- Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C., Martin, P.R., 2008. Impacts of climate warming on terrestrial ectotherms across latitude. Proc. Natl. Acad. Sci. USA 105, 6668–6672.
- Dillon, M.E., Wang, G., Huey, R.B., 2010. Global metabolic impacts of recent climate warming. Nature 467, 704–707.
- Fonseca, A., Justino, M., Souza Jr., C., Veríssimo, A., 2015. Deforestation report for the Brazilian Amazon (November) SAD (p. 10). Belém: Imazon.
- Gunderson, A.R., Leal, M., 2012. Geographic variation in vulnerability to climate warming in a tropical Caribbean lizard. Funct. Ecol. 26, 783–793.
- Hoffman, A.A., Sgro, C.M., 2011. Climate change and evolutionary adaptation. Nature 470, 479–485.
- Huey, R.B., Stevenson, R.D., 1979. Integrating thermal physiology and ecology of ectotherms: a discussion of approaches. Am. Zool. 19, 357–366.
- Huey, R.B., Deutsch, C.A., Tewksbury, J.J., Vitt, L.J., Hertz, P.E., Alvarez-Perez, H.J., Garland Jr, T., 2009. Why tropical forest lizards are vulnerable to climate warming. Proc. R. Soc. Lond. B Biol. Sci. 276, 1939–1948.
- Huey, R.B., Kearney, M.R., Krockenberger, A., Holtum, J.A., Jess, M., Williams, S.E., 2012. Predicting organismal vulnerability to climate warming: roles of behaviour, physiology and adaptation. Philos. Trans. R. Soc. B. 367, 1665–1679.
- IPCC, 2014. Climate Change 2014. Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge.
- INMET, 2016. Instituto Nacional de Meteorologia. http://www.inmet.gov.br/portal/ (Accessed 1 February 2016).
- Kearney, M., Porter, W., 2009. Mechanistic niche modelling: combining physiological and spatial data to predict species' ranges. Ecol. Lett. 12, 334–350.
- Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G., Jarvis, A., 2005. Very high resolution interpolated climate surfaces for global land areas. Int. J. Clim. 25, 1965–1978.
- Kirchhof, S., Hetem, R.S., Lease, H.M., Miles, D.B., Mitchell, D., Muller, J., Rodel, M.O., Sinervo, B., Wassenaar, T., Murray, I.W., 2017. Thermoregulatory behavior and high thermal preference buffer impact of climate change in a Namib Desert lizard. Ecosphere 8 (12), e02033. http://dx.doi.org/10.1002/ecs2.2033.
- Kubisch, E.L., Corbalán, V., Ibargüengoytía, N.R., Sinervo, B., 2015. Local extinction risk of three species of lizard from Patagonia as a result of global warming. Can. J. Zool. 94, 49–59.
- Lara-Reséndiz, R.A., Jezkova, T., Rosen, P.C., Méndez-de la Cruz, F.R., 2014. Thermoregulation during the summer season in the Goode's horned lizard *Phrynosoma goodei* (Iguania: Phrynosomatidae) in Sonoran Desert. Amphib.-Reptil. 35, 161–172.
- Lara-Resendiz, R.A., Gadsden, H., Rosen, P.C., Sinervo, B., Méndez-de-la-Cruz, F., 2015. Thermoregulation of two sympatric species of horned lizards in the Chihuahuan Desert and their local extinction risk. J. Therm. Biol. 48, 1–10.
- Llewelyn, J., Macdonald, S.L., Hatcher, A., Moritz, C., Phillips, B.L., 2016. Intraspecific variation in climate-relevant traits in a tropical rainforest lizard. Divers. Distrib. 1–13. http://dx.doi.org/10.1111/ddi.12466.
- Logan, M., Cox, R.M., Calsbeek, R., 2014. Natural selection on thermal performance in a novel thermal environment. Proc. Natl. Acad. Sci. USA 39, 14165–14169.
- Losos, J.B., 2008. Phylogenetic niche conservatism, phylogenetic signal, and the relationship between phylogenetic relatedness and ecological similarity among species. Ecol. Lett. 11, 995–1007.
- Marimon, B.S., De, S. Lima, Duarte, E., Chieregatto, T.G., Ratter, J.A, L.C., 2006. Observations on the vegetation of northeastern Mato Grosso, Brazil. IV. An analysis of the Cerrado-Amazonian Forest ecotone. Edinb. J. Bot. 63, 323–341.
- Moritz, C., Langham, G., Kearney, M., Krockenberger, A., VanDerWal, J., Williams, S., 2012. Integrating phylogeography and physiology reveals divergence of thermal traits between central and peripheral lineages of tropical rainforest lizards. Philos. Trans. R. Soc. Lond. B Biol. Sci. 367, 1680–1687.
- Muth, A., 1980. Physiological ecology of desert iguana (*Dipsosaurus dorsalis*) eggs: temperature and water relations. Ecology 61, 1335–1343.
- Pearman, P.B., Guisan, A., Broennimann, O., Rand, C.F., 2008. Niche dynamics in space and time. Trends Ecol. Evol. 23, 149–158.
- Piantoni, C., Navas, C.A., Ibargüengoytía, N.R., 2016. Vulnerability to climate warming of four genera of New World iguanians based on their thermal ecology. Anim. Conserv. 19, 1469–1795.
- R Development Core Team, 2014. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna.
- Robson, M.A., Miles, D.B., 2000. Locomotor performance and dominance in male tree lizards Urosaurus ornatus. Funct. Ecol. 3, 338–344.
- Sartorius, S.S., Vitt, L.J., Colli, G.R., 1999. Use of natural and anthropogenically disturbed habitats in Amazonian rainforest by the teiid lizard Ameiva ameiva. Biol. Conserv. 90, 91–101.
- Sinervo, B., Mendez-de-la-Cruz, F., Miles, D.B., Heulin, B., Bastiaans, E., Villagran-Santa Cruz, M., Lara-Resendiz, R., Martınez-Mendez, N., Calderon-Espinosa, M.L., Meza-Lazaro, R.N., Gadsden, H., Avila, L.J., Morando, M., de la Riva, I.J., Sepulveda, P.V., Rocha, C.F.D., Ibargüengoytía, N., Puntriano, C.A., Massot, M., Lepetz, V., Oksanen,

E. Pontes-da-Silva et al.

T.A., Chapple, D.G., Bauer, A.M., Branch, W.R., Clobert, J., Sites Jr, J.W., 2010. Erosion of lizard diversity by climate change and altered thermal niches. Science 328, 894–899.

- Sunday, J.M., Bates, A.E., Dulvy, N.K., 2011. Global analysis of thermal tolerance and latitude in ectotherms. Philos. Trans. R. Soc. B. 278, 1823–1830.
- Thuiller, W., Lavorel, S., Araújo, M.B., Sykes, M.T., Prentice, I.C., 2005. Climate change threats to plant diversity in Europe. Proc. Natl. Acad. Sci. USA 102, 8245–8250.
- Van Damme, R., Bauwens, D., Verheyen, R.F., 1986. Selected body temperatures in lizard *Lacerta vivipara*: variation within and between populations. J. Therm. Biol. 11, 219–222.
- Vitt, L.J., 1991. Ecology and life-history of the widely foraging lizard *Kentropyx calcarata* (Teiidae) in Amazonian Brazil. Can. J. Zool. 69, 2791–2799.
- Vitt, L.J., Zani, P.A., Marinho-Lima, A.C., 1997. Heliotherms in tropical rain forest: the ecology of *Kentropyx calcarata* (Teiidae) and *Mabuya nigropunctata* (Scincidae) in the Curuá-Una of Brazil. J. Trop. Ecol. 13, 199–220.
- Werneck, F.P., Giugliano, L.G., Collevatti, G.G., Colli, G.R., 2009. Phylogeny, biogeography and evolution of clutch size in South American lizards of the genus *Kentropyx* (Squamata: teiidae). Mol. Ecol. 18, 262–278.
- Westphal, M.F., Stewart, J.A.E., Tennant, E.N., Butterfield, H.S., Sinervo, B., 2016.
- Contemporary drought and future effects of climate change on the endangered bluntnosed Leopard Lizard *Gambelia sila*. PLoS One. 11, e0154838. http://dx.doi.org/10. 1371/journal.pone.0154838.
- Wickham, H., 2009. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York.
- Winter, M., Fiedler, W., Hochachka, W.M., Koehncke, A., Meiri, S., De la Riva, I., 2016. Patterns and biases in climate change research on amphibians and reptiles: a systematic review. R. Soc. Open Sci. 3, 160158. http://dx.doi.org/10.1098/rsos.160158.
- Wood, S.N., 2006. Generalized Additive Models: an Introduction with R. Chapman and Hall/CRC.



Emerson Pontes-da-Silva is an Master in Ecology by the Ecology Program of the Instituto Nacional de Pesquisas da Amazônia. He received his B.S. in Biology at the Federal Institute of Education, Science and Technology of Amazonas. He is interested in the impacts of the global warming on thermal physiology, ecology and evolution of Amazonian lizards and amphibians.



Gabriel Henrique de O. Caetano is graduate in Biological Sciences (2011) and Master in Ecology by University of Brasília (2014). Currently attending the Ecology and Evolutionary Biology Ph. D. program at University of California Santa Cruz. Main interests are population ecology, ecophysiology, climate change and conservation with emphasis in herpetology.



Donald B. Miles Professor (Ohio University) and a Research Associate of California Academy of Sciences. He is an evolutionary biologist who conducts research on Ecology, Morphology and Biotics impacts of climate change.



Guarino R. Colli is an Associate Professor and Curator of Herpetology at Universidade de Brasília, and an Affiliate Research Associate at the Sam Noble Oklahoma Museum of Natural History. His research focuses on the ecology, evolution and conservation of the Cerrado herpetofauna.



William E. Magnusson is a researcher of the Instituto Nacional de Pesquisas da Amazônia where he studies the ecology of tropical systems, and principally techniques of biodiversity monitoring. He is a member of the Brazilian Academy of Sciences and obtained his B.Sc. and Ph.D. in Biological Sciences from Sydney University.



Luisa M. Diele-Viegas is a PhD student at the Ecology and Evolution Graduate Program in State University of Rio de Janeiro, Rio de Janeiro, Brazil. She received her B. S. in Zoology at the Federal University of Rio de Janeiro, and her M.S. in Zoology at the Federal University of Pará, in association with the Museu Paraense Emílio Goeldi. Her primary research interests includes the influence of the global warming on thermal physiology, ecology and distribution of Amazonian lizards, and the definition of priority conservation areas for those species.



Barry Sinervo Full Professor, University of California, Santa Cruz, is an evolutionary biologist who conducts research on Behavioral Ecology, Game Theory and Biotic Impacts of Climate Change. He is currently researching contemporary extinctions of reptiles and amphibians and changes in plant communities driven by climate change, leading a multinational research team of scientists developing physiological models of biotic impacts of climate on diverse biological systems from equatorial sites to polar regions. He is also Director of the UC-wide Institute for the Study of the Ecological and Evolutionary Climate Impacts, a research consortium funded by a UC Presidential Research Catalyst Award.



Jéssica Fenker is a PhD student at Australian National University (ANU), Canberra, Australia. She received her M.S. in Animal Biology at the Universidade de Brasília (UnB), Brasília-DF, Brazil. Her interests includes biogeography, ecology and evolution of Squamates, with special focus on savanna environments.



Juan C. Santos is an Assistant Professor at St. John's University (New York, USA) and was previously affiliated to Brigham Young University (Utah, USA). He received his B.A. in Biology at the Pontificia Universidad Católica del Ecuador, M.S. in Statistics and Ph.D. in Ecology, Evolution and Behavior at the University of Texas at Austin. He is interested in the evolution of complex phenotypes, multivariate comparative methods, ecophysiological models of species distributions, and genotype-phenotype relationships using genomic/transcriptomic data. His biological systems of research include poison frogs (Dendrobatidae) and New World lizards.



Fernanda P. Werneck is a Researcher and Herpetology Curator at Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Brazil. She received here B.S. in Biology and her M.S. in Ecology at the Universidade de Brasília (UnB), Brasília-DF, Brazil, and her Ph.D in Integrative Biology at the Brigham Young University (BYU), Provo-Utah, United States. Her main research lines integrate ecological and evolutionary approaches to investigate the processes responsible by patterns of diversity, genetic variation, and diversification of Neotropical amphibians and reptiles, with a special focus on the evolutionary history and future of herpetofauna at the Amazonia-Cerrado transition.